





HOMŒOPATHIC

DOMESTIC PHYSICIAN;

CONTAINING

THE TREATMENT OF DISEASES,

WITH POPULAR EXPLANATIONS ON ANATOMY, PHYSIOLOGY,  
HYGIENE, AND HYDROPATHY,

ALSO

AN ABRIDGED MATERIA MEDICA.


BY J. H. PULTE, M. D.

CINCINNATI:

H. W. DERBY & CO., AND H. F. DAVIS.

NEW YORK: A. S. BARNES & CO.


1850.



WBK  
P983h  
1850

Feb 11 1850  
Derby

Entered according to act of Congress, in the year of our Lord, 1850,  
BY H. W. DERBY & Co.,  
in the Clerk's Office of the District Court for the District of Ohio.



CINCINNATI:  
*Wright, Ferris & Co., Printers.*

## P R E F A C E .

---

OF late years a great deal of attention has been bestowed upon the introduction and perfection of popular works on the treatment of diseases, by the medical profession, not merely by Homœopathists, but, also, by practitioners of other medical schools. This shows, evidently, that the hitherto backward world of medical science has been caught, at last, with the proper spirit of modern progress, whose distinguishing feature in scientific matters consists, undoubtedly, in the tendency to popularize abstruse sciences, and make them useful and accessible to the many.

Homœopathy has not been in the rear as regards these popular instructions, and many valuable volumes have been issued from the press within a short time, which facilitated in a great degree the domestic use of the homœopathic medicines. This service, rendered by the profession to the public in a liberal and generous spirit, has been amply rewarded

by the latter, in extending all over the country, more rapidly the blessings of our beloved healing art, and establishing its title, as a benefactor of mankind, more firmly in the hearts of the people.

The author well recollects that the first "domestic physicians" which appeared, were looked upon with distrust by some of the profession. But the people in general hailed them as welcome friends in their domestic afflictions, where counsel and advice is so often needed at times, when the medical aid can not be obtained. Imperfect as the first efforts of this kind naturally must have been, their beneficial results for families, and for the spread of Homœopathy, were such that the improvement of these necessary medical guides was rapidly undertaken and accomplished. All honor and praise, however, must be given to those few in Germany and America, who made the first attempts to popularize the practice of Homœopathy.

At this time scarcely one homœopathic family is without one or more books on domestic practice, which is a sufficient evidence of their practical utility.

Thus have these messengers of mercy and usefulness travelled with, and often ahead of, the regular practitioners of Homœopathy—the silent, but efficient missionaries of truth, declaring it everywhere by facts and conquests over disease, won by the people

themselves. The profession in general seems now to regard them as necessary allies in the great work of reforming the medical state of the world, and bestows a great deal of care on their constantly increasing perfection, by making them more practical and definite, progressing in their improvement as the science itself progresses.

In this spirit the present volume is written and given to the public by the author, who has attempted to embody in it the results of a practice of more than sixteen years.

In the pathology of the work he has followed mostly the views of Hufeland, who is generally practically useful, and of Schænlein, who is precise and scientific.

In the therapeutics he has endeavored to consult the best authors of our school, carefully comparing their views with his own results, which are embodied in this work.

To increase its general usefulness, popular treatises on Anatomy, Physiology, Hygiene, and Hydropathy, have been added, branches and parts of medical science as a whole, which, when properly understood and digested by the people, will in a great degree aid in fostering the salutary results expected from the use of a domestic physician. Besides this, a condensed extract of the Homœopathic Materia Medica, given at the close of the book, enables the reader to

## PREFACE.

---

correct his prerscriptions, and enlarge his general knowledge of the homœopathic science. Thus it is hoped that, by these additions, the work is rendered more perfect in a twofold manner: first, by giving a more extended knowledge of the human system and its laws through the treatises on Anatomy and Physiology ; and, secondly, by making the reader acquainted with those practical branches of medicine, Hygiene and Hydropathy, of which, particularly, the latter has, within the last few years, very much engrossed the attention of the profession and of invalids. In this respect the author himself does not hesitate to declare that he considers Hydropathy, applied within her proper limits, the handmaid of Homœopathy, and that these two, combined in this relationship, are destined to conquer the medical world.

THE AUTHOR.

*Cincinnati, October, 1850.*

## TABLE OF CONTENTS.

---

LIST OF MEDICINES.....	xiii
INTRODUCTION.....	xvii
DIETETIC RULES.....	xxi

### PART I.

#### DESCRIPTION AND TREATMENT OF DISEASES.

##### CHAPTER I.

###### *General Diseases.*

Rheumatism—Acute—Chronic. Gout. Pains in the small of the Back and Loins. Pain in the Hip. Hip-disease. Inflammation of the Knee-joint. Inflammation of the Psoas-muscle. Cramp in the Legs. Sweating Feet. Goitre. Sleeplessness. Nightmare. Palsy. Delirium Tremens. Epilepsy. St. Vitus' Dance. Tetanus. Trismus. Lockjaw. Somnolency. Lethargy. Fatigue. Overheating and Mental Exhaustion. Fainting. Swooning. Apparent Death from Hunger—from Drowning—from Freezing—Lightning—Hanging—Choking—from Noxious vapors—Fall or Blow—Violent Mental Emotion..... 3

##### CHAPTER II.

###### *Casual Diseases.*

I. External Injuries. Fractures of Bones. Dislocations of Joints. Wounds and Bruises. Sprains and Concussions. Burns and Scalds. Poisoned Wounds—Bites and Stings of Insects. Frozen Limbs.

II. Poisons—Antidotes. Animal Poisons. Vegetable Poisons. Mineral Poisons.

III. Medicinal Diseases. List of Drugs causing Diseases, and their Remedies..... 30



## CHAPTER III.

*Cutaneous Diseases.*

1. Eruptive Fevers. Rash. Nettle-Rash—Hives. Erysipelas. St. Anthony's Fire. Measles. French Measles. Scarlet Rash. Scarlet Fever. Chicken Pox. Small Pox. Varioloid.

2. Chronic Eruptions. Irritation of the Skin—Itching—Pruritus. Chilblains. Excoriations—Intertrigo. Ringworm—Herpes Circinnatus. Tetter—Herpes Itch—Scabies. Milk-crust—Crusta Laetea. Scald Head—Tinea Capitis. Corns—Indurations of the Skin. Warts—Verrucae. Whitlows. Felons. Panaris. Ulcers. Abscesses. Tumors. Swelling of glands. Boils—malignant Boils—Carbuncles. Seirrhus. Cancer..... 47

## CHAPTER IV.

*Fevers.*

Explanation of the term Fever. Irritative, Inflammatory, Torpid form of Fever. Common or Ephemeral Fever. General Inflammatory Fever. Nervous or Typhoid Fever. Typhus—Different forms of Typhus. Yellow Fever. Intermittent Fever—Fever and Ague—Chills and Fever—Febris Intermittens..... 84

## CHAPTER V.

*Affections of the Mind.*

Sudden Emotions. Hypochondria. Melancholia..... 110

## CHAPTER VI.

*Affections of the Head.*

1. Congestion or determination of Blood to the Head—Congestion ad Caput. 2. Vertigo. Giddiness or Dizziness. 3. Weakness of Memory. 4. Apoplexy. 5. Inflammation of the Brain. 6. Sunstroke. 7. Headache. *a.* Headache from Congestion of Blood to the Head. *b.* Headache from Catarrh and Cold in the Head. *c.* Headache from Rheumatism. *d.* Headache from Constipation and Gastric Derangement. *e.* Headache from External Causes. Sick-Headache. Nervous Headache—Neuralgia in the Head—Megrim. Falling off of the Hair—Allopecia. .... 115

## CHAPTER VII.

*Affections of the Eyes.*

General Remarks. 1. Inflammation of the Eyelid and its Margins. *a.* Inflammation and Swelling of the Eyelid. *b.* Inflammation of the Margins of the Lids. 2. Inflammation of Eyeball—Ophthalmia. 3. Chronic Inflammation of the Eyes. *a.* From Gout and Rheumatism.

<i>b.</i> From Scrofula, Scrofulous Sore Eyes. <i>c.</i> From Syphilis or Venereal Disease. <i>d.</i> From Abuse of Mercury. 4. Stye on the Eyelid. 5. Weeping or Watery Eyes. 6. Cataract—Glaucoma. 7. Weakness of Sight—Amblyopia—Blindness—Sudden and Incipient—Anaurosis. 8. Short Sightedness—Myopia. 9. Far-Sightedness. 10. Squinting—Strabismus. 11. Falling of the Eyelids—Paralysis. 12. Foreign Substances in the Eye. ....	140
--	-----

## CHAPTER VIII.

*Affections of the Ears.*

1. External Ear and its surrounding parts. <i>a.</i> Eruptions and Swelling of the External Ear. <i>b.</i> Mumps—Parotitis. 2. Internal Ear. <i>a.</i> Inflammation of the Ear—Otitis. <i>b.</i> Earache—Otalgia. <i>c.</i> Humming, Buzzing of the Ears. <i>d.</i> Hardness of Hearing—Deafness. <i>e.</i> Running of the Ears—Otorrhœa. <i>f.</i> Polypus of the Ear. <i>g.</i> Foreign Substances in the Ear, .....	152
--	-----

## CHAPTER IX.

*Affections of the Nose.*

1. Inflammation or Swelling of the Nose. 2. Eruptions on the Nose—Itching of the Nose. <i>a.</i> Eruptions on the Nose. <i>b.</i> Itching of the Nose. 3. Bleeding of the Nose—Epistaxis. 4. Cold in the Head—Coryza—Suppression of the Catarrh or Cold in the Head. 5. Ulceration in the Nose—Catarrh—Ozaena Narium. <i>a.</i> Ozaena Narium—Catarrh. <i>b.</i> Ozaena Antri Highmori. 6. Polypus of the Nose. Cancer of the Nose. 7. Foreign Substances in the Nose.....	163
--	-----

## CHAPTER X.

*Affections of the Face, Lips, and Jaws.*

1. Inflammation and Swelling of the Face. <i>a.</i> Erysipelas of Face and Lips. <i>b.</i> Swelling of the Cheek. 2. Eruptions of the Face. 3. Faceache—Neuralgia of the Face—Prosopalgia—Tic Douloureux. 4. Warts and Spots on the Face—Cancer of the Face. 5. Paralysis of the Face and Jaw. 6. Lockjaw.....	174
--	-----

## CHAPTER XI.

*Affections of the Teeth, Gums, and Mouth.*

1. Toothache—Odontalgia. 2. Caries of the Teeth—Carious Teeth. 3. Offensive Breath. 4. Bad Taste in the Mouth. 5. Inflammation and Swelling of the Tongue—Glossitis. 6. Canker of the Mouth—Stomacace. 7. Inflammation of the Palate. 8. Abscess in the Gums—Gumboils—	
--	--

Swelling of the Jawbone. 9. Scurvy—Scorbutus. 10. Salivation—Ptyalism. 11. Paralysis of the Tongue. 12. Difficulty of Speech—Stammering. 13. Ranula or Swelling under the Tongue.....	180
---	-----

## CHAPTER XII.

*Affections of the Throat.*

Sore Throat—Quinsy—Angina Faucium. <i>a.</i> Common Sore Throat. <i>b.</i> Ulcerated and Putrid Sore Throat—Malignant Quinsy—Angina Gangrenosa. Inflammation and Swelling of the Tonsils—Amygdalitis. Swelling or Elongation of the Palate.....	199
---	-----

## CHAPTER XIII.

*Affections of the Windpipe and Chest.*

1. Hoarseness—Raucitas—Aphonia. <i>a.</i> Acute Hoarseness. <i>b.</i> Chronic Hoarseness. 2. Ulceration of the Windpipe—Bronchitis. 3. Croup and Croupy Cough. 4. Influenza. 5. Cough—Hooping-Cough—Tussis Convulsiva. Inflammation of the Lungs—Pneumonia. Pneumonia in Old People—Pneumonia Notha. Remarks on Pneumonia. Pleurisy. <i>a.</i> Real Pleurisy. <i>b.</i> False Pleurisy—Stitch in the Side. Diseases of the Heart. <i>a.</i> Inflammation of the Heart—Corditis. <i>b.</i> Palpitation of the Heart. Congestion of the Chest—Plethora Pectoris. Hemorrhage of the Lungs—Hæmoptysis. Asthma. Asthma of Millar. Dropsy in the Chest—Hydrothorax. Pulmonary Consumption—Phthisis Pulmonalis. Contusion of the Chest by a Fall or Blow.....	210
--	-----

## CHAPTER XIV.

*Affections of the Stomach and Bowels.*

1. Want of Appetite—Anorexia. 2. Morbid Appetite—Bulimy. 3. Dyspepsia—Weak Stomach. 4. Waterbrash—Heartburn—Indigestion—Flatulency—Sour Stomach—Pyrosis. 5. Biliousness—Gastric Derangement. Nausea—Vomiting. Sea-Sickness. Spasms and Pain in the Stomach—Cardialgia—Neuralgia of the Stomach. Gripping Colic—Enteralgia—Stomach-ache. Congestion and Stagnation of Blood in the Abdomen. Inflammation of the Diaphragm—Diaphragmitis. Abdominal Inflammation. 1. Inflammation of the Stomach—Gastritis. 2. Inflammation of the Bowels—Enteritis. Inflammation of the Liver—Hepatitis. Jaundice. Tetanus. Inflammation of the Spleen—Splenitis. Worms. Diseases caused by the Presence of Worms. Constipation—Costiveness. Diarrhea—Looseness of the Bowels. Dysentery—Bloody Flux. Cholera
--

Morbus. Asiatic Cholera—Cholera Asiatica. Piles—Hemorrhoids. Itching of the Anus. Prolapsus Ani. Rupture—Hernia. Dropsy of the Abdomen—Ascites. External Injuries of the Abdomen.....	257
---	-----

## CHAPTER XV.

*Affections of the Urinary and Genital Organs.*

Inflammation of the Kidneys—Nephritis. Inflammation of the Bladder—Cystitis. Strangury. Dysury. Ischury. Anury. Incontinence of Urine—Enuresis. Diabetes. Piles of the Bladder. Polypus of the Bladder. Gravel and Stone in the Bladder—Calculus. Hemorrhage with the Urine—Hæmaturia—Mictus Cruentus. Affections of the Penis. Affections of the Testicles. Seminal Weakness—Onania. Eruptions on the Genital Organs. Syphilis. Gonorrhea. <i>a.</i> Syphilis—Chancre. <i>b.</i> Gonorrhea—Clap—Gleet.....	340
---	-----

## CHAPTER XVI.

*Diseases of Females.*

1. Menstruation. Obstructions of the First Menses. Chlorosis. Suppression of the Menses. Painful Menstruation. Menstruation too Early. Menstruation too Late and too Scanty. Menstruation too Copious—Flooding. Menstruation of too long Duration. Critical Period. Cessation of the Menses—Change of Life. Prolapsus Uteri—Falling of the Womb. Leucorrhœa—Fluor Albus—Whites. Pregnancy. Diseases during Pregnancy. Nausea and Vomiting. Pruritus—Itching. Varicose Veins. Spots on the Face. Miscarriage—Abortion. Premature Birth. Regular Birth. Labor-Pains. Adherence of the After-birth—Placenta. Treatment after Delivery. After-pains. Duration of Confinement. Breast—Nipples—Secretion of Milk. Sore Nipples. Ague in the Breast. Gathered Breast. Deficiency of Milk. Suppression of Milk. Milk Bad—too Thin, or Repugnant to the Child. Milk—Excessive flow of it. Weaning. State of the Bowels. Sore Mouth of Nursing Women. Discharge from the Womb during Confinement—Lochia. Pain—Inflammation in the Abdomen—Metritis. Puerperal Fever. General Complaints during Confinement.....	353
---	-----

## CHAPTER XVII.

*Treatment of Children.*

Apparent Death of a New-born Infant—Asphyxia. Diet of a New-born Infant. Colic—Crying of Infants. Elongation of the Head. Snuffles—Obstruction of the Nose—Coryza. Swelling of the Breasts in Infants. Restlessness—Sleeplessness. Inflammation of the Eyes. Thrush, or Sore
--

Month of Infants—Aphthæ. Heat—Red Gum—Heat Spots. Excoriations—Rawsness of the Skin. Jaundice. Erysipelas. Convulsions or Spasms. Teething—Dentition. Constipation. Diarrhea. Fevers. Dropsy of the Brain—Hydrocephalus. Asthma—Choking Fits—Asthma Millari. Remittent Fever of Infants. Vaccination. Cholera Infantum. Summer Complaint.....	381
---	-----

## PART II.

### I. ANATOMY AND PHYSIOLOGY.

Structure of the Human Organism. Elements of the Human Organism. Anatomy of the Bones. Physiology of the Bones. The Muscles. The Teeth. Digestive Organs. The Urinary System. The Respiratory and Circulatory Organs. The Nervous System. The Skin.....	397
---	-----

### II. HYGIENE AND HYDROPATHY.

Hygiene. Food and Drink. Sleep. Air and Exercise. Clothing and Occupation. Hydropathy. Diseases in which to apply it. Different Modes of applying the Cold Water. 1. Ablutions. 2. Shower-Baths. 3. Partial or Half-Baths. 4. Full-Baths. 5. General Plunging-Baths. 6. Local Baths. 7. Head-Baths. 8. Sitting-Baths—Sitz-Baths. 9. Foot-Baths. 10. The Drop-Bath. 11. The Douche. 12. Local Applications (Umschläge) Wet Bandages. 13. Applications round one-half the Body. Wet Linen Sheets. 14. Sweating.....	428
---	-----

III. MATERIA MEDICA.....	475
--------------------------	-----

GLOSSARY.....	541
---------------	-----

INDEX.....	549
------------	-----

# LIST OF MEDICINES,

## THEIR SYNONYMES AND ANTIDOTES.

1. ACONITE. Aconitum Napellus (plant.) *Antidotes*—Camphor, Nux vom., Wine, Vinegar.
2. ANTIMONIUM CRUM. Antimony (mineral.) *Antidotes*—Hepar sulph., Mercury, Pulsatilla.
3. ARNICA. Arnica Montana (plant.) *Antidotes*—Camphor, Ignatia, Ipecac.
4. ARSENIC. Arsenicum album (mineral.) *Antidotes*—For poisoning with it: rust of iron. For its dynamic effects: China, Hepar sulph., Ipecac., Nux vomica, Veratrum.
5. AURUM. Aurum metallicum. Gold. (Metal.) *Antidotes*—Belladonna, China, Cuprum, Mercury.
6. BELLADONNA. Deadly nightshade (plant.) *Antidotes*—Coffea, Hyosciamus, Hepar sulph., Pulsatilla.
7. BELLADONNA CC. Is the two hundredth potency of Belladonna.
8. BRYONIA. Bryonia alba (plant.) *Antidotes*—Aconite, Chamomile, Ignatia, Nux vomica.
9. CALCAREA CARBONICA. Carbonate of lime. *Antidotes*—Camphor, Nitric acid, Sulphur.
10. CAMPHORA. Camphor. *Antidotes*—Opium, Nitri spiritus.
11. CANTHARIS. Spanish fly (animal.) *Antidote*—Camphor.
12. CAPSICUM. Spanish pepper (vegetable.) *Antidote*—Camphor.
13. CARBO VEG. Carbo vegetabilis. Wood charcoal. *Antidotes*—Arsenic, Camphor, Coffea, Lachesis.
14. CAUSTICUM. Caustic of the alkalies. *Antidotes*—Coffea, Colocynth, Nux vomica.
15. CHAMOMILE. Chamomilla (plant.) *Antidotes*—Aconite, Cocculus, Coffea, Ignatia, Nux vomica, Pulsatilla.
16. CHINA. Cinchona. Peruvian bark (vegetable.) *Antidotes*—Arnica, Arsenic, Belladonna. Calcareo carbonica, Carbo veg., Ipecac., Sulphur.
17. CINA. Wormseed (vegetable.) *Antidotes*—Ipecac., Veratrum.
18. CINA CC. Is the two hundredth potency of Cina.
19. COCCULUS. Indian Berries. Coccle (vegetable.) *Antidotes*—Camphor, Nux vomica.
20. COFFEA. Coffee Berries (vegetable.) *Antidotes*—Aconite, Chamomile, Nux vomica.
21. COFFEA CC. is the two hundredth potency of Coffea.

22. COLCHICUM. Meadow Saffron (plant.) *Antidotes*—Nux vomica, Coccus, Pulsatilla.
23. COLOCYNTHIS. Colocynth (vegetable.) *Antidotes*—Camphor, Causticum, Coffea, Chamomile.
24. CONIUM. Hemlock (vegetable.) *Antidotes*—Coffea, Spiritus Nitri.
25. CROCI. Saffron (vegetable.) *Antidote*—Opium.
26. CUPRUM. Cuprum metallicum. Copper (metal.) *Antidotes*—Belladonna, China, Ipecac., Mercury, Nux vomica.
27. DROSER. Round-leaved Sun Dew (vegetable.) *Antidotes*—Camphor.
28. DULCAMARA. Bitter Sweet. Woody Nightshade (plant.) *Antidotes*—Camphor, Ipecac., Mercurius.
29. EUPHRASIA. Eye-bright (plant.) *Antidote*—Pulsatilla.
30. FERRUM. Ferrum metallicum. Iron (metal.) *Antidotes*—Arnica, Arsenic, Belladonna, Ipecac., Mercury, Pulsatilla.
31. GRAPHITES. Plumbago. Pure Black Lead (metal.) *Antidotes*—Arsenic, Nux vomica, Zinc.
32. HELLEBORUS (NIGER.) Black Hellebor (plant.) *Antidotes*—Camphor, China.
33. HEPAR SULPHURIS. Lime of Sulphur. Sulphuret of Lime. *Antidotes*—Vinegar, Belladonna.
34. HYOSCIAMUS (NIGER.) Henbane (plant.) *Antidotes*—Belladonna, Camphor, China.
35. IGNATIA. St. Ignatius' Bean (vegetable.) *Antidotes*—Pulsatilla, Chamomile, Coccus, Arnica, Camphor, Vinegar.
36. IODINE. Iodum. *Antidotes*—Arsenic, Camphor, Coffea, Phosphorus, Sulphur.
37. IPECAC. Ipecacuanha (vegetable.) *Antidotes*—Arnica, Arsenic, China.
38. JALAPPA. Jalap (vegetable.) *Antidotes*—Camphor.
39. KALI HYDRIOLICUM. Hydriodate of Potassa (mineral.)
40. LACHESIS. Poison of the Lance-headed Serpent (animal.) *Antidotes*—Arsenic, Belladonna, Nux vomica, Rhus tox.
41. LYCOPodium. Club Moss. Wolf's Claw (vegetable.) *Antidotes*—Camphor, Pulsatilla.
42. MERCURIUS (VIVUS.) Quicksilver (mineral.) *Antidotes*—Arnica, Belladonna, Camphor, Hepar sulph., Iodine, Lachesis, Sulphur.
43. MURIATIC ACID. Muriatis acidum (mineral.) *Antidotes*—Camphor, Bryonia.
44. NATRUM MURIATICUM. Muriate of Soda (mineral.) *Antidotes*—Arsenic, Camphor, Nitri spiritus.
45. NITRI ACIDUM. Nitric acid (a mineral acid.) *Antidotes*—Calcarea carb., Conium, Camphor, Hepar sulph., Sulphur.
46. NUX VOMICA. Nux vomica (vegetable.) *Antidotes*—Aconite, Camphor, Coffea, Pulsatilla.
47. OPIUM. White Poppy (vegetable.) *Antidotes*—Camphor, Coffea, Calcarea carb., Hepar sulph., Sulphur.
48. PETROLEUM. Rock oil (mineral oil.) *Antidotes*—Aconite, Nux vomica, Coccus.
49. PHOSPHORUS. Phosphor (mineral) *Antidotes*—Camphor, Coffea, Nux vomica.



50. PHOSPHORI ACIDUM. Phosphoric acid (a mineral acid.) *Antidotes*—Camphor, Coffea.
51. PLATINA (metal.) *Antidotes*—Pulsatilla, Belladonna.
52. PULSATILLA. Meadow Anemone (vegetable.) *Antidotes*—Chamomile, Coffea, Ignatia, Nux vomica.
53. RHEUM. Rhubarb (vegetable.) *Antidotes*—Camphor, Chamomile, Nux vomica.
54. RHUS TOXICODENDRON. Rhus. Sumach. Poison Oak (vegetable.) *Antidotes*—Belladonna, Bryonia, Camphor, Coffea, Sulphur.
55. SAMBUCUS (NIGER.) Elder (vegetable.) *Antidotes*—Arsenic, Camphor.
56. SANGUINARIA (CANADENSIS.) Common Blood-root.
57. SECALE (CORNUTUM.) Ergot of Rye (vegetable.) *Antidotes*—Camphor, Opium.
58. SEPIA. Inky juice of the Scuttle Fish (animal.) *Antidotes*—Aconite, Spiritus Nitri, Vinegar.
59. SILICEA. Siliceous Earth (mineral.) *Antidotes*—Camphor, Hepar sulph.
60. SPIGELIA (ANTHELMENTICA.) Indian Pink (vegetable.) *Antidotes*—Camphor, Aurum.
61. SPONGIA. (Marina Tosta.) Burnt Sponge. *Antidotes*—Camphor.
62. STANNUM. Pure Tin (metal.) *Antidotes*—Coffea, Pulsatilla.
63. STAPHYSAGRIA. Stavesacre (vegetable.) *Antidote*—Camphor.
64. STRAMONIUM. Thorn Apple. Gympsum weed (vegetable.) *Antidotes*—Belladonna, Nux vomica.
65. SULPHUR (mineral.) *Antidotes*—Aconite, Camphor, Mercury, Nux vom., Pulsatilla.
66. SULPHURIC ACID. Oil of Vitriol (a mineral acid.) *Antidote*—Pulsatilla.
67. TARTAR EMETIC. Tartarized Antimony (mineral.) *Antidotes*—Cocculus, Ipecac., Pulsatilla.
68. TABACUM. Tobacco. *Antidotes*—Camphor, Ipecac., Nux vomica.
69. THUJA (OCCIDENTALIS.) Arbor Vitæ Tree (vegetable.) *Antidotes*—Camphor, Pulsatilla.
70. VERATRUM (ALBUM.) White Hellebore (vegetable.) *Antidotes*—Ipecac., Arsenic, Camphor, Coffea, Aconite, China.

## TINCTURES.

- |                         |                            |
|-------------------------|----------------------------|
| 1. ARNICA, Tincture.    | 2. RUTA, Tincture.         |
| 3. SYMPHYTUM, Tincture. | 4. URTICA URENS, Tincture. |

These tinctures are intended for external use only. In preparing them for this purpose, put from four to ten drops in a teacupful of pure water; mix well, and wash with it, or lay cloths, dipped in the mixture, on the parts affected.

## NOTICE TO HOMŒOPATHIC PHARMACEUTISTS.

It will be observed, that in this "Domestic Physician," *Coffea*, *Belladonna*, and *Cina*, are used in two preparations—the third and the two hundredth. The latter has <sup>cc.</sup> attached to it, as *Belladonna* <sup>cc.</sup> showing, in the Roman numerals, the potency. It is expected that the preparations of these latter for the boxes are made in the greatest purity and genuineness. They can be procured already in every large city of the Union.

The potencies of all the other remedies are designed for those from the mineral and animal kingdoms in the sixth, and for those from the vegetable kingdom in the third potency. The tinctures mentioned in the "List of Medicines" should accompany the boxes.

## INTRODUCTION.

---

To aid in the search for the proper place in the work, where a disease may be found, its different popular names have been always given, and recorded carefully in the index. Should, however, the identification of a disease be difficult or doubtful, it is only necessary to refer, in the book, to the divisions made as to the parts of the system which are affected, such as "Affections of the Head, Chest," etc., where all the ailments of these parts are generally put together. But there will scarcely ever arise a difficulty in this respect; the index has been made very complete, in view of saving the reader time and unnecessary vexation.

If a disease is found, it is expected that every thing will be first read which is said concerning it in the article, before a selection of a remedy is made; at the end of each article, the proper direction as to administration, diet, and regimen, is given. Wherever the attention is directed to the perusal of the *Materia Medica*, in treating on certain remedies, the medicine in question must be looked for in the *Materia Medica*, and its symptoms carefully compared with those of the patient. If in the directions for diet the

reader is advised to adhere to the usual homœopathic diet in chronic diseases, etc., he will find its rules specified in the following pages, under "*Dietetic Rules.*"

It is necessary to give an explanation in regard to several remedies, whose names will be found different in this work from those used in other similar works. Of the various preparations of *Mercury*, only one kind has been used, and whenever this is mentioned, under the name of *Mercury*, the *Mercurius vivus* is understood by it. This was done on account of not unnecessarily confusing the reader with two different names for mercurial preparations, which are essentially the same in their effects and use. On that account the *Mercurius sublimat.* has been left out entirely. Its use is, at any rate, confined to one disease only (the dysentery), in the treatment of which the *Mercurius vivus* supplies its place completely.

On the other hand, three remedies—*Coffea*, *Belladonna*, and *Cina*—appear in two different preparations in the book, the one commonly used bearing the simple name, while the two-hundredth potency of these remedies have a <sup>cc.</sup> attached to them, showing in the Roman numerals the degree of potency. Although a great many more remedies could have been treated in the same manner, it was not thought advisable to burden the book with more therapeutical detail, as it might interfere with its other simple and practical arrangements. These three remedies, however, in their higher potencies, seemed to be almost indispensable, in a useful work of this kind, as their effects, when properly indicated, can not well be produced by their lower attenuations. We invite particular attention to the use of *Cina* <sup>cc.</sup>, in the nervous worm symptoms, so frequently experienced in children, in nightly attacks, which quickly disappear after the exhibition of one or two doses of *Cina* <sup>cc.</sup>. Every homœopathic practitioner can soon satisfy himself in regard

to the efficiency of the highest potencies, by making the above trial.

In regard to our views on the preference of higher or lower attenuations, we would remark, that we consider homœopathically legitimate and practically useful, all potencies, from the mother tincture and first trituration up to the highest dilution, but that we do not use one exclusively or all indiscriminately. Practical observation and theoretical researches have guided us in applying the various attenuations with real benefit, which is not so difficult as many practitioners might believe. There are principles according to which the different potencies of a remedy must be applied in different cases. But to enlarge upon this subject here, would carry us too far; we content ourselves by remarking that, for a domestic family-chest, we would propose, as a general rule, the third potency for all vegetable medicines, and the sixth potency for all mineral and animal medicines. If this rule was once adopted by all writers on homœopathic domestic practice, their several works might be used with benefit without subjecting the public to additional expense in procuring different medicine-chests for the works of different authors.

For a successful treatment of diseases, every thing depends, besides the selection of the right, or most homœopathic remedy, on the genuineness of its preparation. We would, therefore, advise our readers to procure their medicine-chests from persons whose competency or recommendations can be relied on; and when in their possession, to take good care of them. The phials must not be opened except when used; each phial must receive its own cork again, and no smelling medicinal substances must be allowed to be near by, when a phial is opened or medicine is taken. If a remedy is prescribed to be taken in water, a clean cup or tumbler must be selected, the pre-

scribed quantity of water, either rain or river water poured in, then the medicine added, and, during its solution, well mixed with a clean teaspoon, and afterward covered well. If more than one medicine is prescribed at a time, in water, each one must be dissolved separately in a teacup, and each one receive its own separate teaspoon, which must not be changed during the administration of the medicine.

If a phial is empty and shall be refilled, it is *indispensably* necessary to cleanse it first thoroughly with warm water and dry it perfectly on a stove or in the sun, before the homœopathic apothecary is allowed to fill it. Persons ought to attend to this scrupulously themselves, before presenting them for refilling; because too much depends on it, as regards the genuineness of the medicines.

The medicines should be kept in a dry place, where no odors can reach them. A renewal of their strength, if well kept, is not necessary for years; but if the slightest doubt in this respect should arise, it is better to have it renewed.

## DIETETIC RULES.

---

The observance of a strict diet is essential to the success of a homœopathic treatment, not so much on account of the injury done to the system by the use of improper articles, as because these might be antidotarian in their effects to the medicine. On this account, homœopathists so generally forbid coffee, it being an antidote to most of their remedies. For the same reason, the use of all other things of a medicinal nature is strictly forbidden, such as spices, perfumeries, perfumed waters, soda waters, spiced ices, etc.

It is a general law of nature, to eat and drink only that which can sustain life, without producing any other effects, that is, simple nutriment ; and as medicinal and spicy substances, although sometimes nutritious, are irritating and stimulating, the strict law of nature excludes them peremptorily from the daily use (see “Hygiene.”)

*Diet in Acute Diseases.*—As dietetic rules are given for each acute disease, separately throughout the book, it is not necessary to mention them in this place.

*Diet in Chronic Diseases.*—Consult the chapter on “Hygiene” on this subject.

There are certain principles which must guide us, during the treatment of chronic diseases, in selecting the articles of our diet, even from among those which are gen-



erally allowed. If the patient has symptoms of fever or irritation, in any part of the system, he must avoid all articles of a stimulating character, such as meats, eggs, and butter. In such cases, he should confine himself to a nutritious vegetable diet, such as potatoes, turnips, rice, barley, etc. If the disease appears mostly in the digestive organs, as a chronic diarrhea, all articles of a relaxing nature are in themselves injurious, such as fruits and other green vegetables, soups of meat, eggs, etc. In cases where constipation is present, the opposite course must be pursued.

Another consideration is, that articles of food, which disagree with the patient, although allowed, or wholesome for others under similar circumstances, must be strictly avoided by him.

#### ALIMENTS ALLOWED.

Beef and Mutton baked, roasted, or boiled, fresh or smoked Tongue, lean Ham.

Venison and wild Fowl.

Turkeys, Chickens, Pigeons.

Fresh scale fish, oysters.

Butter which is not rancid, mild new Cheese, raw or soft boiled Eggs, or Eggs in Soups (except in cases of diarrhea), Meat-Soups, Broths, seasoned with salt only.

All kinds of light Bread and Biscuit (except Soda-Biscuit and other Bread made with the addition of unusual substances.)

Puddings, Dumplings or Noodles of Wheat, Indian, Rice, and Oat meal, not too heavy, fat, nor spiced.

Cakes composed of Meal, Eggs, Butter in small quantity, and Sugar.

Food prepared of Arrowroot, Tapioca, Farina, Sago, Salep, Oatmeal, and the like.

Irish Potatoes, Sweet Potatoes, Turnips, Carrots, To-

matoes, Colliflower, Spinage, and green Peas, or Beans (except in Colic and Diarrhea), Rice, Hominy, Pearl Barley, dried Peas, or Beans.

Roasted or boiled, fresh or dried Fruit, as Prunes, dried Currants, fresh ripe sweet Apples, Peaches, Strawberries, Raspberries, Gooseberries, and other sweet berries, and Grapes (except in Colic and Diarrhea.)

Water, pure, or sweetened with Sugar, Currant Jelly, Raspberry, or Strawberry Syrup, Toast water.

Milk, in its different preparations, boiled Milk, fresh Whey, Buttermilk.

Decoctions of Barley, Malt, Oat-meal, Farina, Rice, dried Fruit. Cocoa boiled with Milk or Water, pure plain, Chocolate, weak black Tea.

Salt moderately used, Sugar.

Icecreams flavored with Strawberry and Lemons.

#### ALIMENTS FORBIDDEN.

Old smoked, salted Meat and salted Fish, old rancid Butter and Cheese, Lard, fat Pork, Geese, Ducks, Turtles and Terrapins, Fish not having scales, as Catfish, Eels, Lobsters, Crabs, Clams, and Soups, prepared from these articles of food. All food prepared with Blood, and much animal fat. All kinds of Sausages, particularly such as are smoked. The flesh of all young animals. All Soups highly seasoned, Sauces, different kinds of Hash, drawn Butter, Peppertot.

All kinds of Cakes, or Pastry, prepared with much Fat or Aromatics, Honey. All kinds of colored Sugar-work prepared by the confectioner, red Sugar, sugared Almonds, or Peach-kernels. Nuts of all kinds.

Radishes, Celery, Horse-radish, Garlic, Parsely, Red or Cayenne Pepper, Mustard, Saffron, Nutmeg, Vanilla, Laurel leaves, bitter Almonds, Cloves, Cinnamon, Allspice,

Coriander, Fennel, Anise, Coffee, green Tea, spiced Chocolate.

All kinds of spiritous liquors, Brandy, Rum, Whisky, Gin, Spirits, etc. Liquors, or Cordials, and other drinks, cold or warm, which are prepared with spiritous liquors, or spices.

All artificial and natural Mineral Waters, Mead, Spruce Beer, Soda Beer, Porter, Ale.

N. B. During the homœopathic treatment especial care should be taken by the patient to avoid allopathic medicine of every description and form, pills, herb-teas, etc., internally and externally ; particularly all salves, aromatic waters, hartshorn, smelling bottles, etc. ; also, all perfumery, musk, cologne water, or other aromatic substances, tooth-powders containing such ingredients, etc. ; especially be careful in using matches, not to inhale the vapor of the burning sulphur ; matches must be lit only where the vapor can easily escape in a fireplace, or open hall. Moderate exercise in the open air, as, also, a careful observance of all other hygienic rules (see the article on Hygiene) is strongly recommended.

PART FIRST.

---

DESCRIPTION AND TREATMENT  
OF  
DISEASES.



## CHAPTER I.

### GENERAL DISEASES.

#### RHEUMATISM.

DIAGNOSIS.—Pains in the muscles and membranes, together with swelling of the surrounding cellular tissue, light redness, and heat; caused by having taken cold.

This is the general appearance of all rheumatic affections; yet their degrees and complications are so numerous, that it would almost require a separate treatise to do justice to such an extensive subject. We confine ourselves, here, to the more practical subdivision of *acute* and *chronic* rheumatism.

Beside, we would remark that rheumatism, particularly in the acute form, can exist in any part of the system, creating there such an inflammation as the affected organ is naturally disposed to, adding to it only its own rheumatic character. For instance: a patient is attacked with rheumatism in the arms or limbs; this at once leaves those muscles, and falls on the pleura, where it generates a *pleurisy*, not of the usual kind, but one which has the rheumatic character—a *rheumatic pleurisy*. In the same manner, we speak of rheumatic pneumonia, rheumatic inflammation of the eye, heart, etc. In such cases, we refer the reader to the respective chapters, where the diseases of these organs are specially treated.

There exists an intrinsic difference between rheumatism and gout. (See this article.)

*a.* ACUTE OR INFLAMMATORY RHEUMATISM.

This form of rheumatism, especially, develops itself after taking cold, or after any check of perspiration, sudden and severe enough to unfit the whole skin for its proper function of exhaling the imperceptible gases. First, an uncomfortable sensation and restlessness will appear, followed by chilliness and feverishness in alternation, thirst, constipation, and accelerated pulse; then the rheumatic pains appear in places where they either fix themselves, creating swelling, heat, and redness, or wander to other parts, where they repeat the same process, while those first attacked heal gradually and get well. In acute rheumatism, we sometimes find a dry skin; at other times, a very moist one, drenched with a watery perspiration, which does not relieve the patient. The perspiration which will relieve the patient is greasy, thick, glutinous, and smells acrid or sour.

TREATMENT.—In the beginning, when the fever is high, give, immediately,

*Aconite*, dissolved, 12 glob. in half a teacupful of water, every two hours a teaspoonful, for six or eight times, or until the fever has subsided. After it,

*Bryonia*, in the same manner, when the pains are worse by *motion* or *at night*, when there is headache, fever, swelling of the joints, and gastric derangement. (See Bilious Rheumatic Affection.) Or,

*Belladonna*, when *Bryonia* does not relieve, or when the parts are much swollen, *very red, shining*, and the patient is sleepless at night, complaining of dryness in the mouth and throat. Both remedies may be given in alternation, with advantage.

*Chamomile*, when, in the parts affected, there is a sensation of *numbness* or *paralysis*, worse at night, the patient is feverish, restless, agitated, *irritable*.



*Nux vomica*, in alternation with *Chamomile*, when the parts are numb, *cold air aggravates*, temper is irritable, bowels constipated.

*Pulsatilla*—the same feeling of numbness and paralysis, but *relieved by exposure to cool air*, the patient wants to be uncovered, and is of a mild temper and whining mood, worse in the night; the pains have a great tendency to change places. (*Shifting or wandering Rheumatism*.)

*Mercury*—pains *increased* in the heat of the bed, or *toward morning*—also, when the patient *perspires profusely without being relieved by it*. *Lachesis* suits well after *Mercury*.

*Rhus*—principal indication for its use is: *pains worse during rest*, or as if the flesh was torn from the bones; worse in *cold, damp weather*. This remedy, when the latter symptoms are present, alternates well with *Bryonia*. (See Bilioid Rheumatic Affection.) It is, also, of great use, when there is *paralytic weakness* or *trembling of the extremities*, on moving them.

ADMINISTRATION — Give the above medicines in solution, as stated under *Aconite*, but pause, after four teaspoonfuls of each one have been given, for eight or twelve hours, during which time the effect must be looked for before another remedy is selected. As soon as the patient is better, all medicine must be stopped, until he gets much worse again.

DIET AND REGIMEN.—As regards nourishment, the patient must absolutely abstain from all meat, or soups of meat, even long after the convalescence. He can have, however, gruels of farinaceous substances, oatmeal, farina, rice, etc., toast-water and cold water; during the fever, warm lemonade and oranges. Cover the patient well with blankets, as many as his comfort dictates. Around the swollen joints and other parts affected, breast or neck, put wool, just from the sheep, or, at least, unwashed. This

has, frequently, a very good effect. If possible, consult a homœopathic physician.

*b.* CHRONIC RHEUMATISM, or commonly called RHEUMATISM.

If a person is severely troubled by chronic rheumatism, let him at once apply to a homœopathic physician, who will give him relief, if it is at all in the power of medicine to do so. But, frequently, the reactive force of nature is already so much weakened, that it first requires to be strengthened, which can only be done by a systematic hydropathic treatment. However, in cases of rheumatism, where the system is yet strong, the medicines, as detailed below, may be used, and will be found efficacious in a majority of them. Beside the remedies recorded under "Acute Rheumatism," which may also be used for chronic rheumatism, the following are the principal ones: *Ignatia*, *Thuja*, *Arnica*, *Dulcamara*, *China*, *Veratrum*, *Arsenic*, *Phosphorus*, *Caustic*, *Sulphur*, *Sepia*.

SYMPTOMATIC DETAIL. — *Ignatia*: *Contusive pains*, or as if the flesh were detached from the bones; worse at night, ameliorated by change of posture.

*Thuja*: *Tearing, pulsative pains*, as from ulceration under the skin, with a feeling of coldness and torpor of the parts affected. Pains worse during repose, or in the warmth of the bed. Pain in the right shoulder and arm, *better on movement*.

*Arnica*: Pains as if the parts were *strained or bruised*, with a feeling as if *they were resting on too hard a surface*; a tingling in the hand, red and *shining* swelling, worse when attempting to move. (Suitable before or after *China*, *Arsenic*, *Rhus*.)

*Dulcamara*: Pains after getting cold, wet, worse at night, during repose, and without much fever.

*China* : Pains, worse on the slightest touch, with easy perspiration, the sore parts feel very weak, almost paralysed.

*Veratrum* : Pains, as if from a bruise ; worse by warmth and bad weather ; better by walking ; very weak and trembling.

*Arsenic* : Burning, tearing pains, insupportable at night, worse by cold air, and mitigated by external heat.

*Phosphorus* : Tearing and drawing pains, excited by the slightest chill ; headache, vertigo, and oppression of the chest.

*Caustic* : The pains are *insupportable in the open air* ; less severe in a room, or in bed ; also, when there is paralytic weakness with *rigidity and incurvation of the parts affected*.

*Sulphur* : In almost all cases of chronic, and after a spell of inflammatory rheumatism, when the pains yet linger about. Often after *Aconite, Belladonna, Bryonia, Mercury, or Pulsatilla*.

*Sepia* : For rheumatic affections in tall, slender persons, especially females.

ADMINISTRATION. — In chronic rheumatism, the remedies ought not to be repeated often ; every three or four days, one dose (four to six glob.) is enough. After a remedy has been tried for ten or fourteen days, another may be chosen, if no improvement has appeared.

DIET AND REGIMEN. — The patient must abstain, during the treatment, from all meats, at least, as much as possible ; must not expose himself to the changes of the weather, and should wear, on his skin, flannel underclothes, except while using hydropathy ; in which case, he can do without them.

#### GOUT. (*Arthritis*.)

DIAGNOSIS. — The symptoms of this disease are very similar to those of inflammatory rheumatism ; consequently, the remedies will be almost the same. It is characterized

by pains in the joints, with inflammatory swelling; or, in chronic cases, with a swelling of the joints, caused by deposits in them of a calcareous substance, which impedes their movements and causes them to make a cracking noise. There is always connected with an attack of gout, flatulency, acid stomach, and other derangements of the digestive organs. The principal differences between rheumatism and gout are: rheumatism attacks more the muscles and membranes—gout, more the joints; rheumatism is hardly ever complicated with derangement of the digestive organs; gout is never without that—has, besides, permanent swelling of the joints, by calcareous deposit, which rheumatism never has. Rheumatism is caused by taking cold; consequently, depends on external causes; while gout is generated by internal causes, amounting, sometimes, to hereditary predisposition. Both diseases, however, can intermix with each other, one taking the form of the other.

We again acknowledge an *acute* and *chronic* form of gout; recommending for both the same remedies, which we re-recorded under the two forms of rheumatism. We would add only one more remedy to this list; it is

*Antimon. crud.*, when the attack is attended with nausea and a white-coated tongue; pains are worse *after eating*, in the night, after drinking wine, and in the heat of the sun; mitigated during repose, and in the cool air.

To make the selection of remedies easier, we will give a tabular view of them, in the different gouty complaints. Their details will be found under "Rheumatism."

For *inflammatory* gout: *Aconite*, *Antimon. crud.*, *Bryonia*, *China*, *Nux vomica*, *Arsenic*, *Pulsatilla*, *Arnica*.

For the same, with *gastric derangement*: *Antimon. crud.*

For the *shifting, wandering, flying* gout: *Pulsatilla*, *Arnica*, *Nux vomica*.

For gout with *great swelling*: *Antimon. crud.*, *Arnica*, *Bryonia*, *Rhus*, *China*, *Sulphur*.

For the gouty *nods and lumps*: *Antimon. crud.*, *Bryonia*, *Calcarea*.

For gout in persons addicted to *spiritous liquors*: *Nux vomica*, *Aconite*, *Sulphur*, *Calcarea*.

For gout in persons called *high livers*: *Pulsatilla*, *China*, *Antimon. crud.*, *Sulphur*, *Calcarea*.

For those who sometimes *work in the water*: *Pulsatilla*, *Sulphur*, *Calcarea*, *Dulcamara*, *Rhus*.

For gout confined to *the big toe (Podagra)*: *Pulsatilla*, *Arnica*.

ADMINISTRATION, DIET AND REGIMEN, the same as in "Rheumatism."

#### PAINS IN THE SMALL OF THE BACK AND LOINS.

##### *Notalgia. Lumbago.*

DIAGNOSIS. — Violent pain in the region of the small of the back and loins, more or less permanent, sometimes periodical, but mostly excited by seemingly external causes. This species of lumbago (*back and loin-ache*) occurs, frequently, after any quick motion of the back, as in rising from a stooping position. The patient is suddenly seized with a violent pain as if produced by an arrow shot into the part; it pins him, as it were, into a fixed attitude, from which he can not stir without suffering torture, and forces him to keep quiet.

Although this disease is essentially rheumatic in its character and process, yet its foundation is constitutional; sometimes a predisposition is acquired by overstraining the muscular system by hard labor during exposure to wet and cold, or, even without that, a disposition to hemorrhoidal (*piles*) congestion of blood to these parts, which may sometimes increase to a real *inflammation of the spine*, the *spinal*

marrow (*myelitis*), or go over, if not cured, into a chronic spinal affection.

TREATMENT.—*Aconite* is the principal remedy in the commencement, and does more than the lancet can accomplish, which the old school always applies in such cases, but ought not to be used, since *Aconite* fulfills all our expectations.

*Arnica*, alternately with *Rhus*, if this disease is really the result of falling, overlifting, or any other mechanical injury.

*Bryonia*, if the pains in the back are *pressing*; the patient can not walk erect; worse after the slightest motion, or current of cold air; patient feels chilly; head and limbs ache.

*Rhus*, in alternation with *Bryonia*, if the patient feels very weak, trembling, has to get up sometimes to ease himself; throws himself about in the night. After external injury, see *Arnica*.

*Bellad.*, after *Aconite* and *Bryonia*, under similar symptoms, but more suitable for fleshy females during the (change of life) critical period; in alternation with *Rhus*, when the feverishness attending the attack is complicated with *restlessness*, dryness of mouth and throat, headache.

*Nux vomica*, if the parts feel as if tired, or *very much fatigued*, during the attack; pains are worse by *motion* and *turning in bed*; constipation, irritability of temper. Suitable for persons with a disposition to piles, or addicted to spiritous liquors.

*Mercury*, in alternation with *Nux vomica*, if catarrhalic diseases, influenza, etc., prevail, or a tendency to dysentery; or if the patient feels very weak, *perspires* a great deal without relief, worse at night.

*Pulsatilla*: Resembling the pains under *Nux vomica*, but caused by obstructions or irregularity of the courses



(*menses*) in younger females of a mild, sensitive, or phlegmatic character.

ADMINISTRATION.—Same as in “Rheumatism.”

DIET AND REGIMEN.—Also the same. We would advise those afflicted with this disease, to lie down immediately on a mattress, or something similar, and not attempt to brave it out, as the medicine will not be able to cure as quickly, and chronic weakness of the back remains, together with a liability to a return of the disease.

### PAIN IN THE HIP. HIP DISEASE.

#### *Sciatica. Coxalgia. Coxagra.*

DIAGNOSIS.—Pain in the region of the hip joint, extending to the knee, even to the foot, accurately following the course of the sciatic nerve. Its continual severity may impede the motion of the foot, producing stiffness and contraction, disturbing the rest at night, and thereby inducing general uneasiness and emaciation.

In *sciatica*, or *pain in the hip*, the pain manifests itself only on the *outside* of the hip and leg, during *repose* as well as during *motion*, showing its *neuralgic* character; while, in *coxagra* (*morbus coxarius*), the pain appears only during *motion* and in *stepping*, showing its *inflammatory* character; running in front to the knee, protruding and lengthening the leg. It can terminate in suppuration (*white swelling*).

These diseases are of great importance, and ought not to be neglected, but immediately put under the care of a skillful homœopathic physician, as scrofula is either their remote cause, or excited by them, especially in children, in whom they often occasion the so called *spontaneous limping* (*coxalgia infantilis*) and *white swelling*, particularly when badly treated at first by allopathic remedies.

TREATMENT.—The principal remedies in these diseases

are those again already enumerated under the headings of "Rheumatism" and "Gout." We will give them here again, to facilitate the selection :

In *sciatica (ischias)*: *Chamomile, Bryonia, Rhus, Arsenic, Ignatia, Nux vomica, Pulsatilla.*

In *coxagra (hip disease)*: *Colocynth, Belladonna, Hepar, Lachesis, Silicea, Mercury, Sulphur.*

In *spontaneous limping*, *Mercury* and *Belladonna* in alternation; every few days, a dose: *Rhus, Colocynth, Pulsatilla, Sulphur, Calcare.*

SYMPTOMATIC DETAIL. — *Aconite*, in alternation with *Belladonna* or *Bryonia*, where the attack is accompanied with a great deal of fever.

*Belladonna*, in the inflammatory stage (see *Aconite*), when the pain is increased on the slightest movement, with limping, and on the parts affected the skin is red and shining; dryness of the throat; burning fever. After it, *Mercury*.

*Bryonia*, under similar symptoms as *Belladonna*, but particularly where the parts affected are more painful than red; constipation. After it, *Rhus*.

*Rhus*: Darting, tearing, or dragging pains in the hip-joint, with tension and stiffness in the muscles; worse during repose, or when rising from a sitting posture.

*Chamomile*: Pains worse at night in bed, after having recently taken cold; patient is irritable; does not know what he wants.

*Mercury*: Sharp, cutting, burning pains; worse at night, or during movement; profuse perspiration without relief. After *Belladonna*; before *Hepar*.

*Ignatia*: Cutting pains, ameliorated by change of position, temperament mild, yet vacillating between high and low spirits.

*Pulsatilla*: Pains worse in the evening or night, and



when seated; *relieved in the open air*; temperament phlegmatic; mild disposition.

*Nux vomica*: Pains worse in the morning; *worse in the open air*; temperament irritable, morose; constipation.

*Arsenic*: Burning pains, with great restlessness, obliging the patient to move the limb; pains appear *periodically*, or in spells, (alternate with *China*); mitigated by external heat; patient wants to lie down, feels very weak; emaciation of the limb after long suffering, (in alternation with *Silicea*.)

*Colocynth*. This is the first and principal medicine in hip diseases, acute or chronic, particularly when there is the sensation of a *tight band around the hips and back*, and the pains *run down from the region of the kidneys into the leg*; *spontaneous limping*; pains are worse lying on the back, or after a fit of anger or indignation.

*Hepar* after *Mercury*, when this has not relieved.

*Silicea*, when the disease assumes the chronic form, after *Hepar*; or, where emaciation has taken place, in alternation with *Arsenic*, and, afterward, with *Iodine*.

*Sulphur*, in chronic cases, where other remedies have not entirely relieved.

*Lachesis*: Emaciation, with tearing and contracting pain in the joint; *dread of exercise*; *deeply penetrating suppurations*, (in alternation with *Silicea*.)

ADMINISTRATION.—In the *sciatica* (*ischias*), as well as the febrile stage of the *coxagra*, give the medicines as required in the acute form of "Rheumatism," (see this article.) In the chronic form (suppurative stage) of *coxagra*, give the remedies as stated in "Chronic Rheumatism."

DIET AND REGIMEN.—The same as in "Rheumatism," but without outward applications, save cold-water bandages, in any stage of the disease, when the patient can bear them, and feels relieved after them.

INFLAMMATION OF THE KNEE JOINT. (*Gonitis.*)

Inflammation and swelling of this important joint, requires the use of *Aconite* and *Belladonna*, *Bryonia*, *Rhus*, *Lachesis*, *Mercury*, one after the other, if the first two have not allayed the inflammation. If suppuration threatens, which is indicated by the swelling not going down after the severest fever is over, give *Sulphur*; in eight days, *Silicea*, and consult medical aid.

ADMINISTRATION of *Aconite*, *Belladonna*, and the other remedies, the same as in "Acute Rheumatism."

DIET AND REGIMEN.—Also the same, but without any outward application, save cold-water bandages, when the patient feels relieved by them, at any stage of the disease.

INFLAMMATION OF THE PSOAS MUSCLE. (*Psoitis.*)

DIAGNOSIS.—Pain in the region of the kidneys, hip, and downward to the leg, which can not be stretched, or drawn near to the abdomen, without pain; increased, also, by turning when lying and lifting, with a feeling of numbness of the affected side; walking is possible only by hobbling, with the body bent forward. Although not often fatal, yet it is very important in its consequences, as its issues may give rise to lingering diseases. If suppuration takes place, by not preventing it in time, the matter discharges itself in the abdomen, and causes death; or it sinks down farther and farther, until it reaches, sometimes, even the knee, before it escapes; frequently, the spine becomes affected and is rendered carious in such cases. The causes of this disease are, beside external injuries, rheumatism and piles.

TREATMENT.—Give, first, *Aconite* and *Bryonia*, in alternation, every two, three, or four hours, a dose (four glob.); afterward, *Bellad.* or *Nux vom.*, if not relieved within twenty-four hours, and when the pains increase

during motion; *Rhus* and *Pulsatilla*, however, when the pains are worse during rest; *Mercury* and *Chamomile*, when the pains are worse at *night*.

*Staphysag.*, when the pains are beating, indicating incipient suppuration.

*Colocynth*, when the disease assumes the chronic form.

If possible, consult a physician in this important disease.

ADMINISTRATION, DIET, AND REGIMEN, as in "Affection of the Knee Joint."

### CRAMP IN THE LEGS.

Some persons are habitually addicted to cramp in the legs, from various causes. The first remedy, which every one naturally will resort to, is, to rub the parts taken with the cramp, either with the hands or a rough towel, which, in a short time, will destroy the spasm. Another expedient is, to jump on the cold floor of the room, if the cramp occurred on the calves of the legs, and while in bed; but, if it should occur too frequently in the *night*, take

*Veratrum*, every night a dose (four glob.), for a few nights; or, if it occurs in pregnant females,

*Secale*, in the same manner.

If in the day time, take *Rhus*, in the morning, in the same manner.

*Sulphur* may finish the cure, if necessary.

DIET AND REGIMEN, as in all chronic diseases.

### SWEATING FEET.

This is a complaint with which a great many persons are afflicted. It is generally connected with a qualitative corruption of the secretions, by which a bad smell is emitted, thereby rendering the complaint very disagreeable. It can be easily suppressed by alum and sugar of lead

ablutions; but he who ventures to do so, risks being taken with blindness, deafness, asthma, consumption, etc.

Homœopathy possesses remedies, which, when applied rightly, will insure a permanent and safe cure.

TREATMENT. — *Rhus*, inwardly, every third evening, six glob., to be continued for, at least, four weeks, when two weeks must elapse before the next remedy may be taken. During the first four weeks, bathe the feet every other evening, in cold water, into which four drops of the mother tincture of *Rhus* (to be had in any homœopathic pharmacy) have been dropped.

*Silicea* is the next remedy after *Rhus*, if this has not already ameliorated the complaint; to be taken in the same manner. In this way, alternate with these two remedies until better, or apply to a homœopathic physician, who has more remedies at his command, which can be chosen, however, only according to the individual constitutionality of each one's case.

#### GOITRE. (*Struma*.)

This disease consists in a swelling of the thyreoid gland, in front of the throat. It is, in the majority of cases, of a scrofulous origin; depends, sometimes, however, on an endemical cause; it occurs most frequently on mountains, especially at their base, and in their valleys.

TREATMENT. — Take, internally, *Spongia*, every evening a dose (six glob.), and wash externally with a solution of the tincture of *Iodine* diluted in alcohol, every evening, until it disappears.

If this treatment does not disperse the goitre within two months, apply to a homœopathic physician, who has other remedies, suitable for each individual case.

DIET, as usual in chronic diseases.

SLEEPLESSNESS. (*Agrypnia.*)

This disease consists in an impossibility to sleep, without any other disease, or any external or internal cause to disturb the sleep. If it continues for months and years, a serious disturbance of all functions must ensue. In infants, particularly, it becomes a distressing circumstance for mothers and nurses, as we often are unable to find out its cause.

TREATMENT.— *Coffea* and *Belladonna*, two doses in alternation (in children, *Coffea* <sup>cc.</sup> and *Bellad.* <sup>cc.</sup>), every hour one dose (four glob.) frequently allays the overexcitement of the nervous system.

If this will not do one night, try, on the next,

*Hyoscyamus*, especially after severe illness.

*Ignatia*, when caused by *grief*, *indignation*.

*Nux vomica*, when from study and meditation in hypochondriacs.

*Opium*, after fright and fear, or in old people, or where frightful visions appear when closing the eyes.

*Pulsatilla*, when having indulged too freely in eating.

*Aconite*, when caused by agitating events and anxiety.

If caused by drinking tea or coffee, see their antidotes in the articles respecting these substances.

In *children*, beside *Coffea* and *Bellad.*, are recommended, *Cham.*, *Jalap*, and *Rheum.*, when it is caused by *colic*. See this article in "Diseases of Children."

Beside these remedies, a sponge-bath of cold water every evening when going to bed, is strongly recommended.

NIGHTMARE. (*Incubus.*)

A well-known troublesome disease, consisting in a heavy pressure on the precordial region, which impedes breathing, creating thereby many images of fancy, monsters, robbers,

bears, etc. It occurs mostly in the first hours of sleep, and, if recurring every night, would certainly injure the general state of health.

It is caused either by an overloaded stomach, congestion of the blood to the abdomen, or to the precordial region, when lying on the back.

TREATMENT.—Persons liable to this disease must first avoid the above-mentioned exciting causes, and then the following remedies will have their effect:

*Aconite*: Especially in women and children, when feverishness, oppression of the chest, anxiety, and inquietude, prevails.

*Nux vomica*, after drinking spiritous liquors, eating a full meal in the night, or by sedentary habits.

*Opium*: The principal remedy in severe attacks, snoring respiration, eyes half open, face covered with cold perspiration, and convulsive movements of the limbs.

Beside these remedies, daily exercise in the open air, the use of cold water internally, and externally in shower or sponge baths, are recommended.

### PALSY. (*Paralysis.*)

This affection of the nerves of voluntary motion, is one of those diseases, to remove which, it requires the greatest skill of a physician; and, by bringing it under our notice here, we intend merely to warn our readers not to waste time, health, and money, in using a variety of nostrums which might be recommended for it; but to apply at once to a skillful homœopathic physician, who can effect a cure in a majority of cases. Besides, we will record a few remedies, with which such a cure may be commenced.

If from *debility* caused by loss of fluids, *China*, *Ferrum*, *Sulphur*.



If from suppression of an eruption or habitual discharge, *Lachesis, Sulphur, Caustic.*

If from *rheumatism, Bryonia, Rhus, Arnica, Lycopodium.*

If from *apoplexy, Ipecac., Lachesis, Lycopodium.*

If from handling *whitelead*, or exposure to the fumes of *lead, Opium, Bellad., Platina, Pulsatilla.*

If the *facial muscles* are paralysed, *Bellad., Graphites, Caustic.*

If the tongue, *Bellad., Opium, Stramonium, Hyoscyamus, Lachesis.*

If the *arms, Bellad., Lachesis, Nux vomica, Opium, Lycopodium.*

If the *lower limbs, Cocculus, Opium, Nux vom., Stannum, Silicea.*

*Electricity* and *Galvanism* may be used in this disease with the greatest advantage, as also the cold water in bathing and drinking. See “Hydropathy.”

#### DELIRIUM TREMENS. (*Mania a potu.*)

This terrible disease is almost exclusively confined to drunkards and opium-eaters, who are taken, especially when exhausted otherwise, with a delirium and frenzy, in which appearances of horrible monsters, animals, figures of all kinds, frighten their imagination, combined with ravings, convulsion fits, and complete inability to sleep. In the long and sleepless hours, they converse incessantly with these supposed realities, by which they wear out their strength more and more.

TREATMENT. — *Opium* is the specific in this disease, if it is caused by ardent spirits, as this drug is itself able to produce such a disease. Give of it, every hour or two hours, a dose (one or two drops of a diluted tincture of opium), for at least twenty hours, to see its effects. After this, give *Bellad., Nux vom., Sulphur*, one after the other,

each, for twenty-four hours, every three or four hours a dose (six glob.) If nothing will produce sleep, give *Lachesis* and *Arsenic*, in alternation, every three hours a dose, (four glob.)

Let the patient drink freely of *ice water*, as the best stimulus which can be substituted. For solid nourishment, give him hard toasted bread ; as soon as he is convalescent, give him good beefsteak, bread, and water, on which he has to subsist for a long time. In this way, I have cured very bad cases, and had the satisfaction to see them not only restored to health again, but to usefulness and their friends, as they never relapsed into their former error.

### EPILEPSY. (*Epilepsia*.)

DIAGNOSIS.—Convulsive motions, with loss of consciousness ; falling down, with cries, foaming at the mouth, the thumbs flexed into the palms of the hands. The loss of consciousness is the most essential symptom, not the violence of the convulsions. The weakest convulsion, with unconsciousness, is epilepsy, while the most violent convulsion, but with consciousness, is *not* epileptic.

Most of the epileptic patients can be cured, but it requires the attendance of a skillful homœopathic practitioner. We refrain, therefore, from mentioning here any more of the treatment of this disease, but that which relates to the attack itself.

TREATMENT.—A patient in an epileptic fit must be placed in a position in which he can not hurt himself ; give him, however, full liberty of his own actions, without holding him, or forcing open his thumbs, which is of no use. If convenient, put between his teeth a cork or piece of wood, to prevent his tongue being injured. As soon as possible, give him one dose (four glob.) of *Belladonna* ; or if his face is dark and congested, his breathing very hard and snoring, a dose of *Opium*, (four glob.)



ST. VITUS'S DANCE. (*Chorea.*)

DIAGNOSIS.—Involuntary motions of single members of the body, mostly in such a manner as the patient was in the habit of using them, wandering from one part to the other, with full consciousness, which is a distinctive feature of this disease from epilepsy. It varies very much in degree; occurs most frequently at the time of the development of puberty, from the seventh to the sixteenth year, more among the female sex, more in moist regions on the sea-coast than in more elevated places. It may occur, also, as an epidemic, and is then infectious, particularly when large crowds meet. It is not a dangerous disease, but troublesome, and may lead to other derangements; ought to be attended to, therefore, immediately, as, in such a case, it can easily be cured.

TREATMENT.—As we recommend our readers to apply, in a case of this kind, to a homœopathic physician at once, we limit our remarks on the treatment to a few remedies, which may be given in the beginning.

*Ignatia*, every evening a dose (six glob.) for eight days; after it,

*Sulphur*, every other evening a dose (six glob.) for eight days, and then discontinue for three or four weeks, to await the effect.

DIET AND REGIMEN.—No greasy substances, no coffee or tea, but good plain food. Try to divert the attention of the patient from his disease; never speak of it in his presence.

TETANUS. TRISMUS. (*Lockjaw.*)

DIAGNOSIS.—Constant spasmodic contraction of one muscle, or all the muscles. According to the parts affected and the direction in which the body is drawn, the disease has received different names, which distinction does not

come, however, within the limits of our description, as such severe diseases as these will require medical aid. We intend here to speak more especially of one form of tetanus, called *trismus* or *lockjaw*, which is of frequent occurrence in southern latitudes, and, from its quick termination, requires prompt action and, therefore, domestic attention.

### LOCKJAW. (*Trismus.*)

It arises immediately after a wound has been inflicted, in consequence of the violent pain and nervous irritation, in which case it proves fatal in a very short time ; or it occurs in the first few days after the infliction of a wound, during its inflammatory stage, or from eight to ten days afterward, while the wound is healing and suppurating, without any pain and inflammation. The exciting causes are, mental affections, taking cold, corrupt air, foreign bodies in the wound, tension and distraction of single fibres in the wound. The most dangerous are the stitch-wounds in tendinous parts, as in the sole of the foot and the palm of the hand. The pulse frequently remains normal, the head free, yet there is great anxiety and oppressed breathing.

TREATMENT. — Enlarge the wound, if possible, and poultice it with bread and milk. Besides, give the following remedies :

*Arnica*. In the beginning, externally, in a wash or fomentation ; internally, in globules or drops, every two hours a dose (four glob. or one drop.)

*Belladonna* and *Lachesis*, in alternation, in the same dose and time, if the spasms increase. If no improvement after twelve hours takes place, give *Opium* and *Hyoscyamus*, in the same manner.

*Rhus* and *Ignatia*, in case the body is bent backward, in the form of an arch. Administer it in the same manner.

*Stramonium*, in the same form of lockjaw, if the two former remedies were of no avail.

DIET AND REGIMEN. — The same as in fevers.

#### SOMNOLENCY. LETHARGY.

DIAGNOSIS. — A sleep, continued beyond the natural time, for days or weeks. During this time, short intervals of waking intervene, but the patient soon relapses into sleep again. Nourishment can be given only by injecting fluid aliments ; otherwise, the functions of life are not disturbed.

In such cases, a physician will be consulted ; but, up to the time of his coming, the following medicines may be given :

*Opium*, if the pulse is full and slow, the breathing snoring, and the face very red, even dark ; every three hours a dose, or until better.

*Belladonna*, if the head is hot and the feet are cold ; pulse accelerated and hard.

*Lachesis*, if the pulse is very weak ; besides, *Aconite*, *Veratrum*, *Pulsatilla*, *Phosphoric acid*, one after the other, if necessary.

#### FATIGUE, OVERHEATING, AND MENTAL EXHAUSTION.

*Arnica*, for a feeling as if the body were *bruised*, after over-exertion of the body, and too long and fast traveling on foot.

*Rhus* : Pains in the joints. After *Arnica*, if this was not sufficient.

*China* : Weakness after loss of fluids, or heavy perspirations.

*Veratrum*, if persons are so fatigued as to cause fainting, in alternation with *Ipecac*.

*Coffea* and *Camphor*, in alternation, when the system is exhausted by disease, abstinence from food, or violent exercise.

*Cocculus* and *Nux vom.*, if caused by long night-watching.

*Aconite*, if, with weakness, there is palpitation of the heart, pain in the side, difficulty of breathing, aching in the limbs, from running fast; and then in alternation with *Bryonia*.

ADMINISTRATION.—Dissolve twelve glob. of a remedy in half a teacupful of water, and give, every half-hour or hour, a teaspoonful, until better.

If *overheated* by bodily exercise in the summer, take a little brandy and water, and no cold drinks until restored again; or, if possible, take a warm bath for twenty minutes. If the limbs feel sore, rub with *Arnica*, or alcohol in which soap is dissolved.

For *mental exhaustion* by overstudy or anxious night-watching, take *Nux vomica*; if very much excited and sleepless, *Coffea*; if with fullness of the head, *Belladonna*.

#### FAINTING, SWOONING. (*Syncope*.)

Nervous persons, particularly females, are subject to fainting fits, excited by various causes, external or internal. The first thing to be done is, to lay the patient quietly on a bed or couch, where the fresh air is accessible; loosen everything tight about the neck, chest, and stomach, and sprinkle cold water in the face, for a minute or two; during this time some one has procured spirits of camphor, which now may be held under the nose, to be inhaled. This, in most cases, is sufficient to restore the patient for the time, who must now be let alone, to gather strength. If the cause of the fainting can be ascertained, one of the following remedies may be given, to destroy the bad consequences which the attack may have on the nervous system.

If caused by *fright*, *Aconite*, *Opium*, *Sambucus*, *Staphysag.*, *Veratrum*.

By excessive joy, *Coffea*, *Aconite*, *Opium*.

By *anger*, *Pulsatilla*, *Platina*, *Nux vom.*, *Chamomile*.

By *excessive pain*, *Veratrum*, *Aconite*, *Chamomile*.

By the *slightest pain*, *Hepar*.

By *grief*, *mortification*, *Ignatia*, *Colocynth*, *Platina*, *Mercury*, *Phosphor. acid*, *Staphysag*.

By *fear*, *Ignatia*, *Pulsatilla*, *Veratrum*, *Opium*.

By *depletion*, *blood-letting*, etc., *China*, *Carbo veg.*, *Veratrum*, when, also, a little wine or brandy and water may be given.

ADMINISTRATION.— Dissolve twelve glob. of the selected medicine in half a teacupful of water, and give, every five or ten minutes, a teaspoonful. If not relieved in twenty or thirty minutes, prepare and give another remedy in the same manner.

#### APPARENT DEATH.

Whenever a sudden extinction of life appears, our suspicion must be aroused, as regards the real or apparent death of the individual, in as far as we frequently might be able to restore the apparently dead to life again, if we would only take the trouble and have the patience to use the requisite means. And even when these are applied, it is often done in an unsystematic manner.

As most cases of apparent death occur under violent circumstances, such as drowning, hanging, etc., it is quite natural that the minds of the bystanders become agitated and confused, not knowing what is first to be done. This uncertainty and hurry of action, however, can not produce any favorable results in resuscitating the apparently dead.

It is of the utmost importance to remain self-possessed, to reflect well, and then to do only one thing at a time,

until all available means to restore life are exhausted. There are a great many accidents in life, by which its existence is put in jeopardy. We will enumerate them here, and the means which ought to be used. On poisoning, we give a separate chapter.

### 1. *Apparent Death from Hunger.*

If starvation was the cause of an apparent suspension of life, inject small quantities of warm milk mixed with a very little brandy or Madeira wine ; beside, lay cloths, dipped in warm milk and brandy, on the stomach. Do not attempt to give any nourishment by the mouth, until after the patient has commenced breathing again, at which time the warm milk may be given to him, drop by drop, through the mouth. Increase the doses of milk very gradually, until the patient can take a teaspoonful ; then a few drops of wine or brandy will be salutary. After a while, give him small quantities of beef-tea, or other broth. Solid food is not allowed, until after the patient has had a sound, healthy sleep ; and even then, he has to be very careful in not eating too much at a time, or indulging in anything indigestible.

### 2. *Apparent Death from Drowning.*

The following cautions ought to be observed :

1. Be quick, but not rough, in all that has to be done.
2. Do not roll the body on casks.
3. Do not hold it up by the feet.
4. Do not rub the body with salts or spirits, nor inject smoke or infusion of tobacco.
5. Do not bleed the patient.

But do the following, immediately :

1. Convey the body, carefully, in a raised position, to the nearest house, if possible ; or, if not, lay it on a dry, sandy place, in the hot sun.



2. Strip the body, and rub it dry ; then wrap it in warm blankets, either in a warm bed, or in heated sand or ashes, until other means of warming the body can be procured, such as bottles of hot water, warming pans, heated bricks, etc., which may be applied successively on the stomach, spine, thighs, under the armpits, and soles of the feet.

3. Wipe and cleanse the mouth, nostrils, and throat, carefully ; during this operation the body may be turned on its side, the head bent forward, to allow the water to run out of the mouth ; all the water which can run out is in the mouth.

4. Rub, continually and briskly, the whole body with the hands or with warm cloths. Do not suspend, however, the use of the other means.

5. From time to time, try to inflate the lungs of the patient, by introducing into one of the nostrils the pipe of a pair of bellows, carefully closing the mouth and the other nostril ; blow the bellows gently till the chest rises, then set the mouth and nostrils free and press lightly on the chest, to eject the air from the lungs ; after which, the same process must be repeated and continued for some time.

6. Immerse the body, if possible, in a warm bath, at blood heat.

7. Electricity or galvanism may also be used.

8. Put a few globules of *Lachesis* and, afterward, *Opium* on the tongue of the patient.

9. Continue this treatment for, at least, four or six hours ; even if then no signs of life appear, do not remove the body immediately from its position, but wait until signs of decomposition are manifest.

If, however, the patient recovers and can swallow, give him small quantities of warm wine or brandy and water. Besides, watch him and attend to his further wants carefully.

### 3. *Apparent Death from Freezing.*

See "Frozen Limbs," page 39.

### 4. *Apparent Death from Lightning.*

Remove the body into the fresh air, and, immediately, dash cold water on face, neck, and breast. If possible, cover the body all over (except the face) with newly excavated earth. These means are the best, because water and wet ground are good conductors of electricity, with which the patient's system is surcharged at the time. Inwardly, give him, from time to time, a few globules of *Nux vomica*, of which, also, a solution may be injected.

If the patient recovers, remove him to a light, sunny room, where he should be kept quiet for some time, without mental excitement.

### 5. *Apparent Death from Hanging, Choking, or Suffocation by Burdens and Pressure.*

Tight clothing must be removed, and the patient placed in an easy, half-erect position; the neck not bent forward too much. Then rub him gently, but steadily, with the hands or warm cloths, and give, inwardly, from time to time, a few drops of the following mixture: five drops of laudanum or tincture of opium in four tablespoonfuls of water; of which, also, injections may be made; five drops of the mixture to each injection. Afterward, apply means for warming the body, such as heated bricks, bottles of hot water, etc.

If this has been continued for an hour or two, and still no signs of life appear, mix a drop of Prussic acid in a tumblerful of water, or pound a bitter almond fine and mix it in a tumblerful of water, and put a few drops of either of these mixtures on the tongue from time to time.



After recovery, treat the patient as stated under the head "Apparent Death from Drowning."

6. *Apparent Death from Noxious Vapors.*

Remove the body into cool, fresh air; dash water over neck, face, and breast, and treat it, in general, as is recommended under the head "Apparent Death from Drowning."

7. *Apparent Death from a Fall or Blow.*

After the sufferer has been placed in a half-erect position on a bed or couch, put on his tongue a few globules of *Arnica*, and wash the parts which have been hurt by the fall or blow with a solution of the tincture of *Arnica* and cold water (twelve drops of the tincture in half a pint of water); also, give injections of the same mixture. Then examine carefully the patient's condition as regards fractures of bones or other injuries, and do not allow him to be bled, if the pulse is small and quick; but if full and slow, he may be bled, with advantage.

Continue, however, the use of *Arnica*, internally and externally, except after bleeding, when a dose (four glob.) of *China* may be given, if much blood has been taken.

8. *Apparent Death from Violent Mental Emotion.*

See the article on "Fainting, Swooning," page 24.

## CHAPTER II.

### CASUAL DISEASES.

#### I. EXTERNAL INJURIES.

As this subject commands such a wide sphere of action, including as it were the whole art of surgery, it would be impossible to do it justice in a treatise like this, except we were able to condense it in a lucid manner, showing the application of the few medicinal agents and mechanical appliances, and pointing at those principles, on the strength of which they are used. Thus, every one can easily be prepared to do, if not all, what could be done right; at least, not to do wrong, in cases where circumstances require him to do something.

In the term “external injuries,” are comprehended,

1. *Fractures of Bones.*
2. *Dislocations of Joints.*
3. *Wounds and Bruises.*
4. *Sprains and Concussions.*
5. *Burns and Scalds.*
6. *Poisoned Wounds, Stings of Insects, etc.*
7. *Frozen Limbs, etc.*

#### 1. FRACTURES OF BONES.

It is all-important, from the first, to know whether a fracture has taken place, and to what extent. Its immediate reduction is not so necessary, as this can be done,

just as conveniently and better, after the first wound-fever has disappeared.

Put the patient in a comfortable position; move the broken limb or part as carefully as possible, and compare its formation with the corresponding healthy one. If it is possible, without much exertion, to straighten its form, do so, even if temporary extension of the broken parts would be required to bring them together; then put a compress around the broken parts, on top of which place four strips of pasteboard or shingles, which are to be kept in their places by a circular bandage, not fastened too tight. Over the whole pour, from time to time, a mixture of cold water and *Arnica tincture* (twelve drops of the tincture to half a pint of water, well mixed); put the limb in a comfortable posture, by pillows, etc.; keep the fractured parts wet, and give, internally, a drop or two of *Arnica tincture* every six or eight hours.

The attention of a surgeon must be sought, at any rate; but, if he can not be had, a repetition of the above bandaging on the third, sixth, and twelfth day, etc., will be sufficient to insure the healing of the fractured parts.

In fractures of the skull, do nothing more than apply cold water and *Arnica*, as above stated, until a surgeon arrives.

## 2. DISLOCATIONS OF JOINTS.

Compare the dislocated part with the corresponding healthy one, and consider, quietly, without being confused, whether it can easily be brought back, for which purpose a few trials may be made. If, however, this should be too hazardous, desist from any further attempts, but use cold water and *Arnica*, as stated in "Fractures of Bones," and wait for the arrival of a competent surgeon. If bandages are necessary, their use must be to confine the joint for some time in the same position after it becomes set.

### 3. WOUNDS AND BRUISES.

Wounds are either *incisive*, *lacerated*, *contused*, *punctured*, or *gunshot* wounds, according to the various instruments which have inflicted them.

Our object is to heal them as quickly as possible, as circumstances will allow. This is done, by bringing their edges immediately in close contact and keeping them there by adhesive plaster or sutures, which is the quickest mode of healing wounds, particularly those of the *incisive* kind. In some wounds the edges can not be brought together, because the flesh is lacerated or bruised. Cases of this kind have to heal by means of suppuration, by which healthy granulation is produced, gluing as it were the parts together. This process applies to all the other kinds of wounds, including the *punctured* and *gunshot* wounds. One important consideration must always be kept in view: never to allow the surface of a wound to heal or close before it is certain that its deeper parts have firmly adhered. This can be prevented by introducing lint to the bottom of the wound, renewing it as often as necessary, at least once a day, and cleaning it with lukewarm water.

The first thing to be done, in attending to a wound, is to clean it, by cold water, from all foreign bodies which may be in it. Splinters must be extracted carefully, and the hemorrhage stopped, which, in most cases, ceases after the application of cold water mixed with a few drops of *Arnica tincture* (see "Fractures"); if it does not stop, however, or if the blood gushes out of the wound in jets, and is of a bright red color (arterial blood), try to compress the wound with lint dipped in arnica-water, overlaid with a sponge; and if this does not succeed, compress with the fingers the artery above the wound (on a place nearer to the heart), which can be found on the inside of a limb,

indicated by the beating of the artery ; if this can not be done, bind around the whole limb, above the wound, a handkerchief, as tight as is necessary to stay the hemorrhage. When this is done, dress the wounds, either with adhesive plaster (if they are cut, or incisive wounds), or loosely with lint kept constantly wet by cold water mixed with *Arnica tincture* (see "Fractures.") Give, internally, a few drops of *Arnica tincture* in water, and place the patient in a comfortable position.

Contused and bruised wounds allow of a certain degree of compression, by adhesive plaster, after the inflammation has left, to make the edges adhere more closely, which can be done afterward.

Another important consideration is, to keep healing wounds well cleansed, by means of allowing the secreted matter to escape easily in the lint spread over the wound to keep off the air. Every day a suppurating wound ought to be cleansed and bathed freely in lukewarm water. If a wound inflames, becomes hot, swollen, and painful, put a warm bread-and-milk poultice over it, until it feels easier ; internally, give *Chamomile*, *Belladonna*, and *Hepar*, alternately, every two hours a dose (four glob.)

If a *wound fever* sets in, give *Aconite*, *Chamomile*, and *Belladonna*, in alternation, in the same manner.

If *lockjaw* should appear, see page 22.

N. B. Although *Arnica tincture* is recommended above in *all* kinds of wounds as the proper outward application, yet *Calendula tincture* is best after *Arnica*, in all contused and lacerated wounds.

If a wound suppurates too freely, or secretes unhealthy matter, give *Hepar* and *Silicea*, in alternation, every evening a dose (four glob.), until better.

#### 4. SPRAINS AND CONCUSSIONS.

These are caused by falls, lifting of heavy weights, jerks, false steps, etc. Bruises, wounds, or fractures, which may attend them, have to be treated first. If the patient is free from them, however, put him at once in a comfortable position, and use, externally, a mixture of cold water and *Arnica tincture*, particularly when the parts are *black and blue (bloodshot)*; after it, *Rhus* in solution, when the *joints, membranes, and tendons*, are more affected. This latter remedy is also suitable for the bad consequences of lifting too heavy weights (*strains*.)

#### 5. BURNS AND SCALDS.

Slight degrees of these injuries heal quickly by holding the scalded parts to the fire (if that is possible), or fomenting them with warm alcohol, or covering them with a plaster of Castile soap. But burns and scalds increase in importance and danger, in proportion to the depth and extent of the injury, which, if very great, threatens the life of the patient. If more than one-half of the surface of the skin is deeply scalded, the sufferer will seldom recover.

A great many remedies are recommended for burns and scalds, but their usefulness must be determined according to the quickness with which they can be procured, their easy applicability, and, lastly, their specific effect on the burns, as such.

TREATMENT.—The best remedy in slight cases is already mentioned above. In severer cases, and even the worst, use *Castile soap*, scraped and mixed with water to a thick lather; spread it on strips of linen or cotton cloth, to the thickness of the sixth of an inch; then spread it over the wound, taking care to cover with soap every burned part, and keep it moist for a while with cold water, which may



be dropped on top, very sparingly, from time to time. Let these plasters remain until they drop off themselves ; or, if matter forms underneath, remove them and dress the wounds with a salve made of equal parts of sweet oil or linseed oil and lime-water, which is a very good remedy in the beginning, if it only could be had quick enough.

Another remedy, easy of access, is *raw cotton*, with which the burned parts must be covered, pressing it lightly on the wounds.

If a person has inhaled hot steam, or has burned his throat with hot liquids, give him, of a solution of Castile soap in whisky or alcohol (which can be made very soon, by scraping soap into the liquor and shaking the bottle well), every five or ten minutes, two or three drops in water, lengthening the intervals as the patient gets better.

N. B. Every steamboat ought to be provided with a bottle of spirits of soap, decidedly the best remedy for these internal burns, as it acts specifically on the injured nervous system and composes immediately.

Another remedy is the tincture of *Urtica urens* (*Burning nettle*), of which each homœopathic medicine chest contains a bottle. Apply it in a manner similar to the *Arnica tincture* (twelve drops to a pint of water.) For internal burns, put three drops of it in a teacupful of water, and give, every five or ten minutes, a teaspoonful, until better.

If the burns ulcerate, wash them with a solution of twelve globules of *Caustic* in half a teacupful of water, three times a day, and give, internally, *Hepar* and *Silicea*, alternately, every evening a dose (six glob.) until better.

DIET AND REGIMEN. — The diet must be light, as in fevers ; but, when ulceration takes place, give stimulating diet, beef, etc., even brandy and water. If a *diarrhea* ensues during the healing of the burns, do not disturb it by giving medicine for it (except it be too excessive), as it is a

critical discharge, the intestines supplying, by their action, the interrupted functions of the skin. In dressing the wounds, take care that the patient is not exposed to currents of air; but, if he should have taken cold, in consequence of which the wounds pain very much and become inflamed, red, swollen (a kind of erysipelas in the scalded parts), give *Aconite* and *Belladonna*, in alternation, every two hours a dose, and dress the wounds with dry lint; but if the wounds were already suppurating when this took place, give *Chamomile*, *Belladonna*, and *Hepar*, in the same manner, and dress the wounds with a warm poultice of bread and milk, until suppuration commences again and the fever leaves.

6. POISONED WOUNDS. BITES AND STINGS OF INSECTS—  
*Bees, Spiders, Bugs, Mosquitoes, Snakes, Mad Dogs.*

In injuries from the above named animals, except the two last (which are treated of below), cover the injured part with wet or damp earth, immediately, and then wet it afterward with a mixture of cold water and *Arnica tincture*, (twenty parts to one.) Internally, give *Arnica* and *Camphor*, every three or four hours a dose (four glob.) until better.

In the treatment of wounds inflicted by *venomous snakes* and *mad dogs*, and of their immediate or distant consequences, *hydrophobia*, etc., there exists yet a great deal of uncertainty and diversity of opinion. The old school has at least retired, as it seems, from further investigation on the subject (their last trial was that of using Chloroform and Ether); and it befits homœopathists the more to promote farther investigations, as they alone possess the true means to shorten the labor and discuss the merits of a proposed remedy. As experiences in this class of diseases occur so rarely, the observations and opinions will be slow



in forthcoming. We prefer to give those of Dr. Hering, as follows :

The best remedies against the bites of *venomous serpents, mad dogs*, etc., is the application of *dry heat* AT A DISTANCE. Whatever is at hand at the moment, a red-hot iron or live coal, or even a lighted cigar, must be placed as near the wound as possible, without, however, burning the skin, or causing too sharp pain ; but care must be taken to have another instrument ready in the fire, so as never to allow the heat to lose its intensity. It is essential, also, that the heat should not exercise its influence over too large a surface, but only on the wound and the parts adjacent. If oil or grease can be readily procured, it may be applied round the wound, and this operation should be repeated as often as the skin becomes dry ; *soap*, or even *saliva* may be employed, where oil or grease can not be obtained. Whatever is discharged in any way from the wound ought to be carefully removed. The application of burning heat should be continued in this manner till the patient begins to shiver and to stretch himself ; if this takes place at the end of a few minutes, it will be better to keep up the action of the heat upon the wound for an hour, or until the affections produced by the venom are observed to diminish.

Internal medicines must be judiciously administered at the same time. In the case of a BITE FROM A SERPENT, it will be advisable to take, from time to time, a gulp of *salt and water*, or a pinch of *kitchen salt*, or of *gunpowder*, or else some pieces of *garlic*.

If, notwithstanding this, bad effects manifest themselves, a spoonful of *wine* or *brandy*, administered every two or three minutes, will be the most suitable remedy ; and this should be continued till the sufferings are relieved, and repeated as often as they are renewed.

If the shooting pains are aggravated, and proceed from

the wound towards the heart, and if the wound becomes bluish, marbled, or swollen, with vomiting, vertigo, and fainting, the best medicine is *Arsenic*. It should be administered in a dose of four globules in a teaspoonful of water; and if, after this has been taken, the sufferings are still aggravated, the dose should be repeated at the end of half an hour; but if, on the contrary, the state remains the same, it should not be repeated till the end of two or three hours; if there is an amelioration, a new aggravation must be waited for, and the dose ought not to be repeated before its appearance.

In cases in which *Arsenic* exercises no influence, though repeated several times, recourse must be had to *Belladonna*; *Senna* also frequently proves efficacious.

Against chronic affections arising from the bite of a serpent, *Phosphor. ac.* and *Mercury* will generally be most beneficial.

For the treatment of persons bitten by a *mad dog*, after the application of dry heat, as directed and described above, see "Hydrophobia," below.

If morbid affections or ulcerations exhibit themselves in consequence of a bite from a rabid *man* or *animal*, *hydrophobine*, administered in homœopathic doses, will often render essential service.

For wounds that are envenomed by the introduction of animal substances in a state of putrefaction, or of pus from the ulcer of a diseased man or animal, *Arsenic* is generally the best medicine.

Lastly, as a PREVENTIVE against bad effects, when obliged to touch morbid animal substances, envenomed wounds, or ulcers of men and animals under the influence of contagious diseases, the best method that can be pursued is the application of *dry burning heat at a distance*. To effect this purpose, it will be sufficient to expose the hands for five or

ten minutes to the greatest heat that can be borne; and after this, it will be proper to wash them with soap.

The use of *Chlorine* and muriatic acid, in similar cases, is well known.

**HYDROPHOBIA.**—Apply *distant heat* to the recent wound, as described under “Poisoned Wounds,” or until shudderings appear; and continue this practice three or four times a day, until the wound is healed, without leaving a colored cicatrix.

At the same time the patient should take, every five or seven days, or as often as the aggravation of the wound requires it, one dose of *Belladonna* or *Lachesis*, or also of hydrophobine, till the cure is completed.

If, at the end of seven or eight days, a small *vesicle* shows itself *under the tongue*, with feverish symptoms, it will be necessary to open it with a lancet or sharp-pointed scissors, and to rinse the mouth with salt and water.

If the raging state has commenced before assistance can be procured for the patient, the medicines that ought to be administered will be according to circumstances, especially *Bellad.* or *Lachesis*; or else, again, *Cantharides*, *Hyoscyamus*, *Mercury*, or also *Stramon.* or *Veratrum*.

## 7. FROZEN LIMBS, ETC.

If limbs, ears, fingers, nose, etc., are frost-bitten, rub them with snow, or put them in the coldest water, and then, by degrees, let the water be warmer, until a natural feeling returns again. If the person is in a state of insensibility from being frozen APPARENTLY TO DEATH, undress him carefully (cut the clothes off from him) and cover him all over with snow, leaving the mouth and nostrils free. As the snow melts, renew it. If no snow can be had, put him into a bath of water, cold as ice, in which he may lie from ten to fifteen minutes; afterward, continue to rub with

snow, or cover him with bags containing pounded ice. If, in this way, the body has thawed by degrees, and the rigidity of the muscles relaxed, dry the body carefully, and place it in a cold bed in a cool room, and begin to rub, under the bedclothes with the warm hands only, all over. In this way continue for hours.

If signs of life show themselves, inject a little camphor and water; also, put a drop of spirits of camphor on the tongue, from time to time.

If more signs of life appear, give warm black coffee (without milk), in injection and by the mouth, until he can take more of the latter.

If severe pains in the whole body appear, give *Carbo veg.* (twelve glob. dissolved in half a teacupful of water), every fifteen or twenty minutes a teaspoonful, in alternation with *Arsenic*, if necessary, until better.

The patient must avoid the heat of fire or of a stove for some time.

## II. POISONS.

Of the poisonous substances which can be introduced into the system, those are the most deleterious and pernicious which have the readiest facility of assimilation; consequently, the animal poisons rank first, then come the vegetable, and, lastly, the mineral poisons.

The animal poisons show their effects immediately on the nervous system, through the blood. Bites of snakes, for instance, while introduced into the stomach, are of no dangerous consequence. The vegetable poisons must be carried into the blood by digestion before their effects become dangerous. The mineral poisons kill only by corroding the surface, with which they come in contact, creating inflammation, which has a great tendency to pass quickly over

into gangrene or mortification;\* their bad effects on the system, afterward, are more lasting.

In cases of poisoning, the first thing to be done is, to *eject the poison as soon as possible* from the system, by provoking vomiting,† or to *neutralize its action* by means of suitable antidotes.

---

\* This peculiarity of the mineral poisons, to kill only by corrosion of the mucous membrane of the intestines, renders their use as remedial agents in diseases quite safe and expedient, as they can easily be divested of this poisoning or corroding quality, by reducing their crude bulk (which alone can corrode or inflame) by trituration and subsequent dilution, without diminishing in the least the curative effects of these minerals. For instance: one grain of *Arsenic*, in its crude state, would certainly kill; yet, if triturated with one hundred grains of sugar of milk, one grain of this mixture would have lost all corroding or poisonous power, retaining, at the same time, all the curative quality, and this in a more developed form. In this way, Homœopathy can avail itself of all the healing effects of the most deadly poisons, with the greatest ease and safety; while Allopathy stands trembling and in fear before these dreadful agents, not knowing how to tame their useless fury, or to harness these wild medicinal steeds to the car of Esculapius.

† That, in such cases, homœopathists make use of emetics and other violent means to eject poisons from the stomach, is not a proof that they resort to allopathic means, as allopathists, quite sophistically, would make people believe. It only shows that, in cases of this kind, the homœopathic law can not come into operation until these substances are removed, which act, as it were, as external injuries all the time, while they are buried in the stomach. In the same way, Surgery might be pronounced insufficient in its doings, because a surgeon can not heal a wound as long as the knife continues to lacerate and even enlarge the wound.

It is evident that external injurious substances (and such are all poisons) must be removed, just as a burning coal or a stinging blister, before the injuries done to the system can be repaired by the proper homœopathic agents; and this is done by Homœopathy more safely and quickly than by Allopathy, as the knowledge of antidotes is further advanced by the former, which is of the utmost importance in curing the after-effects of the poison.

To produce *vomiting*, the following means may be resorted to, in preference to the so called *emetics*, which the allopathic school has hitherto employed :

1. *Tepid water*, in large quantities and often repeated.
2. *Snuff* or *mustard*, mixed with *salt*, on the tongue.
3. *Tickling* of the throat with a feather or fine straw ; or, lastly,
4. *Injections* of tobacco smoke into the anus through a pipe-stem.

To *neutralize* or *mitigate* the injurious effects of poison, the following means may be used as each case requires, which will be shown below under the heading of each poison named.

1. *White of egg*, dissolved in water, and thus drank in large quantities, is of the greatest efficacy in poisoning by *metallic* substances, particularly *Corrosive sublimate*, *Verdigris*, *Tin*, *Arsenic*, *Lead*, *Mercury*, *Sulphuric acid*, particularly when there are violent pains in the stomach and bowels, with or without diarrhea.

In all cases in which the poison imbibed is unknown, the white of an egg is first given when violent pains in the abdomen are present.

2. *Soap* (*white household soap*), one part dissolved in four parts of hot water, is given in all cases where the white of an egg, when indicated, was insufficient.

In cases of poisoning by *alkaline* substances, it must be omitted as injurious.

3. *Vinegar* (made from wine or cider) diluted with water, as a draught or in injections, alternately with slimy things, is the principal remedy against *alkaline* substances, but is very injurious in cases of poisoning by minerals (the very opposite of *soap-water*.)

4. *Sugar*, diluted and given in water, can be administered in cases of poisoning by *mineral*, as well as *alkaline*



substances, and is, therefore, one of the most valuable remedies in domestic practice, as it can be most easily procured.

5. *Milk* and *Olive oil* are substances not so well calculated to envelope poisons in the stomach, as slimy, mucilaginous drinks, made of pearled barley, linseed, or rice; yet they may be used with advantage against *alkaline* substances and *corrosive acids*, such as *Nitric* and *Sulphuric acid*.

6. *Mucilaginous* drinks, in draughts and injections, are very useful against *alkaline* poisons, particularly in alternation with vinegar. They are made of barley, rice, linseed, farina.

7. *Coffee* — *strong black coffee* — taken very warm, is the principal remedy for all *narcotic* poisons, such as *Opium*, *Nux vomica*, *Belladonna*, *Stramonium* (Jamestown weed.)

8. *Camphor* is principally efficacious against all *vegetable* poisons of an acrid, *corrosive* nature; also, against *Cantharides* (Spanish fly.)

The following table shows the proper and successive application of the above named remedies in cases of poisoning with the different substances alluded to.

*Table of Antidotes to the most powerful Poisons.*

POISONS.	ANTIDOTES.
I. ANIMAL POISONS.	
a. <i>Cantharides</i> (Spanish fly.)	<i>Camphor</i> , internally or by smelling; externally, use white of eggs and gruels.
b. <i>Poisonous fish, clams, mussels, etc.</i>	<i>Charcoal, Sugar and water, Black coffee, Camphor.</i> — For eruptions and swelling of the face, <i>Bellad.</i> , <i>Rhus</i> .



POISONS.	ANTIDOTES.
c. <i>Poison of fat</i> in half putrified meat ( <i>Sebacic acid</i> ), as it appears often in cheese, sausages, etc.,	<i>Vinegar and water</i> , internally and externally as a gargle, <i>Lemon juice</i> , <i>Black coffee</i> , <i>Black tea</i> . For the remaining symptoms, <i>Bryonia</i> , <i>Phosphoric acid</i> , <i>Creosote</i> .
II. VEGETABLE POISONS, in general,	Require mostly <i>Camphor</i> , by smelling, and <i>Black coffee</i> in a drink.
a. <i>Opium</i> ( <i>Laudanum</i> , seeds of <i>Poppy</i> ), <i>Stramonium</i> ( <i>Gymson weed</i> ), <i>Nux vom.</i> , <i>Tobacco</i> ,	<i>Narcotics</i> , require particularly <i>Black coffee</i> and <i>Vinegar</i> diluted with water.
b. <i>Gamboge</i> , <i>Euphorbia</i> , and all other burning, corrosive vegetable substances which produce violent pains,	Require mostly <i>Soap-water</i> and <i>Milk</i> .
c. <i>Sumach</i> ( <i>Poison vine</i> ) . . .	No external application, but, internally, <i>Belladon.</i> , <i>Bryonia</i> , <i>Rhus</i> .
d. <i>Camphor</i> and <i>Saffron</i> , . . .	<i>Black coffee</i> , until the patient vomits; afterward, <i>Opium</i> .
e. <i>Spirits of Turpentine</i> , . . .	<i>Opium</i> , <i>Belladonna</i> , <i>Bryonia</i> .
III. MINERAL POISONS.	
a. ACIDS— <i>Prussic</i> or <i>mineral acids</i> , . . . . . <i>Sulphuric</i> , <i>Muriatic</i> , <i>Nitric</i> , <i>Phosphoric acid</i> .	<i>Spirits of Hartshorn</i> . <i>Tepid Soapsuds</i> ; <i>Magnesia</i> ; <i>Chalk</i> , powdered and mixed with water; <i>Wood ashes</i> mixed with water; <i>Potash</i> or <i>Soda</i> .
b. ALKALINE SUBSTANCES— <i>Pot</i> and <i>Pearl ashes</i> , <i>Lapis infernalis</i> , <i>Salt</i> , and <i>Oil of tartar</i> ,	<i>Vinegar</i> , <i>Lemon juice</i> , and other acids; <i>sour milk</i> ; mucilaginous drinks and injections.
c. METALLIC SUBSTANCES— <i>Arsenic</i> ,	<i>Soapsuds</i> ; <i>white of eggs</i> with water; <i>Sugar-water</i> ; <i>Milk</i> ; <i>Rust of Iron</i> .

POISONS.	ANTIDOTES.
<i>Corrosive sublimate, Copper, Verdigris, . . . . .</i>	<i>White of eggs in water; Sugar-water; Milk; Starch from Wheat flour.</i>
<i>Lead, . . . . .</i>	<i>Epsom salt; Glauber salt.</i>
<i>Lunar caustic, . . . . .</i>	<i>Common salt, dissolved in water.</i>
<i>Tin, . . . . .</i>	<i>Sugar, white of eggs and milk.</i>

### III. MEDICINAL DISEASES.

The abuse of medicines, prescribed often by allopathic physicians, and taken in large quantities, for a long time, produces, frequently, artificial diseases, well-known under the name "Medicinal Diseases." The most obstinate and deleterious of these maladies are those produced by vegetable drugs, as their readiness to assimilate with the human system is far greater than that of minerals. From this it is evident how foolish or deceptive the conduct of those physicians is, who pretend to use in their practice only "vegetable medicines," as if this kingdom of nature did not contain the most noxious and violent poisons.

In the following table, the drugs are put together with their antidotes, which may be given as the indications mentioned may require. In administering them, it must be understood, that all these cases are more of a chronic nature; consequently, the repetition of doses is not required to be made so often, say from half a day to two, three, or more days, until improvement takes place, or another remedy has to be selected.

Medicinal diseases are not so easily cured as natural ones, because the reactive force of nature (so important in the treatment of diseases in general) can not altogether be relied on in cases of this kind.

*List of Drugs causing Diseases, and their Remedies.*

DRUGS.	REMEDIES.
1. <i>Opium, Laudanum, Paregoric.</i>	<i>Coffea, Ipecac., Belladonna, Nux vomica, Mercury.</i>
2. <i>Mercury, Calomel.</i>	<i>Hepar sulph., Nitric acid.</i> Against salivation and sore mouth, <i>Hepar, Nitric acid, Carbo veg., Sulphur.</i> Against ulcers in the throat, <i>Carbo veg., Nitric acid, Hepar, Belladonna, Lachesis, Sulphur.</i> Against nervous weakness, <i>China, Hepar, Lachesis.</i> Against nervous excitability, <i>Chamomile, Pulsatilla, Carbo veg.</i> Against sensitiveness to changes of the weather, <i>Carbo veg., China.</i> Against rheumatic pains, <i>Neuralgia; Carbo veg., China, Lachesis, Sulphur.</i> Against ulcerations and swellings of glands, <i>Buboes; Carbo veg., Hepar, Lachesis, Nitric acid, Sulph., Thuja.</i> Against dropsical affections, <i>China, Dulcamara, Hellebor., Sulphur.</i>
3. <i>Quinine, Peruvian bark.</i>	Against Rheumatic pains, <i>Arn., Puls.</i> — Dropsical swellings, <i>Arsenic, Ferrum.</i> —Congestions to the head and bowels, <i>Bellad., Mercury.</i> — <i>Fever, intermittent, Ipecac., Pulsat., Carbo veg., Arsenic.</i>
4. <i>Sulphur.</i>	<i>Puls., Merc., Chin., Sep.</i>
5. <i>Magnesia.</i>	<i>Colocynth, Ars., Rheum., Puls.</i>
6. <i>Arsenic. (Fowler's solution.)</i>	<i>China, Ferrum, Ipecac., Veratrum, Nux vomica.</i>
7. <i>Lead (used in ointments and in washes, as sugar of lead water, etc.</i>	<i>Opium, Belladonna, Platina, Nux vomica.</i>
8. <i>Iodine.</i>	<i>Bell., Phos., Hep., Spong., Chin., Ars.</i>
9. <i>Rhubarb.</i>	<i>Chamomile, Colocynth, Mercury.</i>
10. <i>Chamomile.</i>	<i>Aconite, Ignatia, Pulsatilla.</i>
11. <i>Cantharides (Spanish fly.)</i>	<i>Camphor, Aconite, Pulsatilla.</i>
12. <i>Asafœtida.</i>	<i>China, Mercury, Pulsatilla, Caustic.</i>

## CHAPTER III.

### CUTANEOUS DISEASES.

THE skin, as the external covering of the whole body, is, on account of its exposure and extent, liable to many and various diseases—from the slightest redness, or rash, to the most inveterate itch and leprosy. The importance of its preservation is, therefore, obvious. We divide the diseases of the skin into,

1. *Acute Eruptions, Eruptive Fevers, etc.*
2. *Chronic Eruptions and Ulcers.*

#### 1. ERUPTIVE FEVERS.

##### *Rash.*

DIAGNOSIS.—Frequently, after taking cold, particularly with a disordered stomach, persons are troubled with a rash on different parts of the body, which appears in red spots, of the size of a pin-head, scarcely visible; however, more so in a warm bed; with intolerable itching, particularly at night, preceded by shivering toward evening, disturbing sleep, and followed by feverishness during the night.

In such cases, the common practice heretofore has been, and is yet, where homœopathy is not known, to put scorched flour on the parts affected. This is not, however, without danger, and ought not to be resorted to immediately. It would be better first to give some of the following remedies,

which will relieve, without driving the disease to internal parts.

TREATMENT.—If the itching is intolerable, with either shivering or heat, restlessness, sleeplessness, give *Aconite*, every hour one dose (four glob.) If not better within two or three hours, give *Chamomile*, every two hours a dose (four glob.) If perspiration follows this treatment, keep it up as long as necessary, and give, the next day, a dose of *Sulphur* (four glob.)

DIET AND REGIMEN.—The diet ought to be very light—farinaceous substances, gruels, toasted bread and toast-water—but no meat, or soups of meat. The patient must be kept in a dry, warm room.

#### *Nettle Rash—Hives.*

DIAGNOSIS.—A disease similar to the former in origin and appearance, only that the red spots, with a whitish tinge, appear in groups, somewhat elevated, and itch and prick intolerably. They resemble that eruption caused by contact with nettles. Hence the name, nettle rash. Sometimes the body is covered with these spots, and then the patient feels sick, has no appetite, with a feeling of fullness in the pit of the stomach.

TREATMENT.—When caused by indigestion, particularly from rich and fat food, give *Pulsatilla*, morning and evening (four glob.) When caused by taking cold, and accompanied with diarrhea, bitter taste in the mouth, and slimy coated tongue, give *Dulcamara*, morning and evening a dose (four glob.) When caused by taking cold in damp, wet weather, exhibiting itself with pains in the limbs, shivering, and headache, give *Bryonia* and *Rhus*, alternately, every three or four hours a dose (four glob.), followed, in twenty-four hours, by a dose (four glob.) of *Sulphur*, if necessary. If not better the third day, give, evening and morning, a

dose (four glob.) of *Carbo veg.* When these remedies are of no avail, or there is a burning sensation, with itching, great restlessness, and even an uneasy feeling, give two doses of *Arsenic*, every two or three hours one (four glob.); and, the next day, of a solution of *Urtica* (one drop in six tablespoonfuls of water), every three hours a teaspoonful.

Sometimes this rash strikes in suddenly, when the patient feels oppressed, sick at the stomach, and weak. In such a case, give, immediately, *Ipecac.* (four glob.); in an hour afterward, *Bryonia* (four glob.), and, after that, in two hours, if not better, *Arsenic* (four globules.) At the same time, cover the patient well, to produce perspiration, which will also be promoted by warm drinks, particularly warm lemonade, if he has no diarrhea at the time.

If a person is troubled with this disease for a longer time than common, or liable to it on the slightest occasions, let him take *Calcarea*, every four days a dose (four glob.), in the evening, and, if not better in four or six weeks, *Lycopodium*, in the same manner.

DIET must be simple; no meat nor heating drinks; nothing but water, black tea, gruels, dry toast, baked apples, and stale bread.

### *Erysipelas, or St. Anthony's Fire.*

DIAGNOSIS. — It is characteristic of this disease, that it appears, suddenly, on different parts of the body, with heat, redness, swelling, tingling, and other painful sensations; in severer cases, the heat becomes intense on the parts affected, the surface almost shines, the pains become burning and shooting, and the skin rises in blisters filled with water (*erysipelas bullosum*.)

In such cases, an intense fever is present, with its concomitant gastric nervous symptoms, such as a quick, full, wiry pulse; high fever preceded by shivering or chill;



tongue coated, white or dry brown; great thirst; pains all over the body; scanty and deep-colored urine; intense headache, with sleepiness or wakefulness; very sensitive to noise and light.

The nearer to the brain this disease appears, as on the face, the ears, or the scalp of the head, the severer the symptoms of the head will be, and, consequently, the more dangerous is the attack, so that delirium often ensues.

In the so-called "wandering erysipelas" (*erysipelas erraticum*), the morbid spot disappears from one place, changing into a yellowish hue, to reappear again on another, in the form of redness and heat, so that it frequently wanders from the left side of the face, around the ear and neck, to the right side, before the whole disease disappears. In such cases, the face is very much swollen and disfigured, closing, for a time, the eyes, nose, and ears. If the scalp of the head is attacked, the hair often falls off afterward.

CAUSES.—In most cases, exposure to cold immediately after the system has been heated or excited by over-exertion or mental emotions, particularly with a previous derangement of the stomach; or, in females, during menstruation. Some persons are predisposed to it. Certain kinds of food, at certain seasons, will also produce it, such as lobsters, oysters, and other shell-fish.

TREATMENT.—All outward applications ought to be avoided as dangerous—at any rate, all which are greasy and damp. Erysipelas does not bear anything wet. Salves and ointments are very dangerous. If the itching and burning is too violent, we may mitigate it a little, by applying powdered starch made of wheat flour, but not until after having given some of the following medicines internally. The internal remedies are the most important.

If the fever is very high, pulse full, skin dry, give first a few doses of *Aconite*, one every two hours (four glob.),



followed by *Belladonna*, two doses, one every four hours (four glob.), which, in less severe attacks of this disease, is alone sufficient. In such cases, the patient improves after twelve or sixteen hours, and no more medicine is needed; he simply remains at home a few days, until there is no more danger of taking cold.

In severer cases, however, when after the above treatment the symptoms do not abate, give *Lachesis*, two doses, every three hours one (four glob.), particularly if the patient complains of dryness in the throat and pain in swallowing; also, coughs without raising.

If no better after this, or if there be great sensitiveness to noise and light; the rash expanding in radiation; shining redness on its surface, which is very sensitive to the touch; sometimes raised in blisters; give *Belladonna* and *Rhus tox.* in solution (twelve glob. in half a teacupful of water, each cup having its separate teaspoon), every two hours a teaspoonful, alternately, until four teaspoonfuls of each are taken. Then wait twelve or more hours. If the patient is then very drowsy, give a dose of *Opium* (four glob.); or, if very wakeful and restless, give *Coffea* <sup>cc.</sup> and *Bellad.* <sup>cc.</sup>, alternately, every hour a dose (six glob.) for three or four hours, and wait again twelve hours; after which, repeat *Belladonna* and *Rhus tox.*, in solution as above. If the symptoms are yet the same, give two doses of *Hepar*, every three hours one (four glob.), when the skin looks less shining and inflamed, but heat, pain, and swelling, are the same.

The above treatment suits severe cases of erysipelas of the face, head, or ears, with high fever, and, also, the vesicular form, where it appears in blisters, and the "wandering erysipelas." In the latter, however, do not expect so rapid a termination of the disease. If erysipelas appears

on the joints, give *Bryonia*, three doses, every six hours one (four glob.), alternating with *Rhus*, and followed by *Sulphur* after thirty-six or forty-eight hours, one dose (four glob.), if necessary.

If it appears on the scrotum, *Arsenic* is necessary, administered in the same manner as *Bryonia*. If the vesicles become gangrenous or of a dark color, the patient is weak, or black diarrhea sets in, *Arsenic* and *Carbo veg.* ought to be given alternately, every three hours one dose (four glob.) until a homœopathic physician takes charge of the patient.

Sometimes a swelling of the affected parts, showing the pressure of the finger, combines itself with erysipelas (*erysipelas œdematodes*.) This requires *Rhus tox.*, every evening a dose (four glob.) for three or four days.

If erysipelas terminates in ulceration of the parts affected, give *Rhus* and *Sulphur* alternately, every evening a dose (four glob.) for four days, and, afterward, a dose of *Silicea* for two evenings, and wash the ulcer three or four times a day with lukewarm water, in half a teacupful of which twelve globules of *Silicea* are well dissolved.

The disposition to this disease will be taken away by the alternate use of *Rhus* and *Graphites*, every four days a dose (four glob.), if continued for four or six weeks. Beside this, however, change the mode of living, if it was too luxurious; avoid highly seasoned or salted food, but drink freely of cold water, and wash and bathe frequently and systematically.

DIET.—During this disease the diet should be as in fevers, very light; dry toast, gruels, and black tea; warm lemonade is very good and refreshing, if no diarrhea prevails; stewed prunes may also be allowed, particularly in convalescence.

If, at any time during this disease, the bowels are con-

fined, even for four or six days, it will not exert, in the least, a detrimental effect upon the patient ; on the contrary, it is better than the opposite state.

Avoid taking cold during the convalescence, as it is often followed by dangerous results, dropsies, etc. If, at any time during the attack, the erysipelas suddenly disappears, and the brain, in consequence, becomes affected, which shows itself in drowsiness, difficulty of breathing, and spasmodic twitchings, give *Cuprum*, in solution (twelve glob. in half a teacupful of water), every half hour a dose, until a homœopathic physician takes charge of the patient.

Before we leave this subject let us here remark, that erysipelas, as such, is *no infectious* disease whatever, and has lost its dread entirely, since homœopathic remedies have been used in the treatment of it.

#### *Eruptive Fevers with an Infectious Epidemic Character.*

We now come to a class of eruptive diseases which appear, generally, as epidemics, and are infectious. These are *measles*, *French measles*, *scarlet rash*, *scarlet fever*, *chicken pox*, *varioloïds* and *small pox*. Their epidemic appearance depends upon the general laws of disease growing out of the changing conditions of the seasons and atmosphere. As a general rule, we can say that persons are attacked only once in their lifetime with these diseases, though some few exceptions may happen. Their contagiousness does not commence, as is generally believed, during the fever period, or when the eruption first appears ; but *only after* the pustules are well filled, or drying up, that is, from the seventh day onward, to three or four weeks, as long as the system throws out through the skin the infectious matter. Before the seventh day of either of the above diseases, no infection can take place, for the same reason as no vaccination is possible out of a pustule before that time. The

infecting matter must be first fully ripe in the system before it can affect another; be, therefore, rather more careful during the convalescence of a patient, or after his death, (when the contagion is in its highest state of perfection), than in the beginning of his sickness. This is the case as well in measles as in scarlet fever, varioloids and smallpox.

*Measles. (Morbilli.)*

In adults, this disease generally assumes a severer character than in children, because it developes in them more readily those constitutional germs of diseases which often lie hidden in the system a long time, such as consumption, sore eyes, etc. In children, its attacks are rarely dangerous, except when other causes render it complicated, or it is badly treated.

DIAGNOSIS.—For the first three or four days the patient seems to have taken a cold; he has a dry, hoarse cough, and sometimes a sore throat; his eyes water constantly, and can not bear the light; these symptoms and more or less feverishness (the catarrhal fever) precede and accompany the eruption of the measles, which appear in small red spots of irregular shape and size, (the skin between them preserving its natural color), the first day on the temples, face and neck, the second day, on the breast, or stomach and back, and the third day on the arms and limbs. In mild cases they disappear after the fifth or sixth day in bran-like scales.

TREATMENT.—*Aconite* and *Pulsatilla* are the principal remedies, and are regarded almost as specifics, which, in milder cases, without complication, they prove to be.

ADMINISTRATION.—Dissolve of each remedy twelve globules in two teacups half full of water, each (put to each a separate spoon), and give alternately every two, three or six hours, a teaspoonful (children under two years, half a

teaspoonful), beginning with *Aconite*; and if the patient remains, or becomes more comfortable, lengthen the intervals still more. If, at any time, there is great restlessness and sleeplessness, introduce a dose of *Coffea*,<sup>cc.</sup> repeating it as often as necessary, particularly in the night.

If the cough is very dry and hoarse (resembling croup), give a dose (four globules) of *Hepar* in alternation with *Coffea* (if the patient is very much agitated and irritable), at intervals of one or two hours.

If, however, the cough excites severe pain in the upper part of the breast and sides (complication with bronchitis and pneumonia), *Aconite* and *Bryonia* (dissolved in water and administered the same as *Aconite* and *Pulsatilla*) are necessary.

If the measles strike in at once, and look pale, or if sickness at the stomach and feebleness ensue, give *Ipecac.* and *Bryonia* alternately every hour or half hour a dose, (four glob.) and if the skin looks bluish brown, give *Arsenic* and *Pulsatilla* in alternation every hour or two hours a dose, until better. If the head is affected in children, give *Cuprum* every half hour a dose, (three glob.) twice or three times; and if not better in three hours, give *Belladonna* and *Hellebor.* in solution (twelve glob. of each to half a teacupful of water) every hour, alternately, a teaspoonful until better.

*Disorders consequent upon Measles.*—If sensitiveness to light continues too long after the eruption is out, give *Belladonna*, morning and evening one dose (four glob.); this same remedy, applied similarly, suits, if the head is affected (*congestion of the head*), in alternation with *Stramonium*.

Constipated, or not too open bowels, are not injurious during or after this disease; if a mucus diarrhea ensues

give *China*, *Mercury*, *Pulsatilla*, at intervals of six hours, until better. If not better, give one or two doses of *Sulphur*—every six hours one dose, (four glob.). If the diarrhea is watery, and connected with typhoid symptoms, loss of consciousness, dry tongue, give *Arsenic* and *Phosphorus* in the same manner.

If earache ensues, give *Pulsatilla*, every two or three hours a dose; if this is accompanied by running of the ear, give after *Pulsatilla* a dose (four glob.) of *Sulphur* on two successive evenings, and if not better in two or three days, give *Carbo veg.* in the same manner.

If the glands below and in front of the ear swell, (*parotitis*, or mumps), give *Arnica* and *Rhus* every four or six hours a dose (four glob.), till better.

DIET must be light—as gruels, water (children's milk and water), toasted bread; and only gradually more nourishing.

N. B. Do not allow the children to go out of the room too early, or even to go to the door or window, or play on the floor; we can not be too careful to avoid taking cold.

As a preventive of measles, give *Aconite* and *Pulsatilla* every third day a dose (four glob.), alternately, during the epidemic; it may prevent an attack, and will certainly mitigate its severity.

#### *French Measles.*

In this disease, which resembles the former, but is milder in every respect, it is only necessary to mitigate the fever and restlessness, if there is any, with alternate doses (four glob.) of *Coffea* and *Aconite*, one every two, three, or six hours, after which, the eruption soon disappears.

DIET AND REGIMEN the same as in measles, but when complicated with *catarrhalic* symptoms, give a dose or two (four glob.) of *Pulsatilla*; and when the head is congested,



give a couple of doses of *Belladonna* in alternation with *Aconite*.

### *Scarlet Rash.*

This disease is different from scarlet fever, in as far as the redness of the eruption is darker, and the finger, on slight pressure, leaves no white imprint, besides there are numerous small granular elevations felt under the skin, sometimes distinctly seen. There is another distinctive difference from scarlet fever; this latter appears, invariably, first on the face, next on the body, and lastly on the extremities. Such a regularity is not found in scarlet rash—it may appear irregularly or at once over the whole body. It is frequently and mostly seen in combination with scarlet fever, which is known by its smooth redness on the surface.

TREATMENT. — If it appears alone, it is not a dangerous disease, and yields easily to a few doses of *Aconite*, every three or four hours one dose (four glob.), and in alternation with *Coffea*, if there is great restlessness and irritability.

If it is combined with scarlet fever, the symptoms sometimes become very severe; see, for their treatment, the next article, “Scarlet Fever.” In case it disappears suddenly, *Ipecac.* and *Bryonia* should be given every half hour, or hour, a dose (four glob.), twice or three times, at the same time covering the patient well; but when symptoms of the head appear, with drowsiness, give *Opium* (four glob.), or, if the patient starts on closing the eyes, *Belladonna* (four glob.), once or twice in four hours.

DIET the same as in measles, and the same caution as regards taking cold afterward.

N. B. This disease does not strictly *prevent* patients from having the real scarlet fever afterward, although it often diminishes their liability to it, as I have had occasion to



observe in many instances. It bears the same relationship to scarlet fever as French measles (or measles-rash, as we may call it) does to ordinary measles.

*Scarlet Fever.*

A well known scourge to the world of children, and dreaded by the allopathic physician more than any other. It is not the least triumph of the glorious Hahnemann to have found the true specific against this disease, diminishing its terrors, and furnishing a remedy, which the allopathists have adopted to a great extent.

DIAGNOSIS.—After a fever of more or less severity, characterized by a very quick pulse, and sometimes accompanied with sore throat, headache, thirst, and stupor, or uncommon liveliness, the eruption appears in the form of bright red (scarlet) blotches indefinitely marked, on which, when pressed by the finger, a white spot remains, which soon becomes red again, from the centre to the circumference. If it is not combined with scarlet rash, no grains can be found in the skin by rubbing over it with the hand; it appears, the first day, on the face and neck, the second on the breast and region of the stomach, and the third day, on the arms, hands, and limbs. But more frequently it appears in combination with scarlet rash (see this article), in which case it assumes a more dangerous and destructive character.

If this disease exists alone, it generally terminates in five or six days, the skin coming off in large pieces.

TREATMENT.—*Belladonna* is the specific for this disease, which is cured by its use alone, except in complicated cases.

The symptoms requiring its use, are dry burning fever, quick pulse, starting on closing the eyes for a few minutes, great thirst, dry, red, or whitish coated tongue, urine scanty and highly colored, scarlet eruptions on the face, or over

the entire body. Sometimes this external redness does not appear, but instead of it the throat is more or less affected, in which case the tongue has always a reddish gloss. Yet, even this wandering of the scarlet from the outside skin to the mucus membrane of the throat, does not materially increase the danger of the case, if the glands only do not enlarge too much. Under all these circumstances, as indeed always at first, give *Belladonna* in the following manner: Dissolve twelve globules in half a teacupful of water, and give of this solution every two, three, or four hours, a teaspoonful for four or six times. If the disease gets worse soon after giving *Belladonna*, discontinue its use, or let the patient smell of spirits of camphor a few times and wait, with the application of the *Belladonna*, till the child gets worse again; this, however, will not generally be found necessary.

In this manner continue to use the *Belladonna* for three or four days, until the scaling process commences, when it is only necessary to keep the patients away from the slightest draught; it is by far the safest to keep them in bed closely for four or five days after the fever has left, when a dose of *Sulphur* (four glob.) may be given to complete the cure.

But if the case is a complicated one from its beginning, or becomes so in its progress, other remedies besides *Belladonna* are needed. Cases of this kind ought to be treated by a homœopathic physician; yet, where none is to be had, the following prescription will be found beneficial:

If there is great restlessness and sleeplessness, irritability, whining and tossing about, give *Coffea* and *Belladonna* in solution (twelve glob. in half a teacupful of water), every hour a teaspoonful; or, if there is violent fever, with *dry heat*, full and quick pulse, congestion of the head, occasional delirium or lethargy, with starting when awaking

and dry, short, painful cough, give *Aconite* and *Belladonna* in solution, alternately, in the same manner.

If the patient is better under this treatment during the daytime, but in the night these symptoms appear more or less, particularly restlessness and sleeplessness, give *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup> every hour alternately, (four glob.), until the patient is more composed.

If, with great drowsiness and tossing about, the tongue is very dry, thirst great, with swelling of glands on the neck increasing, skin shining on the face and neck, head thrown backward, and almost constant delirium, give *Rhus* and *Belladonna* in solution the same manner as above, every hour a teaspoonful, alternately, for six or eight hours, followed by *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>; if very restless during the night, as above, this repeated every day for three or four days, secures, in most cases, a favorable issue; perspiration sets in, and then one or two doses of *Sulphur* (four glob.) are necessary to complete the cure.

If this disease is combined with scarlet rash, as it mostly is at present, the symptoms become more severe, the throat is affected inside and the glands swell outside, sometimes to a great extent. We call this the *malignant scarlet fever*. In such cases, when ulceration of the glands commences, indicated by a very offensive smell, and a great quantity of mucous running from the mouth, give *Mercury* (twelve glob, dissolved in half a teacupful of water), every two hours a teaspoonful, for six or eight times alone, or alternating with *Belladonna*, if the patient starts or jerks on closing the eyes, can not swallow liquids easily, has violent thirst, with sensitiveness to light.

If no improvement takes place within twelve hours, give *Sulphur* in solution the same as *Mercury*, twice or three times, and then wait six or eight hours, giving, in the

meanwhile, only an occasional dose (four glob.) of *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, alternately, if the patient is very restless.

If no better at the end of that time, restlessness still increasing, the saliva excessively fetid, with gritting of teeth, give *Arsenic* in solution, in the same manner as *Mercury*, and after that, in the same manner, give *Nitric acid* and *Lycopodium*, particularly when the stupor of the patient increases and lethargy sets in.

If, in this condition, the breathing resembles snoring, with burning heat of the skin, whether dry or covered with perspiration, give *Camphor* and *Opium* in solution (twelve glob. of each, separately dissolved in half a teacupful of water), every half hour a teaspoonful, alternately; give of each three or four times, and after that, if there is no improvement, give *Cuprum*, (twelve glob. in half a teacupful of water), in solution, every hour a teaspoonful.

If, at any time during this disease, a kind of *stranguria* (difficulty of urinating) ensues, give *Cantharides* (four glob.) once or twice, and in alternation with *Coffea*, if there is great restlessness and irritability. If not better, give *Conium* in the same manner.

If croupy symptoms appear, give *Aconite* and *Hepar sulph.* alternately, every hour a dose (four glob.), until better.

*Diseases consequent upon Scarlet Fever and Scarlet Rush.*—*Earache and Ulceration in the Ear.*—For severe pains in the ear, give *Belladonna*, *Pulsatilla*, and *Hepar sulph.*, alternately, every two hours one dose (four glob.) until better.

For the running of the ear (otorrhœa), give *Calcarea* six times, every evening and morning one dose (four glob.), wait three or four days, and if no better, give *Silicea* in the

same manner. For the swelling of the glands below, and in front of the ear, (*mumps*), give *Carbo veg.* and *Rhus*, alternately, every four or six hours a dose (four glob.), until two doses of each are taken.

If a child, after this disease evinces symptoms of *dropsy of the brain*, (head hot, extremities cold, vomiting on moving, with or without diarrhea, sleeps with eyes half open), give *Belladonna* and *Hellebor.* in solution (twelve glob. in half a teacupful of water) every two hours, alternately, a teaspoonful until it gets more lively again and the above symptoms disappear, when a dose, (four glob.), of *Sulphur* will be serviceable.

In case of *dropsical swelling* of the whole body, give first *Hellebor.* and *Belladonna* in the same manner as above, and afterward *Bryonia* and *Hellebor.*, alternately.

In such cases *Arnica*, *Arsenic*, *Phosphor. acid* and *Sulphur* are also useful; commence with *Arnica*, of which two doses may be given, one in the evening and one in the morning (four glob.); and then wait one day until signs of improvement show themselves. If these do not appear, give the other remedies one after another in the same manner. Keep the patient always covered during this treatment.

For soreness of the nose and face, with swelling of the glands under the chin, give first *Mercury* twice, every evening a dose, (four glob.), then wait two or three days, and if necessary, give one after the other, *Hepar sulph.*, *Silicea*, *Sulphur*, and *Calcareæ*.

If the scarlet strikes in suddenly, the eruption assuming a livid bluish hue, and the child becomes drowsy, with hurried breathing, give first *Bryonia* and *Belladonna*, a couple of doses, every half hour a dose, (four glob.), and then if no better give *Cuprum* in solution (twelve glob. in half a teacupful of water) every fifteen or thirty minutes a

teaspoonful. If not better in two hours, continue the same with *Camphor* in alternation, and cover the patient well.

DIET AND REGIMEN the same as in measles; examine closely "N. B." at the end of article on "Measles."

As a *preventive* against scarlet fever, give, during the prevalence of the disease, to every child a dose (four glob.) of *Belladonna*<sup>cc.</sup> to be taken every other evening.

N. B. It is hardly necessary to urge the necessity of the advice and attendance of a homœopathic physician in this disease, if one can possibly be obtained.

### *Chicken Pox.*

DIAGNOSIS.—This eruption is often mistaken for small pox or varioloid, particularly when a small pox epidemic is raging. We will give the differences between them. Before the chicken pox appears, the patient is only sick from twenty-four to forty-eight hours, and then he has, generally, fever and headache, with bilious rheumatic symptoms, without the severe swimming in the head and the back-ache, which never fail to appear before the small pox breaks out. Instead of that, he complains of stranguria and tenesmus, (see glossary), which is not the case in small pox.

The chicken pox appears irregularly on different parts of the body at once, while the small pox and varioloid always appear on the face first. Again, the pustule of the chicken pox appears at once in the form of a bladder, not as in small pox and varioloid, in the form of a point like the head of a pin. When the chicken pox is developed, the pustule has but one cavity, without any inner divisions, containing the clear liquid, and without a dent on the top of it; while the pustule of the small pox contains the liquid in divisions, like an orange, and has a dent on top. The liquid of the chicken pox very seldom becomes turbid or mattery, as is the case in small pox, but dries up in a spongy



crust, without leaving a mark ; the crusts of small pox are hard, and oftentimes leave marks. Chicken pox is not contagious in the same manner as small pox ; it depends, for its propagation, more on individual predisposition, strengthened by atmospheric influence.

TREATMENT.—For the fever, prior to the eruption, give *Aconite* and *Belladonna* every two or three hours, alternately, a dose, (four glob.), until three doses of each are given. If the patient complains of *bilious rheumatic* symptoms, give *Bryonia* and *Rhus*, alternately, every two hours a dose, (four glob.), until three doses of each are given.

If there are *bilious nervous* symptoms, give *Belladonna* and *Rhus*, alternating in the same manner.

If tenesmus is present, give *Mercury*, every two hours one dose, (four glob.) If there is stranguria, give *Cantharides* and *Conium*, alternately, every two hours a dose (four glob.) until three doses of each are given, or the patient is relieved. If the eruption is very considerable, give two doses of *Antimon. crud.* every twelve hours one (four glob.)

DIET AND REGIMEN as in measles.

N. B. In children severe symptoms of the head may appear, for instance, convulsions, (see this article) or the eruption might strike in ; in the latter case, treat it as stated in measles striking in. (See article “Measles.”) Chicken pox may occur more than once in a lifetime.

### *Small Pox. (Variola.)*

This disease has four important stages which we will first describe, afterward we give their treatment.

First. *Febrile Stage*.—This commences, generally, from seven to fourteen days after the exposure to the contagion, with a chill more or less severe, after which, intense fever sets in with severe pains *in the head and small of the back*, aching in the bones and general soreness ; dry, hot skin,



great thirst, cough, oppression in the stomach, sometimes vomiting of bile; light hurts the eyes, with swimming of the head; the mind wanders, is flighty, anxious expression of countenance and great prostration of strength.

Second. *Eruptive Stage*.—After the frebile stage has lasted from forty-eight to seventy-two hours, the eruptive stage commences, by the appearance of small red pimples on the forehead and face of the size of pin heads, after which the severity of the fever symptoms abates gradually. On the first day the eruption appears on the face; the second, on the breast and body; and the third, on the limbs and arms—this stage, therefore, lasts three days.

Third. *Stage of Suppuration*.—The pustules now grow to perfection, as large as a bean cut in two; the liquid inside is contained in cells like an orange, on top is a little dent, and around them a red circle. This stage lasts three days; but as the pustules first appear on the face, then on the breast, and lastly on the extremities, they are already in perfection on the face when they are still filling on the breast and growing on the extremities.

Thus, a wise Providence has divided the burden in three parts, which would otherwise be unbearable.

At the end of this period, the liquid in the pustules is turbid and mattery.

Fourth. *Stage of Desiccation*.—With the eighth or ninth day of the eruption it begins to dry up; some of the pustules burst, and with the formation of scabs desiccation commences and progresses until the tenth day, at which time, in favorable cases, the fever has entirely disappeared, the swelling of the face has diminished, the scabs have fallen off the upper part of the body, leaving marks of a reddish-brown color.

This is the regular and favorable course of the disease where the pustules are not so numerous as to run together;

but where the small pox is *confluent*, the danger of the case and the length of its duration are considerably increased.

TREATMENT. — *Febrile stage*. — During the chill and the first six hours of the fever, give *Aconite*, every two hours a dose (four glob.). If severe headache (congestion), with sensitiveness to light, and delirium, are present, alternate with *Belladonna*, in the same manner.

If, afterward, severe backache ensues, with pains in the bones and general soreness (bilious rheumatic), give *Bryonia* and *Rhus*, alternately, every two or three hours a dose (four glob.) for twelve or sixteen hours. This is generally sufficient.

If, however, there is vomiting, give one dose (four glob.) *Tart. emet.*

If the headache increases to insensibility, stupor ensues, and snoring-like breathing, give one or two doses of *Opium*, every two hours one (four glob.)

If the patient is very restless, sleepless, and irritable, give *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, every hour a dose (four glob.) until better.

TREATMENT OF THE SECOND OR ERUPTIVE STAGE. — If the delirium, which may have lasted up to this time, does not disappear entirely when the eruption appears, or if the eruption does not appear sufficiently or not at all, give *Stramonium*, in solution (twelve glob. in half a teacupful of water), every two hours a teaspoonful, for twelve hours.

If the lungs suffer, at this stage, with a hoarse, slimy cough, give *Tart. emet.*, a few doses, every two hours one, (four glob.); if with great oppression on the chest, give *Ipecac.* in the same manner.

In children, this stage requires particular attention, and *Belladonna*, in alternation with *Stramonium*, should be given, every two hours a teaspoonful of each (ten glob. in half a teacupful of water), until the eruption is out.

If, however, this stage progresses finely, without severe symptoms, give nothing but *Tart. emet.* and *Thuja*, in alternation, every three hours a dose (four glob.), until six doses of each are taken. These remedies have the specific power to mitigate the eruption, from the similar eruption they produce on the healthy.

TREATMENT OF THE THIRD STAGE.—*Suppuration*.—If this stage has a great deal of fever, give *Mercury*, every three hours a dose (four glob.) until four doses are taken, particularly when there is abundant saliva in the mouth. The swelling of the eyes only requires ablutions of warm water and milk.

If, however, in this stage, the skin between the pustules becomes of a livid hue or dark brown, and the pustules themselves watery and flabby (black small-pox), give *Arsenic* in solution (twelve glob. in half a teacupful of water), every two hours a teaspoonful, alternating, in the same manner, with *China*, if diarrhea ensues. After these remedies, a dose of *Sulphur* (four glob.) will be of service.

This treatment is also serviceable, if, in the confluent form, typhoid symptoms occur, with brown, cracked tongue only, give, in the absence of diarrhea, instead of *China*, *Rhus* in alternation with *Arsenic*.

*Striking in of the eruption* happens in this stage more than in any other, and ought to be treated with *Cupr.*, as directed for "Scarlet Fever," (see this article.)

TREATMENT OF THE FOURTH STAGE.—*Desiccation*.—In the beginning of this stage, give a dose (four glob.) of *Sulphur*; if the patient is tolerably comfortable, he does not require anything more than cleansing the skin by frequent ablutions with tepid water, and a careful attention to diet.

If, however, he still has diarrhea, give *Mercury* in alternation with *Sulphur*, every three or four hours a dose (four glob.) until better.

The itching at this period is alleviated by a couple of doses of *Sulph.*, evening and morning one dose (four glob.)

If the patient has reached this period in a typhoid state, he must be treated accordingly. (See "Typhus.")

DIET AND REGIMEN. — The room of the patient should be kept well ventilated, not too warm, and almost dark, during the whole time.

The diet should be rather cold than warm ; for instance, water, ice cream as much as wanted, lemonade, oranges (the latter three not in diarrhea), gruels, dry toast ; also, well-stewed prunes and roasted apples ; animal food is not allowed, even long after convalescence.

Diseases which may occur after small pox, such as consumption, diarrhea, inflammation of the eyes, etc. See under their proper heads.

As a *preventive* against this disease, vaccination is well and favorably known ; for how long a time, however, this may prevent small pox, is yet a matter of speculation. It varies, certainly, in different individuals ; the shortest period may be seven years ; the longest is not known.

I consider vaccination and revaccination from time to time, our duty and a safeguard against the encroachments of this fearful disease upon society in general.

N. B. See introduction to "Eruptive Infectious Fevers," p. 53.

#### *Varioloid, or Modified Small Pox.*

This disease is similar, in all respects, to the former, yet not the same. It is decidedly milder, and requires no mention of a different treatment from that given under "Small Pox." (See this article.)

N. B. A person who is well vaccinated will, when exposed even to small pox, take varioloid only ; and, thus far, vaccination, which is a protection against small pox, does not prevent the varioloid, although one who has had the

small pox will very seldom take the varioloid. The varioloid rarely leaves scars on the skin.

## 2. CHRONIC ERUPTIONS.

To give a detailed description of all the varieties of cutaneous diseases coming under the above head, would be of no advantage for domestic practice, as most of these chronic skin diseases require the most skillful discrimination of an attending physician; for the reason that their causes and character are too deeply connected with the patient's constitution (in each case, perhaps, differently so); and, thus, it is almost impossible to generalize them and their treatment, within the limits of a medical guide-book like this. We will try to describe some of the most familiar ones, advising the reader, at the same time, however, to apply to a homœopathic physician immediately on finding his own treatment insufficient; because all chronic cutaneous affections are very important for the welfare of the system.

### *Irritation of the Skin.—Itching. (Pruritus.)*

A fine rash under the skin, scarcely perceptible, and colorless, produces a very disagreeable and distressing itching, particularly at night, when undressing, or in bed after getting warm. It is caused, mostly, by exposure to the extremes of heat and cold, and appears, consequently, in the height of summer, as well as in winter. Sometimes, the eating of too much fat or greasy food produces it: if so, the diet ought to be changed.

TREATMENT.—Wash, every evening before going to bed, with water and plenty of Castile soap, without drying it off. If this does not relieve within four or six days, try brandy or alcohol, in the same manner (but take care not to come in contact with a light.) If this does not produce good

results, wash with water mixed with *spirits of Camphor*. Internally, administer as follows :

If the irritation is worse in the warmth of the bed or near the fire, and if worse after scratching, give *Pulsatilla*, every evening a dose (six glob.), and, also, if fat food may have caused it.

If it commences after going to bed, and resembles flea bites all over the body, shifting from one part to another, *Ignatia*, given in the same manner, will be of service.

*Mercury* suits, when it continues through the whole night, and after scratching bleeds easily and freely ; in this case alternate with *Sulphur*, every other evening, one dose (four glob.) for a week.

If an intolerable *burning* accompanies the irritation, amounting almost to feverishness, take *Bryonia* and *Rhus*, alternately, every six hours a dose (four glob.) for twenty-four or thirty-six hours; two days after take *Hepar sulph.*, evening and morning, a dose (six glob.) If not better in four or six days after, take *Carbo veg.*, evening and morning, one dose (six glob.)

If it commences when undressing, take *Nux vom.* and *Arsenic*, in alternation. If old people are troubled with it, give *Opium* and *Secale*, alternately, every evening a dose (four glob.) If it renders the patient *very restless*, so that he perspires, take *Colocynth*. If it is caused by summer heat, take *Lachesis* and *Lycopodium*, alternately, every evening a dose (four glob.)

If it is accompanied with *fine stitches*, like needles, take *Thuja*, every evening a dose (four glob.) for two evenings.

If it does not yield to one of the above remedies, take *Silicea*, every third evening a dose (four glob.), until four doses are taken. If then not better, take *Sulphur*, in the same manner.

If this irritation shows itself around the anus or the pri-



vate parts, (*prurigo*) its cause is rather a constitutional one, and requires *Calcareae*, *Lycopodium*, *Sulphur*, *Nitric acid*, and *Sepia*. If it is around the anus, *Dulcamara*, *Nitric acid*, *Petrol*, *Sulphur*, *Lycopodium*, *Graphites*. If on the *scrotum*, *Thuja*. If on the *puddendum*, *Calcareae*, *Carbo*, *Conium*, *Sepia*, *Sulphur*, *Nitric acid*.

ADMINISTRATION.—In using these remedies, begin with the first—put twelve globules in four tablespoonfuls of water, and take, evening and morning, a teaspoonful. After it is taken, wait four days; then take the next in the same manner, if not better at that time. Beside this, wash frequently in cold water.

### *Chilblains.*

DIAGNOSIS.—Chilblains mostly appear on hands and feet which have been frost-bitten, and, if they burst and ulcerate, are extremely painful.

TREATMENT.—If the parts begin to swell, assuming a dark, reddish-brown color, with itching and beating, worse in the evening or at night, take *Pulsatilla*, in solution (twelve globules in half a teacupful of water), morning, noon, and evening a teaspoonful, and occasionally wash the parts affected in a teaspoonful of the solution.

In a few days afterward, take *Sulphur*, evening and morning, a dose (four glob.) If the parts have a bright red color, and itch more in the warmth, take *Nux vom.*, in the same manner.

If the parts are very painful and burning, take *Chamomile*, prepared in the same manner; but give, every three or four hours, a teaspoonful, alternately with *Arsenic*, or else *Hepar sulph.*, in the same manner.

If the swelling is hard and shining, take *Arnica* internally, and wash externally with two drops of the tincture to a teaspoonful of water, the same as *Pulsatilla*.



If the swelling still increases, and appears of a bluish red, take *Belladonna*, in the same manner as *Chamomile*.

DIET.—Abstain strictly from pork, and all irritating substances, such as pepper, and too much salt and salted meats.

N. B. If the disease will not improve under the above treatment, apply to a homœopathic physician.

### *Excoriations. (Intertrigo.)*

If they appear during the summer, in adults, give *Arnica*, *Carbo veg.*, *Nux vom.*, *Lycopodium*, *Sulphur*, beginning with *Arnica*, every evening a dose (four glob.) for two days, and then wait two days for its effect—taking the next in the same manner, and so on until better.

*Chafing of bedridden patients* requires *Arnica* (ten drops of the tincture to half a teacupful of water,) externally applied, with cloths dipped into the mixture. At the same time place a soft, tanned deerskin under the sheets of the bed, the hairy side down.

If the affected parts look bluish, and there is danger of mortification, give *China* internally, in alternation with *Carbo veg.*, every four or six hours a dose (four glob.), and wash externally, with a solution of *Silicea* (twelve globules in half a teacupful of water), several times a day.

*Disposition to fester.*—There exists in some persons a disposition of the skin to fester, if the slightest injury has taken place, or ulcers do not heal. In such a case, give of the following remedies every week, on two succeeding evenings, one dose (four glob.) until better, *Chamomile*, *Hepar sulph.*, *Lachesis*, *Silicea*, *Sulphur*.

*Chapped hands (rhagades)*, from working in water, requires *Calcarea*, *Hepar sulph.*, *Sepia*, *Sulphur*, administered in the same manner.

Cracked skin, from cold in the winter, requires *Petrol*, *Sulphur*, in the same manner.

RINGWORM. (*Herpes circinnatus*.)

This eruption appears in small rings on various parts of the body; within them the skin looks healthy as usual—at least, this is the case at first. They are more apt to appear in summer.

TREATMENT.—Dissolve *Sepia* (twelve glob.) in half a teacupful of water, and give for three days, morning and evening, a teaspoonful, and wash with part of the solution; repeat the same for eight or ten days, if necessary.

TETTER. (*Herpes*.)

This term comprehends a great many varieties of the same disease, which, when present, show a constitutional disorder that ought to be treated by a homœopathic physician. The tetter may appear on any part of the body—hands, face, lips, the ears, etc., and requires different remedies accordingly; but the easiest direction which can be given for practical purposes, in a domestic way, is to be guided by its discharging quality.

*Dry tetter* requires *Dulcamara*, *Sepia*, *Silicea*, *Sulphur*.

*Running tetter* — *Dulcamara*, *Rhus*, *Graphites*, *Calcarea*, *Lycopodium*, *Sulphur*.

*Bleeding tetter* — *Arsenic*, *Carbo veg.*, *Rhus*, *Mercury*, *Sulphur*.

*Ulcerating tetter* — *Mercury*, *Sulphur*, *Rhus*, *Sepia*, *Lycopodium*.

ADMINISTRATION.—In using these remedies for a tetter having one of the above qualities (its locality on the body does not make so much difference), take a dose of the first named remedy on two consecutive evenings, (six glob.) and repeat this every week for four weeks, then discontinue for

two weeks, taking the next remedy in the same manner, if the tetter shows no improvement during that time. In case of amelioration, however, take no more medicine unless worse again.

Keep the air off the tetter by a simple cover.

DIET as in chilblains.

### *Itch. (Scabies.)*

Not every eruption or pimple that itches is the itch—a malady so much dreaded in a family, that everything is welcome, and indiscriminately used, which will tend to eradicate this loathsome disease; and thus salves and ointments without number are used, but not without danger. Even if apparently cured by these salves, it is only driven *into* the system, laying the foundation for innumerable diseases afterward. Therefore, never do anything of this kind, under any consideration. First, be sure that it is *the real itch*, which is contagious. Frequently, it is only a disease described in the article on “Itching.”

DIAGNOSIS.—The real itch appears in pointed vesicles, filled with a transparent serous fluid, mostly about the wrists, between the fingers, and around the joints. The itching increases in the evening, especially in the warmth of the bed. It never appears on the face.

TREATMENT.—*Sulphur* is the specific for it. Take, every evening, a dose (ten glob.), for eight days; at the same time, wash the parts with half a pint of water, into which is put a grain of powdered *Sulphur*, twice a day, morning and evening, shaking the mixture well before using. Repeat this treatment if, after a week, no improvement has taken place; but, then, alternately with *Mercury*, in the same manner.

If, after another week, there is no improvement, and the eruption is yet small and dry, take *Carbo veg.*, for eight

days, every other evening a dose (six glob.), and, afterward, if necessary, *Hepar sulph.*, in the same manner; but, if, during this time, pustules have made their appearance, give *Caustic*, in the same manner.

The *pustular* or *humid* variety generally spreads more over the body, and also appears frequently on the back, shoulders, arms, and thighs. In this form, *Sulphur* and *Lycopodium* are necessary, given, as stated above in "*Sulphur and Mercury*," for eight days; then wait eight days, and, if no better, take *Caustic*, and, after it, *Mercury*, in the same manner.

*Sepia*, internally and externally, applied in the same manner as *Sulphur*, has frequently been beneficial when nothing else seemed to have the desired effect.

When the pustules are large, and turn yellow and blue, take *Lachesis*, for several evenings a dose (six glob.)

When the itch has been suppressed by external applications, and dangerous results threaten, give *Sulphur* and *Arsenic*, in alternation, every evening a dose (six glob.)

In obstinate cases, apply to a homœopathic physician.

#### *Milk-crust. (Crusta lactea.)*

This disease, as, in fact, all chronic eruptions of any extent, ought to be treated by a skillful homœopathic physician; yet we will here insert as much of the treatment as will warrant its rational commencement and often favorable termination.

DIAGNOSIS. — It consists of numerous small, whitish pustules, appearing in clusters upon a red basis, first on the face, the cheeks and forehead, but spreading, afterward, over the whole head and other parts of the body. The redness and swelling frequently increase, and, with it, the itching becomes intolerable, particularly if a large part of the face and head is covered with the eruption. In such

cases, the glands on the neck and under the ears enlarge in consequence.

Children, from four to eighteen months of age, are liable to it.

TREATMENT.—*Rhus* is the principal medicine, of which is to be given, every fourth evening a dose (four glob.), in alternation with *Hepar sulph.*, in the same manner, so that the child takes one or the other of the medicines every other evening.

If the child at any time is very restless, and the itching seems to be very aggravated, particularly at night, give *Aconite* and *Chamomile*, alternately, every two hours a dose (two glob.) until better.

If the running of the eruption suddenly dries up, and the child becomes drowsy, sleeps with eyes half open, has a hot head and cold feet, give *Belladonna* and *Hellebor.*, as directed under the head of "Measles." Externally, apply nothing but a little sweet cream, or wash occasionally with weak soapsuds.

*Scald Head.* (*Ring worm of the Scalp.*) (*Tinea capitis.*)

There are several varieties of this disease ; but for practical purposes, the distinction of dry and running scald head is sufficient.

DIAGNOSIS.—It appears, generally, on the hairy part of the head, in numerous yellowish pustules, finally forming a thick crust covering the head and neck of children from two to fourteen years of age ; bad as it is, and obstinate to cure, yet its sudden disappearance, after the application of violent external means, creates worse internal disorders, and frequently even death. It is highly contagious ; be careful even with the articles of clothing, particularly on the patient's head ; let no other child come in contact with it.

It generally lasts a long time, and ought to be treated by a skillful homœopathic physician.

**TREATMENT.**—First institute the treatment given under “Milk-crust,” and continue it for four weeks; if no amelioration takes place during that time, give, if the eruption is of the *dry kind*, *Calcarea carb.* every third evening a dose (six glob.) for two weeks, and then, if there is no sign of improvement, give *Sulphur* in the same manner. If it is the *running scald head*, give *Lycopodium*, and afterward *Sulphur*, in the same manner. If scrofulous symptoms appear, such as swelling of the glands on the neck and throat, give *Bryonia* and *Dulcamara*, alternately, every six hours a dose (four glob.) until better. If the discharge of the eruption is very corrosive, causing ulcers, give *Arsenic* and *Rhus* in alternation, every evening and morning a dose, (four glob.) and wash with a solution (twelve glob. in half a teacupful of water) of each medicine, applying it to the edges of the scab, while giving it internally.

**DIET AND REGIMEN.**—The usual homœopathic diet must be adhered to strictly. (See “Introduction.”)

The hair ought to be removed in the beginning of this disease.

### *Corns. (Clavi pedis.) Induration of the Skin.*

*Corns on the Feet.*—If they will not disappear after several cuttings and applications of *Arnica* tincture (six drops mixed with two tablespoonfuls of water), an internal treatment must be resorted to for their eradication.

First, take *Antimon. crud.* every evening a dose (six glob.), if the pains are more pressing, and as if *needles* were running through the corn. If the pain is of a burning nature, take *Calcarea carb.* every other evening a dose (four glob.) If they are inflamed, take *Lycopodium*, *Sepia*, *Silicea* in the same manner, taking every week another



remedy if necessary. If they are particularly troublesome in wet weather, take first *Bryonia* and *Rhus*, alternately, every two hours a dose (four glob.), for eight hours, and then twelve hours afterward a dose of *Sulphur*.

*Induration of the Skin.*—Sometimes the *skin indurates* (gets *callous*) on the *hands and feet*, becomes painful and peels off—in such cases take *Graphites* internally, every second evening a dose (four glob.) until four doses are taken, and use externally the same remedy (twelve glob. in half a teacupful of water well shaken) three times a day.

#### *Warts. (Verrucæ.)*

Neither cauterize nor cut these excrescences ; it is too dangerous ; a better remedy is the application of the tincture of *Rhus* on the wart three times a day. They will quickly disappear.

#### *Whitlows on the Fingers. (Felons.) (Panaris.)*

This disease exists in the form of an abscess, more or less deeply seated on the end of the finger, attended with severe pain and considerable swelling.

In the beginning, its formation may frequently be prevented by dipping the finger quickly into water almost boiling, or by wrapping around it finely powdered wood soot made wet with alcohol, or by washing the finger with a mixture of a drop of creosote in half a teacupful of water. But if this does not avail, keep warm bread and milk poultice around it all the time till it opens, and take internally *Mercury*, *Hepar sulph.*, and *Silicea*, alternately, every six hours a dose (four glob.), for two or three days.

After it opens, wash the finger three times a day in water, in half a teacupful of which twelve globules of *Silicea* are dissolved, or keep the bandage around the finger wet with this solution.



ULCERS.

We can only give their general treatment here, as their origin and appearance are too various to be discussed in a work on domestic practice. They arise, mostly, from a diseased condition, which must be first changed by systematic internal treatment, before the external sign of it, the ulcer, can heal. When it is forced to close itself by violent external means, the disease, of which it was the expression, attacks internal parts, and the danger is thus increased, as the dignity of the organ affected is greater.

Be careful, therefore, in using salves and ointments.

DIAGNOSIS AND TREATMENT. — First. An ulcer may be deep, presenting a *hollow excavation*. In this case fill it once or twice every day with dry scraped lint, previously cleaning it well with warm water and tying it up with a bandage. This promotes healthy granulation, and the hollow gradually fills up. Internally give *Lachesis*, *Hepar*, and *Silicea*, alternately, every second day a dose (four glob.), until four doses of each are given, or improvement is perceptible.

Second. An ulcer may be *flat, superficial*, sometimes presenting *proud flesh*; in this case dress it with finely powdered loaf sugar, or with a bandage dipped in cold water and changed from time to time. Internally, *Petroleum*, *Lycopodium*, and *Silicea*, in the same manner as above.

Third. An ulcer may be *fistulous*: in this case try to compress, if possible, the farthest ends of the fistula, or fill them with dry lint to excite healthy granulation, always trying to heal up the deepest cavity first. Give *Antimon. crud.*, *Calcarea carb.*, *Silicea* and *Sulphur*, in the same manner as above.

Fourth. An ulcer may be *callous*, with thick, hard, broken margins. In this case fill the bottom with dry lint,

and dress around the margins with simple cerate. Give *Arsenic*, *Pulsatilla*, *Lycopodium*, and *Sulphur*, in the same manner as above.

Fifth. An ulcer may be *carious*, proceeding from the bone; dress outside with simple cerate and give *Mercury*, *Sulphur*, *Calcareo carb.*, and *Lycopodium* in the same manner as above.

Sixth. An ulcer may be *varicose*, caused by swelling of the veins, for instance, on the lower limbs; dress outside with simple cerate and lint, or cold water bandage, and give *Pulsatilla*, *Lycopodium*, *Lachesis*, and *Sulphur* in the same manner as above.

We have another condition of the ulcer to take into consideration, in order to determine what medicine has to be given.

First. Ulcers may be *painful*; in this case, give

a. When accompanied by a burning or drawing sensation, *Arsenic*, *Rhus*, and *Sulphur*, in the same manner as above.

b. When *beating*, *eating*, or *pressing*, *Mercury*, *Lachesis*, *Lycopodium* and *Sulphur* in the same manner.

c. When *itching*, *Hepar sulph.*, *Pulsatilla*, and *Sulphur* in the same manner.

Second. An ulcer may be *without pain*; in this case give *Phosphor. acid*, *Carbo veg.*, *Sepia*, and *Sulphur* in the same manner.

Third. An ulcer *smells offensively*; in this case give *Carbo veg.*, *Arsenic*, *Pulsatilla* and *Sulph.* in the same manner.

Fourth. It *spreads* very much, *increasing in size*; in this case give *Mercury*, *Lachesis*, *Hepar sulph.*, *Silicea* and *Sulphur* in the same manner.

*Syphilitic Ulcers* require *Mercury*, *Nitric acid*, *Lachesis* and *Thuja* in the same manner.

*Abscesses. Tumors. Swelling of Glands.*

When a congestive or inflammatory swelling is not dispersed or absorbed, it changes gradually into an abscess; matter forms which finally discharges through an opening of the sore, either prepared naturally, or artificially by a lancet.

TREATMENT.—Before an abscess opens, it may be very painful; in this case, poultice it with bread-and-milk, and take internally *Belladonna*, *Lachesis*, and *Mercury*, alternately, every three or four hours a dose (four glob.), until better.

If it is a long time *maturing*, without being very painful, (*cold swelling*), take *Hepar sulph.*, *Iodine*, and *Sulphur*, every other evening a dose, alternately, and poultice it when it comes near breaking.

After an abscess has opened, wash it frequently during the day with a solution of twelve globules of *Silicea* in half a teacupful of water, and take internally *Silicea* every other evening a dose (four glob.) until three doses are taken.

If it be desired to open an abscess with a lancet, select the lowest part, if it is also the softest, which latter condition is *always* to be chosen first. In other swellings, particularly when they appear hard, with stitches through them, or soft, without much pain, apply nothing externally, until after having consulted a homœopathic physician. Meanwhile, take internally *Calcarea carb.* every three or four days a dose in the evening (four glob.), until six doses are taken; afterward, *Sulphur* in the same manner.

*Enlarged* and indurated *glands* on the neck, etc., require *Mercury* and *Dulcamara* every evening, alternately, a dose (four glob.); when suppurating, *Hepar sulph.* and *Silicea*,

every third evening a dose (four glob.), alternately until four doses of each are taken.

*Boils. Malignant Boils. Carbuncles.*

These are painful, hard tumors, of a pyramidal form, and deep red color, generated sometimes by a constitutional tendency, very often, however, as critical discharges after acute or eruptive fevers, or terminating chronic eruptions, such as itch, etc. They suppurate slowly, and discharge, on breaking, a little pus mixed with blood, and exhibit a core which is gradually discharged.

TREATMENT. — The best application externally is softened bread; all other things, such as roasted onions, soap, and sugar, are too irritating, and enlarge the suppurative sphere. Internally take first *Arnica* evening and morning a dose, (six glob.), and if the boil gets very painful and red, take *Belladonna* and *Mercury*, alternately, every three or four hours a dose (four glob.) If matter has formed, take *Hepar sulph.* every four hours a dose (four glob.) until the tumor breaks.

The predisposition to boils may be removed by taking *Sulphur* every week one dose (six glob.) for six weeks. If a boil becomes *blue*, and *increases* very fast (*malignant boil*), take *Lachesis* every three or four hours a dose (four glob.) in alternation with *Arsenic*. Let the treatment be the same if typhoid fever ensues, and the patient is very weak, restless, and sleepless (*Carbuncle, Anthrax*), and mortification threatens.

*Scirrhus. Cancer.*

We mention these diseases here, not to give their treatment, because the seriousness of their character would not allow of their being treated domestically; but it is necessary to warn patients of this kind against all those who

pretend to cure them with the knife or cauterizing applications. They always end fatally under such treatment, and with increased misery. At the same time, we would advise them not to despair, but to apply immediately to a skillful homœopathic physician, who possesses the only means which can rationally afford relief, and which sometimes effect a complete cure.

*Indurations* under the skin, in glands, on the lips, nose, etc., with stitching pains through them, should receive attention, from the fear that they might be of the *scirrhous* nature ; if so, and if not attended to in time, they will become open cancers.

In such cases, relief is possible, but only by the timely advice of a skillful physician. Meantime, take *Belladonna*, every third evening a dose (four glob.), until four doses are taken.

In open cancers, the distressing burning pains are relieved by *Arsenic*, in solution (twelve glob. in half a teacupful of water), every two or three hours a teaspoonful, until four or six are taken.

## CHAPTER IV.

### FEVERS.

THE term *fever* is frequently misunderstood — people intending to express by it the disease itself, while, in reality, *fever is the reaction of the vital powers against a disease*; and, as every action in nature can be of a threefold kind, either not powerful enough for its purpose, or just powerful enough, or, lastly, too powerful, the fever, also, as an action of nature, allows these three divisions.

Accordingly, we have simple *irritative fevers*, where the effort of nature to remove the disease is just sufficient; or, *inflammatory fevers*, where this effort is greater than is necessary; or, lastly, *torpid fevers*, where the effort is not sufficient to accomplish the removal of the disease.

In all forms of acute disease, fever of one of the above kinds is present to a greater or less extent. It is obvious that one form of fever may, under circumstances favorable to the change, merge into another; for instance, a simple irritative fever may become an inflammatory one, if the patient is overstimulated; or a torpid one, if his vital energy is too much depressed.

How dangerous, therefore, must be the use of allopathic remedies in fevers, as, when a little too strong or too weak, they lead to such awful consequences! But, in Homœopathy, this result is altogether obviated, as the remedial agent used neutralizes, by its specific action, the disease of which the fever was only an attendant symptom.

Thus we see, in pneumonia, under allopathic treatment, that the fever frequently returns with great violence, although the patient, shortly before, had been depleted to exhaustion. The reason is, that the cause of the fever, the pneumonia, had not been removed by the depletion; the reactive force or vital energy to overcome the disease was merely diminished, and, therefore, the system rouses itself up again to a second attempt; if checked or depressed again, it must finally sink to rise no more; it must fall into the *torpid* or *typhoid* form. Not so, under the homœopathic treatment, where the disease itself is specifically reached and as it were absorbed; for when the disease is removed, no effort of nature is necessary, and, consequently, the fever disappears of itself.\*

The action of the unaided vital force of nature, during the fever, is in two directions—by the *nervous*, and by the *vascular* system—to bring about a crisis, sufficient for the extermination of the disease; but if it fails in establishing a sufficient crisis, nature has to surrender, or death ensues; and this, again, either by paralysis of the nervous system (paralysis properly so called), or paralysis of the vascular system (mortification.)

The less the disease, therefore, the less needs be the crisis; and we aid nature, indeed, if we take away the disease, but not if we take away her vital power to overcome it. Herein lies another important difference between

---

\* The best illustration of the *modus operandi* of the homœopathic medicines, or what is meant by specific action generally, is found in the effect of the dynamic agents of nature on each other. Positive and negative electricity, being similar, but not identical in their nature, neutralize or absorb each other; while positive and positive, being identical, repel each other, having no affinity to each other. In the same manner, two similar diseases absorb each other to the zero point; this state is where neither exists, and therefore health.



the two medical systems ; and it is easy to see how salutary the homœopathic, and how destructive the allopathic treatment of fevers must be.

In most cases, Homœopathy takes away the disease so entirely, by its specific method, that no crisis at all appears. If it does take place, however, it manifests itself by diarrhea, perspiration, or other secretions, or by an eruption ; after which, the skin becomes moist and the pulse regular.

DIAGNOSIS.—The symptoms by which the presence of the three forms of fever may be known, are given in the following characteristics :

First. *Irritative form*.—Shivering, preceding a moderate heat ; skin natural or a little moist ; pulse not very much accelerated, but fuller, stronger, and not hard ; urine a little more reddish than common, and with a sediment after the fever leaves.

Second. *Inflammatory form*.—Great lassitude precedes a short but severe chill (which frequently does not appear, however), followed by an intense heat ; skin dry and burning ; eyes sparkling ; tongue dry ; thirst intense ; pulse quicker, hard, and full ; urine red and scanty.

Third. *Torpid form*.—The greatest lassitude prevails during its presence ; the patient feels very weak, although he may be very hot at times ; at others, the temperature is very much diminished ; the skin is now very dry and inclined to crack ; then, again, covered with clammy perspiration.

At one time, the tongue is dry and black ; at another, either natural or covered with a tough mucous. There is often an absence of thirst, although the tongue is dry ; and then, again, thirst is intense with a moist tongue.

The pulse shows the same anomaly, being sometimes full, with a low temperature of the body ; and again, at other times, small and weak, with a high fever-heat of the system.

TREATMENT.—As we have only given here the general character of the three kinds of fever that combine themselves with local diseases of the different organs, we must, also, generalize their treatment here, and refer, for details, to the affections of the respective organs where fever manifests itself. For instance : fever of the lungs, *pneumonia*, inflammatory, or torpid (*typhoid*.) See “Affections of the Lungs.”

The essentials in the treatment of all fevers are as follows :

Quiet and rest of body and mind, as much as possible ; cool and even temperature of the patient's room, say from fifty to sixty degrees, according to comfort ; pure air and thorough ventilation, without producing it by drafts passing on or over the patient. Let the patient lie on mattresses, and be covered with quilts, light, but sufficiently warm.

As the patient himself does not desire solid food, we ought not to give it ; but, when thirsty, we must never refuse drink. The best drink is cold water, as much as the patient desires, if it agrees with him ; frequently give toast water, barley or rice water, or lemonade, except when the patient has a diarrhea, or is taking *Aconite*, with which no acid will agree.

In convalescence, baked apples and stewed prunes are very salutary, if there is no diarrhea at the time ; the latter, however, should be partaken of sparingly.

We will now treat of several other general forms of fever as they occur.

#### COMMON OR EPHEMERAL FEVER.

(*Febris simplex*.)

DIAGNOSIS.—Shivering, followed by heat, restlessness, thirst, quick pulse ; its termination is generally within twenty-four or thirty-six hours, by profuse perspiration.

It is mostly caused by exposure to sudden changes of temperature, and will then terminate in from twenty-four to forty-eight hours ; but where it is the forerunner of other more serious diseases, such as scarlet fever, measles, etc., it will be superseded by them.

TREATMENT. — *Aconite*, in solution (twelve glob. to half a teacupful of water), every two hours a teaspoonful, until perspiration ensues.

DIET AND REGIMEN. — Cover the patient well, and let him drink cold water, which promotes perspiration. Bathing the feet in warm water before lying down is allowed, but it should be done near the bed.

N. B. If pains are felt in different parts of the body, see respective headings, and treat accordingly.

#### GENERAL INFLAMMATORY FEVER. (*Synocha*.)

DIAGNOSIS. — See “Fevers—Inflammatory form.”

CAUSES. — These symptoms are frequently caused by sudden checks of perspiration and exposure to high degrees of temperature or its sudden change. They are also caused by external injuries or lesions. Plethoric and young persons are most liable to it.

It frequently precedes typhus fever, particularly under allopathic treatment, or, if neglected, seizes upon some organ, where it then runs through its phases. In such cases, see the respective headings.

TREATMENT. — *Aconite* must be given first, the same as in common fever. See preceding article.

But, if, after six hours, there is no improvement, give *Belladonna*.

When the *head is hot*, with *violent pains in the front part*; *face red*; *sensitiveness to light*; *eyes sparkling*; *noise distresses*; very restless or drowsy; great thirst; delirium;

ADMINISTRATION — the same as *Aconite* above, or alterna-

ting with it, until better. But, if, with the general symptoms of inflammatory fever, there is *swimming in the head* on rising or moving; some delirium; *oppression at the pit of the stomach*; *vomiting after drinking*; *constipation*; aching pains in the limbs and small of the back; oppression of the lungs; give *Bryonia*; and, if the weather is damp and rainy at the time, give *Rhus*, in alternation with *Bryonia*, in the same manner as above in *Aconite* and *Bellad.*, until better.

DIET AND REGIMEN—the same as under the head of “Common Fever.”

#### NERVOUS OR TYPHOID FEVER. (*Typhus.*)

These names, signifying the same disease, are only its various grades; but these are distinguished by prefixing the appropriate word in an adjective form, as a *malignant typhoid fever*; a *malignant typhus*; or, a *malignant nervous fever*; all three of which signify the same.

Although we could not advise any one to treat a fever like this on his own responsibility, with no other guide than a work on domestic practice, unless in cases of absolute necessity, it is, nevertheless, advisable to give a sufficient account of the disease and its remedies, to enable the inquirer to distinguish it from others, and commence its treatment with the right remedies.

The progress of the disease may frequently be arrested by early and proper treatment in its commencement. To the realization of this important point our particular attention will be directed, as in the prevention of disease is seen the great advantage of domestic practice.

DIAGNOSIS.—The real nature of typhus consists in an alteration of the blood, which becomes thick, decomposed, and carbonized; but as such a qualitative change of the blood can not take place at once, the disease begins slowly,

but is sure in its progress, unless arrested by the proper homœopathic remedy.

The first indications are, general lassitude, chilliness, followed by occasional feverishness, and, sometimes, pain in the head, chest, or abdomen, followed by drowsiness. Then appear the symptoms given under the heading, "Fevers—Torpid form." Beside these, one characteristic symptom of real typhus is, a pain in the abdomen near the right hip, sensible to heavy pressure; here, in the intestines, ulcerations of the glands invariably take place in real typhus; hence the pain only on pressing heavily. Under such symptoms, it runs its course, if not checked by good homœopathic treatment, perhaps to convalescence, by frequent critical efforts, in about twenty-eight days.

The best crisis is perspiration, coming on gradually; or sleep, in the place of delirium and restlessness. Bleeding of the nose is less favorable; and a still less favorable crisis is the swelling of the glands in front of, and below the ear; or, it ends in death, either by paralysis or mortification. (See "Fevers.")

CAUSES.—In general, any depressing influence on the vital powers of the body may produce typhus, but, particularly, overexertion of the body or mind; excesses of any kind; long and imprudent exposure to cold or damp weather; bad air, water, and mode of living; crowded population in filthy streets and small apartments; exhausting blood-letting in inflammatory diseases; or, lastly, a real typhus contagion, which is generated in one form of the disease. (See below, "Contagious Typhus.")

PREVENTIVES.—The best preventive is, to avoid the above described causes of the disease, and to take, if already threatened with it, *Bryonia* and *Rhus*, every evening alternately a dose (four glob.), for eight days, or until better. Keep quiet and in the house; follow the diet prescribed in

fevers generally ; keep the feet warm and head cool ; drink freely of cold water, and take no allopathic remedies, such as pills, etc., *under any consideration*.

### DIFFERENT FORMS OF TYPHUS.

First. According to the *time of its appearance*, typhus is —

- a. *Primary, idiopathic, or true and contagious typhus ; or,*
- b. *Secondary, following another disease, and pseudo-typhus ; that is, fevers with apparently typhoid symptoms.*

Second. According to its *locality*, typhus is —

- a. Typhus of the *brain* (*typhus cereбрalis*), *typhoid brain-fever, and congestive fever ;*
- b. Typhus of the *lungs* (*typhus pulmonalis*), *typhoid lung-fever ;*
- c. Typhus of the *bowels* (*typhus abdominalis*), *abdominal typhoid, and yellow fever.*

N. B. In either of these three forms, those characteristic symptoms of real typhus (pains, on pressure near the right hip, in the intestines), as mentioned above (see “Diagnosis”), never fail to appear, even if the brain or lungs are primarily affected.

Third. According to its *qualitative appearance*, typhus is —

- a. *Versatile* (*typhus versatilis*), with *excited* nervous action ;
- b. *Stupid* (*typhus stupidus*), with *depressed* nervous action ;
- c. *Putrid* (*typhus putridus*), with tendency to *organic dissolution*.

For practical purposes, we follow the distinctions made under the “third” or last division, as it is the only *qualitative* differences in typhus that make different remedies necessary. The stupid form is oftener found in the typhus of the brain ; the versatile, in typhus of the lungs ; and the putrid, in typhus of the bowels. Yet there are exceptions.



TREATMENT. — At the commencement of all three forms, if there is headache, giddiness, chilliness with alternate heat, rheumatic pains in the limbs and arms, sometimes a slight cough with pains in the back, with or without diarrhea, sickness at the stomach or vomiting, restlessness at night, furred tongue; give *Bryonia* and *Rhus*, every two hours a dose (four glob.), until three of each are taken. Make the patient lie down, well covered, and await the reaction for twenty-four hours; giving nothing, during that time, except one or two doses of *Coffea* <sup>cc.</sup> and *Bellad.* <sup>cc.</sup>, if he is very restless during the night. As soon as the patient perspires and the symptoms gradually disappear, the typhus, as such, is checked, and its return is prevented by giving, on the third day, evening and morning, a dose (four glob.) of *Sulphur*; during which time the patient still keeps his room, and diets as recommended in fever.

If, however, the disease progresses, and the real typhus symptoms appear more and more, give as follows:

If inflammatory symptoms appear, such as full, hard pulse; hot, dry skin; violent thirst; give *Aconite*, two or three doses, every two hours one (four glob.); and alternate with *Belladonna*, in the same manner, if to the above symptoms are added violent congestion of the head or lungs, with severe headache; sensitiveness to light and noise; starting after closing the eyes, and wild expression of countenance. Wait eight or twelve hours for the effect.

If no change for the better takes place, but, on the contrary, the symptoms increase in violence, give *Bryonia* and *Rhus* again, alternately; but now only every four hours a teaspoonful of their respective solutions, (twelve globules in half a teacupful of water), with which continue till four of each are taken, or until better.

After this time, the disease may have changed, so as to make one of the following remedies necessary.



*Phosphor. acid.* Great exhaustion, flighty when awake, always lying on the back in a drowsy state; giving either no reply to a question, or replying incoherently, loquacious delirium, or low muttering, picking the bedclothes, black lips, dry, hot skin; frequent, copious, watery, and involuntary diarrhea, sometimes bloody; give alternately with *Rhus* in the same manner as *Bryonia* and *Rhus* were above directed to be given.

*Arsenic.* If, with these symptoms, there is *extreme prostration of strength*, falling of the lower jaw, open mouth, dull and glassy eyes, burning thirst, profuse diarrhea, pulse barely perceptible; alternate *Carbo veg.* with *Arsenic*, if to the above symptoms is added rattling respiration, cold perspiration on the face and extremities, and very offensive evacuations, to be given in solution as above, every half hour a teaspoonful. These remedies suit principally in the *putrid* form of the disease.

In the *stupid* form, however, after *Bryonia* and *Rhus* have been given, the following remedies may be found necessary:

*Opium.* When there is great drowsiness, with snoring breathing, open mouth, low mutterings, picking the bedclothes, lethargy, all discharges passing involuntarily, give every hour a dose (four glob.), until better, if not give *Phosphor. acid* as above.

*Lachesis.* Under the same symptoms as those under *Opium*, particularly if the tongue is *very red* and shining, as if varnished; in the same manner as above.

*Lycopodium.* Under similar symptoms as *Lachesis*, and then in alternation with it, in solution, every two hours a teaspoonful, until a change takes place; but particularly if there is a strongly marked *redness of the cheeks*, constipation, and screaming, especially on awakening, *dryness of the tongue without thirst*.

In the *versatile* form, after *Bryonia* and *Rhus* have been given, the following medicines may be necessary, as,

*Hyoscyamus*. When there is a desire to escape; twitchings, and grasping at persons that are near; hot, red face, with bluish cheeks, sparkling eyes, dry, brownish tongue; alternately bland and furious delirium, involuntary evacuations; give every three hours a dose (four glob.), alternating with *Belladonna* in the same manner; or, in alternation with

*Stramonium*. If to these symptoms are added frightful fancies, loquacious delirium, staring eyes, spasmodic action, convulsions, and aversion to liquids; no stool or urine, and loss of consciousness.

There are several other remedies, which, under various circumstances, are important in the treatment of this disease. *Arnica*, *Arsenic*, *Camphor*, *China*, *Veratrum*, *Cantharides*, *Cocculus*; see in "Materia Medica" their pathogenetic symptoms, and give them if similar with the state of the patient.

N. B. If, at any time, the patient is very sleepless, give *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, every hour a dose, (four glob.), alternately. If there is clammy *cold* skin, debilitating and clammy sweats, diarrhea or disposition to it, give *Camphor* and alternate with *Coffea* if very restless, every half hour a dose until better. During convalescence, *China*, *Mercury* and *Sulphur* are frequently necessary. (See "Materia Medica" at end of book.)

DIET AND REGIMEN the same as under "Fevers."

*Congestive fever* is a typhoid brain fever, and is treated as already stated, with *Aconite*, *Belladonna*, *Bryonia*, *Rhus*, *Opium*, etc.

The *contagious typhus* is in nothing different from the above, only it is caused by contagion. The attack occurs but once in a person's lifetime, and its course is marked with greater severity.

The *Pseudo-typhus* or *bilious-rheumatic* fever is composed of the incipient symptoms of typhus, and has to be treated accordingly, as stated above, with *Bryonia* and *Rhus*, followed by *Sulphur*, to prevent its running into real typhus. For the treatment of this form, however, compare *Ipecac*, *Pulsatilla*, *Ignatia*, and *Nux vom.* in the "Materia Medica" at the end of the book.

The *ship fever* is a species of the *contagious* and *abdominal* typhus; treat accordingly; compare as in "Yellow Fever."

#### YELLOW FEVER.

As I never had an opportunity to observe and treat this malignant fever, I thought it advisable to insert here the full article of Doctor Marcy on that subject, who had an opportunity to peruse the notes left by Doctor Taft, whose success in the treatment of yellow fever was astonishing:

This fever is exceedingly uncertain in its course, violence, and duration. It may strike its victim suddenly prostrate, overwhelming, in its severity, the whole system, and thus preventing a single rally of the circulatory vessels; or it may advance mildly, differing but little from an ordinary attack of remitting fever. In some instances it bears a strong resemblance to the higher grades of bilious fever. Much depends upon the peculiar circumstances of the individual attacked. If he is recently from a temperate climate, and unaccustomed to hot regions, he will be more susceptible to the action of the poison, than if he had been previously acclimated.

Medical men have supposed that after a certain period of exposure, the system becomes so completely accustomed to the miasm, that it loses all susceptibility to its influence, and in this manner the process of acclimation is accomplished. There is, doubtless, some truth in this idea, but

there are other causes which exercise quite as important an influence in this process. Those persons who abandon a temperate for a residence in a tropical climate, do so in that physical condition which the requirements, habits, and regimen of the former naturally generate. In a previous chapter we have seen that, in cold regions, where the atmosphere is highly condensed, a large amount of animal food is requisite to supply the system with sufficient carbon and hydrogen to resist and neutralize the action of the inspired oxygen. With these habits, appropriate only where a condensed atmosphere is respired, individuals seek the tropics, with bodies abounding in carbon, and continuing, in most instances, their accustomed regimen of animal food and stimulants, thus burdening their systems with an amount of the elements of nutrition far greater than the oxygen contained in the rarefied air which they inhale can decompose.

It is probable, therefore, that one of the chief predisposing causes of yellow fever, is the presence of a greater amount of carbon in the system than the inspired air can properly act upon. The exact equilibrium between the supply of the elements of the food and the absorbed oxygen is disturbed; the carbon predominates, and all of those derangements which proceed from a superabundance of this agent, necessarily ensue.

The inhabitants of tropical latitudes have comparatively but little desire for animal food, but prefer farinaceous diet, vegetables and fruits, in this manner naturally securing to themselves a due proportion between the elements assimilated and the oxygen absorbed; while the inhabitants of the north find it necessary to consume large quantities of meat and other articles abounding in the elements of nutrition, in order to preserve a healthy equilibrium. We, therefore, most strongly urge it upon those who remove

from cold to hot climates, to adapt their systems by appropriate regimen and strict temperance in all things, for the change, and we confidently predict that they will enjoy as great an immunity from this dreadful scourge as the natives themselves.

DIAGNOSIS. — The premonitory symptoms of yellow fever are giddiness, wandering pains in the back and limbs, slight chills, nausea, and frequent sensations of faintness.

After these symptoms have continued a few hours, a decided reaction occurs ; the circulation becomes excited, the face flushed, the eyes red ; there are violent pains in the head, back, loins, and extremities ; distress of stomach, and vomiting of acid bilious matters ; the surface becomes dry and burning hot ; mouth and throat dry, with intense thirst, and sometimes delirium.

The duration of this paroxysm is usually about twenty-four hours, although occasionally it continues two or three days, after which there is a remission of all the symptoms, except a distressed sensation in the stomach, with nausea and vomiting. The patient remains in this state with a considerable degree of comfort for a few hours, when there is a recurrence of many of the former symptoms in an aggravated form. The stomach now becomes extremely painful and sensitive ; vomiting is violent and incessant ; the fluids ejected are of a darker color ; the skin and eyes acquire a yellow tinge, and the mind becomes confused and wandering.

The duration of this second stage varies from twelve to forty-eight hours, with sometimes slight remissions toward the termination of the paroxysm, when the third or last stage sets in. This stage is characterized by the complete development of the dreaded "*black vomit*." At this period, the powers of the system all sink rapidly ; the pulse flags, and perhaps intermits ; the tongue becomes dry, black and

shrivelled; the breathing irregular and laborious; cramps seize the calves of the legs and the bowels; the whole countenance loses its natural, life-like expression; the extremities become cold; colliquative sweats, diarrhea, hemorrhages, and loss of intellect occur, and, finally, dissolution ends the scene.

This is only a brief outline of the more ordinary symptoms and course of the malady, and will, we trust, serve to aid the inexperienced practitioner in his diagnosis. Each case, however, must necessarily present modifications according to the predisposition, habits, and peculiar circumstances of the individual attacked.

CAUSES.—When animal and vegetable matters are submitted, for a considerable length of time, to the daily influence of intense solar heat, and a certain amount of moisture in the crowded and filthy streets of cities, or other confined places, a miasm is generated, which, under favorable circumstances, will cause yellow fever. Concerning the nature of this miasm we know nothing; but it is evident that the continued high degree of temperature to which these substances are exposed, and the confinement of their noxious emanations within the walls of crowded cities, develop a more virulent morbid agent than is the case when the same matters are exposed in the open country, or to a more irregular and less intense heat, such as usually occurs in more temperate localities.

There are several other causes which act as powerful predisposing influences, one of the most important of which, as before mentioned, is the too free use of animal food and stimulants. We may also include in this category, irregular habits, mental anxiety, depression of spirits, fear, grief, exposure to night air or to a burning sun, and, indeed, whatever else tends to debilitate the organism.

TREATMENT.—The remedies most commonly applicable in



the treatment of this affection are, *Ipecac*, *Belladonna*, *Bryonia*, *Rhus*, *Arsenic*, and *Aconite*. The other medicines likely to prove serviceable are, *Nux vom.*, *Mercurius*, *Veratrum*, *China*, *Sulphur*, *Cantharides*, *Carbo veg.*, and *Lachesis*. The late and much lamented Doctor Taft, of New Orleans, was eminently successful in his treatment of the yellow fever as it occurred in that city. Sometime since, we had the pleasure of perusing a letter from a highly intelligent gentleman of New Orleans, in which he states, that the success of Doctor Taft was so great in this malady as to attract the marked attention of a large number of citizens; and the writer expresses a deliberate opinion, that a new and favorable era would soon have occurred in the management of this formidable affection, if the able and accomplished Taft had survived. The remedies which this physician found most successful, and upon which he chiefly relied, were *Aconite*, *Ipecac.*, *Belladonna* and *Bryonia*, in the *first*, and sometimes *second* stages; in the second and third stages, in addition to the above, *Rhus tox.*, *Arsenic*, *Veratrum*, *Cantharides*, *Carbo veg.*, *Nux vom.* These medicines were usually employed at the first attenuation, and frequently repeated, either singly or in alternation, as the circumstances of each case appeared to require.

When the first symptoms declare themselves, as dizziness, slight chills, pains in the back and limbs, uneasy sensations at the epigastrium, with nausea, vomiting, and sensation of faintness, *Ipecac.*, at the third attenuation, should be immediately exhibited. This remedy may also be found serviceable during the second and third stages, in alternation with some other article. Should the malady continue to progress, the following medicines should be considered, and, in proper cases, promptly administered.

*Belladonna.* Glowing redness and bloated appearance of the face; eyes red and sparkling, or fixed, glistening, and



prominent; tongue loaded with whitish mucus, or yellowish, or brownish; pulse variable. Dry burning heat; sharp, darting and shooting pains in the head; violent throbbings in the head; burning thirst; painful heaviness and cramp-like pains in the back, loins, and legs; pressure, cramp-like, and contractive pains in the stomach; inclination to vomit, or violent vomitings. During the remission, melancholy; dejection; when reaction comes on, great agitation, with continual tossing and anguish.

ADMINISTRATION. — *Belladonna* is for the most part applicable to the first stage of yellow fever. Twelve globules in half a teacupful of water, every two hours a teaspoonful, until better or another remedy is needed.

*Bryonia*. Skin yellow; eyes red, or dull and glassy, or sparkling and filled with tears; tongue dry, and loaded with a white or yellow coating; pulse rapid, and full, or weak and rapid. Severe pain and burning sensation in the stomach; vomiting, particularly after drinking; burning thirst; pains in the back and limbs; headache, aggravated by movement; eyes painful on motion; sense of fulness and oppression in the stomach and intestines. Anxiety, with dread and apprehension respecting the future; loss of memory; delirium.

Administration same as *Belladonna*, or in alternation with *Rhus*.

*Rhus*. Surface of a dirty yellow color; eyes glazed and sunken; tongue dry and black; lips dry and brownish; pulse quick and small; loquacious delirium, or coma with stertorous breathing; constant moaning. Distressing pain and burning in the stomach; nausea and vomiting; paralysis of the lower extremities; spasms in the abdomen; want of power over the abdominal muscles; colic; diarrhea; difficulty in deglutition, and pain on swallowing. Intellect dull and clouded; constant uneasiness; delirium.

Administration same as *Belladonna*. (See *Bryonia*.)

*Arsenic*. Face of a yellowish or bluish color; eyes dull and sunken, with a dark mark under them; sclerotica yellow; nose pointed; coldness of the body, with cold and clammy sweat; lips and tongue brown or black; colliquative sweats; pulse irregular, or quick, weak, small, and frequent, or suppressed and trembling. Sense of extreme debility; dull, throbbing, stunning, or shooting pains in the head; burning or sharp and darting in the epigastrium, or in the region of the liver; limbs feel stiff and useless; frequent evacuations, with tenesmus, or painless and involuntary; oppression at the chest, with rapid and anxious respiration; cramps in the calves of the legs; great oppression at the stomach, with violent vomiting, especially after drinking; drawing and cramp-like pains in the abdomen; sensation as if a weight was pressing upon the abdomen. Indifference; weakness of memory; stupidity; delirium, with great flow of ideas; loss of consciousness and of sense; raving.

ADMINISTRATION. — Dissolve in water, as stated in *Belladonna*. In urgent cases, a teaspoonful may be exhibited every half hour, until some change is produced in the symptoms. In less dangerous instances, the intervals of administration may be lengthened as circumstances require.

*Aconite*. Suitable in the first and second stages, when there are burning and dry skin; red cheeks; full and rapid pulse; red and sensitive eyes; tongue natural or covered with a whitish slimy coat; lips and mouth dry; vomiting of mucus and bile; urine dark red. Violent febrile reaction; sensation of intense heat; great thirst; acute pains in the temples, forehead, or on the side of the head; vertigo on rising, eyes weak and sensitive to light; pains and soreness in the back and limbs; nausea; general sense of debility; great heat and irritability of the stomach; short

and anxious respiration. When the fever is on, great anguish, anxiety, and restlessness; for the most part nightly delirium.

Administration. Dissolve in water as above. *Aconite* and *Belladonna* may sometimes be alternated with benefit in the first period of the disease.

In a majority of cases, a few doses of this remedy will be found indispensable, during the first reaction.

*Nux vomica*. Skin yellow; face pale or yellowish, especially round the nose and mouth; lower part of the sclerotica yellow; eyes inflamed with redness of the conjunctiva; eyes surrounded with a dark circle and full of tears; tongue with a thick white or yellow fur, or dry, cracked, and brown, with red edges; pulse variable. Burning pains in the stomach; pressure or cramp-like pains in the epigastrium; vomiting of acid, bilious, or mucous matters; frequent and violent hiccough; eyes sensitive to light; vertigo, or pains in the head; tremors of the limbs, cramps in different parts; thirst for beer, brandy, or some stimulant; contraction of the abdominal muscles; loose discharges of slimy or bilious matters or blood; burning pains at the neck of the bladder, with difficulty in urinating; coldness, paralysis, and cramps in the legs; feet benumbed and cramped. Excessive anxiety, uneasiness, fear of death; despair, or loss of consciousness and delirium, with moaning or muttering.

Administration same as *Belladonna*.

*Mercurius*. Yellow color of the skin; eyes red, blood vessels of sclerotica injected; eyes sensitive to light; paralysis of one or more limbs; tongue with moist, thick, white fur, or dry and brown mucus; fæces variable; pulse irregular, or quick, strong, and intermittent, or weak and trembling. Excessive inclination to sleep, or restlessness from nervous irritation; sense of fatigue and debility; ra-

pid loss of strength ; dizziness, or violent pain in the head ; violent convulsive vomiting of mucus and bilious matters ; burning pain and tenderness of the stomach ; constipation, or diarrhea with discharges of mucus, bile, or blood ; coldness of the arms and legs, with cramps ; excitability and sensibility of all the organs. Anguish and agitation ; weakness of memory ; apprehensions ; discouragement ; moroseness ; raving.

Administration same as *Belladonna*.

*Veratrum alb.* Face of a yellowish or bluish color, cold and covered with cold perspiration ; eyes dull, clouded, yellowish and watery ; lips and tongue dry, brown, and cracked ; hiccough ; coldness of the hands and feet ; trembling and cramps of the feet, hands, and legs ; evacuations loose, blackish or yellowish ; pulse slow and almost extinct, or small, quick, and intermittent. General prostration of strength ; confusion of head, or vertigo ; deafness ; difficult deglutition ; intense thirst ; violent vomiting of green bile and mucus, or black bile and blood ; burning in the stomach ; great exhaustion ; cramps in the stomach, abdomen, and limbs ; diarrhea. Timid, despondent, restless, loss of sense ; coma or violent delirium.

Administration same as in *Arsenic*, and with it in alternation.

*Sulphur.* Face pale, or yellowish ; eyes red, or yellowish ; aphthæ in the mouth ; tongue dry, rough, and reddish, or with white or brownish coat ; pulse hard, quick, and full ; fæces whitish, greenish, or brownish, bloody or purulent. Dizziness, or sharp pains in the head ; itching and burning pains in the eyes ; roaring in the ears ; nausea, with trembling and weakness ; vomiting of bilious, acid, bloody, or blackish matters ; pressure and pain in the stomach ; pains in the back and loins. Melancholy, sad, timid, undecided, wandering.

Administration. Dissolve in water as above; every four or six hours a teaspoonful.

*Cantharides* is sometimes indicated in the third stage with complete insensibility, cramps in the abdominal muscles and legs, suppression of urine, hæmorrhages from the stomach and bowels, and cold sweat on the hands and feet. It may be employed in water dissolved, every half hour a teaspoonful, until a decided impression is produced.

*Carbo veg.* and *Lachesis* have both proved curative in the third stage of yellow fever, and should always receive due attention in grave cases.

INTERMITTENT FEVER. FEVER AND AGUE. CHILLS AND FEVER.

(*Febris intermittens.*)

The difference between *remittent* and *intermittent* fevers is very marked, and ought to be well understood.

*Remittent* is a fever in which the symptoms continue during the whole time of the patient's sickness, only varying, from time to time, in severity; but never leaving the patient entirely at any time.

*Intermittent*, however, is a fever, in which the symptoms, at certain times, cease entirely, and the patient feels as if entirely free from them, or well. Of this latter class are the chills and fever, or fever and ague, which is caused either by a certain miasma, generated in marshy countries, or follows other diseases, acute as well as chronic; for instance, inflammation of the liver, indigestion, internal obstructions, or peculiar constitutional tendency, etc.

DIAGNOSIS.—The cold stage is preceded by languor, yawning, drowsy headache, numbness of toes and fingers, and blue nails. Then coldness of the extremities is felt, gradually increasing, and sometimes causing the patient to

shake and tremble, his teeth to chatter, and him to become delirious.

During this time, the pulse is weak and oppressed; his thirst variable. The cold stage lasts from twenty minutes to three or four hours, and varies much in severity.

The hot stage shows all the symptoms characteristic of an inflammatory attack: hot, dry skin; thirst; full, quick pulse; congestion of the head and, sometimes, even delirium. Its intensity and duration are variable, the latter being from two to six or eight hours.

The sweating stage indicates an abatement of the fever, although, frequently, there are fever and perspiration at the same time.

TREATMENT.—Certain and general rules for the treatment of this disease can not be given; it depends too much upon the individual, and exciting causes, to make up its general character. But we will try to be as practical on the subject as possible.

Two or three attacks must have passed before we can know, for a certainty, that we have to do with an *intermittent*. These attacks will have been treated already, according to their appearance, either as bilious rheumatic, or gastric fevers, whereby we may have used remedies beneficial in fever and ague. But if this shows itself clearly as such, adopt the following treatment: First, *Ipecac.*, four doses (six glob. each), given in regular intervals between two attacks of fever, during which administer no medicine; but, after it is over, give only one dose of *Nux vomica* (six glob.) If, then, the next attack is lighter, do not give any more medicine, as the disease will disappear, the attacks gradually growing lighter and lighter.

But if this general treatment does not succeed, select the specific remedy among the following, taking great care to find the one most similar to the symptoms of the disease.



For its application, I will here give general directions suitable for every one.

Having found the remedy, give three doses of it (six glob. each), commencing twelve hours before the next attack is expected, and giving a dose once in four hours. During the attack, if its symptoms of chilliness and fever are very severe, give, invariably, *Aconite*, in solution (eight globules in half a teacupful of water), every hour a teaspoonful, until the attack is over. If, then, the next attack is lighter, give no more medicine; if not lighter, select for the next attack another more suitable remedy. In this way proceed for ten or fourteen days, trying to find the right *homœopathic* remedy, before adopting the method spoken of at the end of this article, which subdues the fever, without, perhaps, eradicating it as safely as the *homœopathic* remedy would, if found.

We will give, in the following, the principal remedies useful in domestic practice, leaving more complicated forms of the disease for the treatment of a *homœopathic* physician, if one can be had.

*Arsenic.* For *great debility*, disposition to vomit, or *violent pains in the stomach*; great anguish in the precordial region; lameness of the extremities, or *violent* pains; *imperfect developement* of chilliness and heat, or both, alternately; *drinking very often, but little at a time*. (Compare *China*, *Ipecac.*, *Veratrum*.)

*China.* When, before the fever, there is nausea or thirst, voracious appetite, headache, palpitation of the heart, thirst *between* the cold and hot stages, or *after* the hot stage, or during the perspiration; cold and heat alternately; *no thirst* during the heat; great debility; *uneasy* sleep; *yellow complexion*. (Compare with *Arsenic* and *Lachesis*.)

*Natrum mur.* Violent headache during the heat, with obscuration of sight; pains in the bones, yellowish com-



plexion; fever-blisters on the lips and *ulcerated corners of the mouth*; thirst during the chills, more during the heat; sensitiveness at the pit of the stomach; bitter taste; no appetite. (Compare with *Ignatia*, *Arsenic*.)

*Ignatia*. Thirst only during the chill; gastric symptoms, and eruptions on the lips, as in *Natrum mur.*; *external heat moderates the chill*; external heat, with partial internal shuddering. (Compare with *Carbo veg.* and *Natr. m.*)

*Carbo veg.* Thirst only during the chill; rheumatic pains in teeth or limbs, before or during the attack; nausea; giddiness during the heat; chill comes on in the evening or at night; and, also, appropriate in *damp or wet weather*.

*Sabadilla*. When the attacks return almost at the same hour, with chills of short duration; then thirst followed by heat.

*Ipecac*. Much shivering with but little heat, or the contrary; *increase of the shivering by external warmth* (the opposite of *Ignat.*); oppression at the pit of the stomach; no thirst, or but little, during the chill; violent thirst during the heat; nausea and other gastric symptoms. (Compare *Nux vomica*.)

*Nux vomica*. External heat or internal chill, or *vice versa*; constant desire to be covered, even during the heat; rheumatic pains in the sides, abdomen, and limbs; during the heat, headache, buzzing in the ears, heat in head and face, with redness of the cheeks, and thirst. (Compare *Ipecac*.)

*Pulsatilla*. Gastric symptoms; chills, heat, and at the same time in the afternoon and evening (compare *Sabadilla*); at intervals the patient feels very chilly. It is very suitable when a disordered stomach has caused a relapse, or after *Lachesis*.

*Lachesis*. Chills after a meal or in the afternoon, with pains in the limbs and breast; *violent headache* during the hot stage, with delirium, burning thirst, restlessness, internal

shuddering ; *debility* ; livid complexion during the intervals and when the fever returns easily after eating sour things.

*Veratrum*. External chill and cold sweat, or internal heat with dark red urine ; delirium and red face, or chills with thirst and nausea ; or, vomiting and diarrhea during the heat or in the beginning of the cold stage.

We will now give some general indications and the names of the medicines only. The symptoms may be examined in the *Materia Medica*, at the end of the book, and then the choice made accordingly.

For *marsh-intermittent fevers* : *China*, *Arsenic*, *Ipecac.*, *Carbo veg.*

For *fevers prevailing in damp and cold seasons* : *Carbo veg.*, *Rhus*, *Sulphur*, *China*, *Pulsatilla*, *Lachesis*, *Veratrum*.

For *mismanaged fevers*, by large doses of *Quinine* : *Pulsatilla*, *Arsenic*, *Natrum mur.*, *Lachesis*.

For *daily fevers* : *Arsenic*, *Bryonia*, *Carbo veg.*, *China*, *Ignatia*, *Ipecac.*, *Lachesis*, *Natrum mur.*, *Nux vomica*, *Pulsatilla*, *Rhus*, *Veratrum*.

For *tertian*, or every-other-day fevers : *Arsenic*, *Carbo veg.*, *China*, *Ipecac.*, *Nux vomica*, *Pulsatilla*, *Rhus*.

For *quartan*, or every-fourth-day fever : *Arsenic*, *Pulsatilla*, *Veratrum*.

When the fever returns yearly, give *Lachesis*, *Carbo veg.*, or *Arsenic*.

Having tried to cure the *intermittent* fever with these remedies, for a fortnight, without success, and being unable to procure a homœopathic physician, give the patient the following remedy :

Take four or six grains of *Quinine*, mix with eighteen or twenty grains of loaf sugar, and triturate thoroughly, in a clean mortar, for half an hour ; make six powders of it, and give, twelve or eighteen hours previous to the next attack, one powder ; and, afterward, every two hours one,

until all are given. Give, then, a similarly prepared powder one hour before the next chill is expected to come on, for at least a week, and, afterward, continue to give such a powder every seventh day, in the evening, for thirteen weeks after the last chill. If the fever returns (which it seldom does), use *Pulsatilla*, *Arsenic*, and *Natrum mur.*

N. B. If the disease is complicated with *bilious rheumatic* symptoms, see this article in Index.

DIET AND REGIMEN, as generally in fevers. (See "Common Fever," page 87.)

## CHAPTER V.

### AFFECTIONS OF THE MIND.

SUDDEN emotions are often followed by fatal consequences, either immediately or soon after, and it is not well to neglect them, particularly in children and delicate women.

We will give, in the following remarks, the most frequent emotions, with the principal remedies, whose symptoms are detailed below, and which must be compared with the patient's symptoms before a choice is made.

ADMINISTRATION — All of the following remedies, if selected in a case, should be prepared and given as follows : Dissolve twelve globules of the remedy in half a teacupful of water, and give, every half hour, a teaspoonful, until three or four are given, or until better ; if not better after giving three or four doses, select another remedy, and give it in the same manner. As soon as the patient feels better, give the medicine at longer intervals.

For the consequences of *fright and fear* : *Opium, Aconite, Belladonna, Ignatia, Pulsatilla, Mercury.*

*Excessive joy* : *Coffea, Opium, Pulsatilla, Aconite.*

*Grief* : *Ignatia, Phosphor. ac., Staphysag., Colocynth.*

*Home sickness* : *Phosph. ac., Mercury, Capsicum, Staph.*

*Unhappy love* : *Hyoscyamus, Ignatia, Phosphor. ac.*

*Jealousy* : *Hyoscyamus, Lachesis, Nux vomica.*

*Mortification and insult* : *Belladonna, Ignatia, Platina, Colocynth, Staphysag., Pulsatilla.*

*Contradiction and chagrin* : *Chamomile, Bryonia, Ignatia, Colocynth, Aconite, Nux vomica, Platina, Staphysag.*

If accompanied with *indignation* : *Colocynth, Staphysag.*

*Violent anger* : *Aconite, Nux vomica, Chamomile, Bryonia, Phosphorus.*

SYMPTOMATIC DETAIL. — *Aconite.* Headache, with congestion to the head ; constant fear, especially in children ; or when *Opium* has not been given at once for fright.

*Belladonna.* Loss of consciousness, or constant anxiety, with weeping, fear, *crying with malice*, particularly in children ; also, when *Opium* and *Aconite* have been insufficient for fright.

*Capsicum.* For sleeplessness from home sickness, with heat and redness of the cheeks.

*Chamomile.* For the following consequences of anger : bitter taste in the mouth, nausea, vomiting of bilious matter, *cutting colic, pressure in the stomach*, headache, fever, thirst, restlessness, asthma, *suffocating fits* ; in children, convulsions or gastric derangement after eating or drinking, shortly after an angry fit.

*Bryonia.* Chilliness and shuddering over the whole body ; vehement anger ; loss of appetite ; nausea ; vomiting of bile.

*Coffea.* Nervousness from great joy, with trembling and faintness, especially in females and children.

*Colocynth.* From chagrin or mortification — colic, cramps in the stomach, nausea, bitter vomiting, sleeplessness.

*Hyoscyamus.* From fear — stupefaction and apathy, inability to swallow, convulsions, sudden starts or involuntary laughing during sleep, desire to escape ; and, also,

From unhappy love — jealousy, restlessness, running about.

*Ignatia.* From fright, mortification, chagrin, grief, especially after losing a friend — irresistible grief, headache,

vertigo, vomiting, gastric symptoms or even convulsions, especially in children from fright, or from unhappy love.

*Mercury.* From fright, mortification, or home-sickness—as great anguish, trembling, and restlessness; sudden starting from sleep; congestion of blood on the least motion; sleeplessness; can not bear the warmth of the bed; nervousness; quarrelsome mood; complains of every body, even one's own family; constant shivering; night-sweats.

*Nux vomica.* From anger—particularly after having eaten or drank something, or after *Chamomile* or *Bryonia* has been given without effect.

*Opium.* From joy or fear, to be given *immediately*—loss of consciousness; perspiration about the head, with coldness of the rest of the body; congestion to the head; involuntary *diarrhea*; snoring-like breathing; *fainting fits*; spasms; *epileptic fits*, with spasmodical rigidity of the whole body.

*Phosphoric acid.* From deep grief, unhappy love, home-sickness, or in any case where *Ignatia* is not sufficient, particularly when the patient is dull and listless—when the hair falls out, or turns gray; hectic fever, with profuse sweats; sleepy in the morning.

*Platina.* From anger and mortification—as indifference, sadness, and laughter; pride, with contempt for others; great anguish and dread of death.

*Pulsatilla.* From fear—*diarrhea*, with heat in the abdomen.

From anger—in persons of a bland disposition, or when *Chamomile* is not sufficient.

*Staphysagria.* From anger or indignation and ill humor—the patient pushes violently away whatever is near him; restlessness; fear; also,

From deep grief, as sadness, with a disposition to take everything wrong—dread of the future; sleepy in day-



time and sleepless at night ; falling off of the hair ; feeble and faint voice ; hypochondriac mood.

### HYPOCHONDRIA. MELANCHOLIA.

Hypochondria is a form of diseased mind, which depends mostly on obstructions in the abdomen, and may be cured, if, beside taking the medicines, the patient keeps the general homœopathic diet, and pays attention to the hygienic and hydropathic rules.

*Nux vom.* Ill humor ; *aversion to life* ; *disposition to vehemence* ; disinclination to mental labor ; unrefreshing sleep ; *feels worse in the morning* ; dullness of the head, with aching pains, or sensations as if a pin were sticking in the brain ; aversion to the open air ; *constant desire to lie down* ; great exhaustion after walking ; painful distention in the hypochondria and stomach ; *constipation* of the bowels ; disposition to, or presence of the piles (hemorrhoids.) *Sulphur* follows well after *Nux vom.*

*Sulphur.* Lowness of spirits ; solicitude about one's affairs, health, salvation ; fixed ideas ; anxious impatience ; restlessness ; vehement disposition ; *bodily and mental indolence* ; absence of mind ; can not think ; exhaustion after mental labor ; fullness and pressure in the stomach ; *constipation* ; feels very unhappy. (*Calcarea carb.* follows well after *Sulphur.*)

*Calcarea carb.* Lowness of spirits, with disposition to weep ; anxiety, with congestion or palpitation of the heart ; despair about one's health ; apprehension of illness, misfortune, infectious diseases, insanity, etc. ; dread of death ; can not think, or perform any mental labor. (*Natrum m.* follows well after *Calcarea carb.*)

*Natrum mur.* Lowness of spirits ; weeping and lamenting about the future ; *desire to be alone* ; aversion to life ; ill humor ; disposition to vehemence ; can not perform mental



labor; headache, with want of appetite; indigestion after eating.

ADMINISTRATION.—If one of the above remedies is chosen, give every third evening a dose (six glob.), until four or six doses are taken; then discontinue for the same length of time; giving afterward the next best remedy in the same manner; and so on, until better.

#### MELANCHOLIA.

A state of mind similar to the former, but more deeply seated, with derangement of the system, often requiring skillful medical aid. However, try in addition to the remedies indicated under “Hypochondria,” *Lachesis*, *Belladonna*, *Ignatia*, and *Pulsatilla*. (See “Materia Medica” at the end of the book.”)

## CHAPTER VI.

### AFFECTIONS OF THE HEAD.

#### 1. CONGESTION, OR DETERMINATION OF BLOOD TO THE HEAD.

(*Congestio ad caput.*)

SOME persons have a predisposition to this disease ; others acquire it by wrong habits, such as the use of spiritous liquors, leading a sedentary life, or by intense mental application. It becomes dangerous by long continuance, in which case the patient ought to put himself under the charge of a homœopathic physician.

For the single attacks, we will give the remedies below.

DIAGNOSIS.—In such an attack, the beating of the arteries of the head is felt by the patient, the veins of the neck are swelled, the head feels full and heavy, and giddiness ensues, particularly when stooping, exercising, or walking in the sun, and when turning round suddenly. The head aches mostly above the eyes ; increased by stooping, coughing ; sleep is unrefreshing ; drowsy.

TREATMENT.—In general, the patient ought to abstain from heating or ardent drinks, such as coffee, liquors, and tea ; but drink cold water freely, and wash or bathe in it. Every evening take a cold foot-bath up to the ankles, rubbing the feet briskly after it. Never allow bleeding, as it would only increase the disposition afterward, without tak-

ing away the present attack sooner than the following remedies will do, if properly applied.

The principal circumstances, under which congestion to the head takes place, and their remedies, are :

From *great joy* : *Coffea*, *Opium*.

From *fright or fear* : *Opium*.

From *anger, violent* : *Chamomile*, *Bryonia*, *Nux vom.*

From *anger, suppressed* : *Ignatia*.

From a *fall or blow, causing concussions* : *Arnica*.

From *loss of blood or fluids* : *China*.

From a *cold* : *Dulcamara*, *Bryonia*, *Rhus tox.*

From *constipation* : *Opium*, *Nux vom.*, *Bryonia*.

From *drinking liquors* : *Nux vom.*, *Opium*.

From *sedentary habits* : *Nux vom.*, *Aconite*.

At the *critical periods of girls*, or

From the *stoppage of menstruation* : *Pulsatilla*, *Aconite*, *Belladonna*, *Bryonia*, *Veratrum*.

From *dentition in children* : *Aconite*, *Coffea*, *Belladonna*, *Chamomile*.

Before giving one of the above remedies, examine the symptoms, as detailed below.

*Aconite* is the first remedy to be given, if the head aches violently above the eyes, *as if it would burst*, particularly when stooping or coughing, or when it is caused by *fright or anger*, especially in children. (See "Affections of the Mind.")

ADMINISTRATION the same as in "Affections of the Mind," page 110.

*Belladonna*. After *Aconite*, if necessary, or in alternation with it, if the congestive symptoms are more on one side of the head, and particularly the *right*; or if there is violent pressure in the forehead, increased by motion, stooping, noise, or the glare of light; darkness before the eyes; buzzing in the ears. Administration the same as *Aconite*.

N. B. *Belladonna* and *Aconite* for young girls at the critical age. (See *Pulsatilla*.)

*Opium*. First remedy, if occasioned by fright (see "Affections of the Mind" under *Opium*), or in those serious cases of sudden congestion which occur in summer from a draught of cold water when heated, with the following symptoms: vertigo; buzzing in the ears; *stupor*, with sweat pouring off the head; pulse full and heavy, or quick and weak.

Also, in those cases which arise from extreme *constipation* when the patient has no desire for stool (*Nux vom.* in constipation, with ineffectual effort to evacuate the bowels), with red and bloated face; dry mouth; wandering look; if, with acid regurgitations, nausea, and vomiting, alternate with *Nux vom.*; the same, also, when from *debauch*. Administration the same as *Aconite*.

*Coffea*. In cases of congestion caused by excessive joy, or in children when teething; sleeplessness, and agitation; sometimes in alternation with *Belladonna* or *Opium*; administer same as *Aconite*.

*Chamomile*. Congestion caused by angry passion, particularly in children when teething; followed by *Bryonia* or *Nux vom.*, if necessary. Administer the same as above.

*Ignatia*. Congestion from *suppressed anger*, or *grief*. Administer same as above.

*Arnica*. By a fall on the head or other parts; congestions of the head take place either by actual concussion, or by fright and fear occasioned by a fall. In both cases, *Arnica*, externally applied (twelve drops of the tincture in half a teacupful of water), in wet bandages; and internally, as stated above under *Aconite*.

Also, in cases where there is heat in the head, with chilliness of the other parts; dull pressure in the brain; buzzing

in the ears, and vertigo; obstruction of sight, especially when rising from lying down. Administer same as *Aconite*.

*Nux vom.* Congestion from sedentary habits; intense study, or drinking spirituous liquors; also from violent anger (see *Chamomile* and *Bryonia*), when there is nervousness, with painful feeling in the head when walking or moving; pressure in the temples; dim eyes, with desire to close them without being able to sleep; the symptoms worse in the morning, in the open air, or after a meal; bowels constipated, with ineffectual desire to evacuate. Administer as above.

N. B. In cases where *Nux vom.* does not give perfect relief, in persons fond of liquors, *Opium*, *Calcarea carb.*, and *Sulphur*, at intervals of four or six days, one dose (six glob.), are necessary. (See "*Delirium Tremens*." )

*Pulsatilla.* Congestion at the critical period of young girls, or in phlegmatic temperaments, when there are distressing pressure and pain on one side (particularly the *left*) of the head; relieved in the open air, or from binding the head; whining mood; anxiety; shivering; pale face, with vertigo. Administer same as *Aconite*.

*Mercury.* Congestion with sensation of fullness, or as if the head were compressed by a band; relieved by the pressure of the two hands around the head; worse at night; disposition to perspire—is often suitable after *Belladonna* and *Opium*. Administer same *Aconite*.

*Dulcamara.*—Congestion from the least cold; wet feet, particularly when there is continual buzzing in the ears, with dullness of hearing. Administer same as *Aconite*.

*China.* Congestion after loss of fluids, as blood-letting, or hemorrhage; when there is beating in the head, with palpitation of the heart, and dimness of sight; heavy breathing when lying with the head low; followed, in such

cases, by *Calcarea carb.* and *Veratrum*. Administer same as *Aconite*.

*Bryonia*. Painful pressure in the temples, or sensation as if everything would fall out of the forehead when stooping; nose bleeds without relief; constipation. If in wet weather, alternate with *Rhus*. Administer every two or three hours a dose (four glob.), until three doses of each are given, or till better.

*Rhus tox.* Congestion with burning, throbbing pains, and fullness in the head; rheumatic pains in the limbs; worse after eating. (See *Bryonia*.) Administer as *Bryonia*.

DIET AND REGIMEN.—The diet ought to be light, farinaceous; no meat or broths, except in cases where loss of blood or fluids has preceded the congestion.

In the habitual form of congestion, early rising and daily exercise in the open air, bathing or sponging in cold water, are recommended.

## 2. VERTIGO. GIDDINESS OR DIZZINESS.

It arises either from disordered stomach, irritation of the brain and congestion to the head, or from suppressed eruptions, or ulcers; also from riding in a carriage.

In most cases it accompanies severe disorders of the brain and stomach, when it ought to be treated by an experienced homœopathic physician. Until such help can be had, the following remedies may be resorted to:

TREATMENT.—Vertigo from *disordered stomach*: *Nux vom.*, *Arnica*, *Chamomile*, *Pulsatilla*, *Antimon. crud.*, *China*, *Rhus*, *Cocculus*.

*Irritation of the brain*, from excess of thinking, etc.: *Nux vom.*, *Arnica*, *Belladonna*, *Chamomile*, *Hepar*, *Pulsatilla*, *Rhus*.

*Suppressed ulcers and eruptions*: *Calcarea carb.*, *Sulphur*,



*Riding in a carriage: Hepar sulph., Silicea, 'Cocculus, Petrol.*

*Congestion of blood to the brain: Aconite, Bryonia, Arnica, Belladonna, China, Lachesis, Mercury, Nux vom., Opium, Pulsatilla, Sulphur.*

Before giving one of the above remedies, if selected, read its detailed symptoms below.

ADMINISTRATION of these medicines. Every two, three, or four hours, one dose (four glob.) until relieved, or another remedy is necessary. If in vertigo, caused by congestion to the head, see "Administration," in this article.

*Aconite.* Vertigo on raising the head after lying down or stooping, attended with nausea, vomiting, dimness of sight, loss of consciousness.

*Antimon. crud.* Nausea and vomiting; aversion to food; disordered stomach; is followed well by *Pulsatilla*.

*Arnica.* Vertigo from over-eating, or during meals, with nausea, dimness of sight, red face; is followed well by *Nux vom., Chamomile, Rhus, Lachesis.*

In cases of this kind, frequently very dangerous, seek the advice of a homœopathic physician.

*Nux vom.* Vertigo *during or after a meal, or when walking in the open air; stooping, or thinking;* worse in the morning or evening, *in bed, when lying on the back, with sensation as if turning around or falling; fainting; loss of consciousness.*

*Opium.* From fright, with trembling; stupor; dimness of sight; worse when rising from bed, and obliged to lie down again.

*Pulsatilla.* Vertigo when *looking upward, or when sitting or stooping;* worse in the evening, or after a meal; better in the open air.

*Rhus.* Vertigo when lying down in the evening, with fear of falling or dying.



*Lachesis.* Vertigo on *walking* in the morning, with absence of mind; intoxicated feeling; constipation; sensation as if the head were turning to the left side.

*Mercury.* Vertigo on arising in the morning, or in the evening, with nausea; dimness of sight; or when raising the head with a desire to lie down again. (See *Bryonia*.)

*Bryonia.* Same as *Mercury*, and with feeling of intoxication and congestion when stooping and rising again.

*Hepar sulph.* Vertigo from riding in a carriage.

*Cocculus.* Same as *Hepar sulph.*, if this is not sufficient.

*Petroleum.* Same as *Cocculus*, if that is not sufficient.

*Silicea.* Same as *Petroleum*, if this is not sufficient, or the vertigo seems to rise from the back to the nape of the neck, and thence to the head; worse in the *morning* and *after emotion*.

*Sulphur.* Vertigo in the *morning*, or at *night*, when *ascending an eminence*, or after a meal; after suppressed ulcers and eruptions.

*China.* Vertigo, with fainting when raising the head. (See "Congestion to the Head.")

*Calcarea carb.* Same as *Sulphur*, and when with *trembling before breakfast*; *coldness of the head*; *congestion to the head*.

*Chamomile.* Vertigo in the morning or after eating, and drinking coffee, with *fainting turns*.

DIET AND REGIMEN the same as in "Congestion to the Head."

### 3. WEAKNESS OF MEMORY.

This diseased state of the brain is principally caused by *debilitating loss of fluids*, such as bleeding, purging, sexual excesses, etc., for which give *China*, *Nux vom.*, and *Sulphur*, every week one or two doses (six glob.), in alternation, till better.

Sometimes it is caused by *excess of mental labor*, for which give *Nux vom.* and *Sulphur*, as above.

If caused by *external injuries*, a fall or blow, give *Arnica*. (See "Congestion to the Head.")

If by *congestion to the head*, give *China*, *Mercury*, *Rhus*, and *Sulphur*. ("See Congestion to the Head.")

If by *excessive use of spirituous liquors*, give *Nux vomica*, *Calcarea carb*, *Lachesis*, *Opium*, *Sulphur*, in the same manner as above, under "Debilitating Loss, etc."

DIET AND REGIMEN as in "Congestion to the Head."

#### 4. APOPLEXY. (*Apoplexia*.)

Short, thick-necked persons, of a full and fleshy growth, are predisposed to this disease; it is also hastened on by the too free use of spiritous liquors, by excesses of any kind, and suppression of the habitual perspiration of the feet. An attack of apoplexy is often preceded by heaviness and fullness of the head; buzzing in the ears; dim eyes; dullness of hearing; sleepiness; indistinct speech; numbness of limbs. If this state exists, particularly in old persons, or those predisposed to apoplexy, the greatest care should be taken in the diet, which ought to consist only of simple farinaceous and mucilaginous substances, and cool acidulated drinks; they ought not to exercise very much, particularly in the heat of the day, and should take the following remedies, as it is easier to *prevent* an attack than to cure it:

*Opium*. If in *old people* there is unusual stupor; buzzing in the ears; obtuseness of hearing; redness of the face; constipation; pulse slow and full. Give three times a day a dose (four glob.), until better.

*Nux vomica*. In people of a sedentary habit, or addicted to the use of ardent spirits, in alternation with *Opium* and *Lachesis*, particularly when, with the above premonitory

symptoms, there is headache on the right side, with vertigo and a nervous bilious temperament.

ADMINISTRATION same as under *Opium*.

*Lachesis*. Same as *Nux vom.*; but particularly when the left side of the head suffers, and *lowness of spirits* is connected with it. Administration same as above.

*Belladonna* and *Aconite* under the same circumstances as stated under "Congestion to the Head."

*Arnica*. See the same article under "Congestion to the Head after a Meal." If an apoplectic fit ensues, however, continue the same remedies as indicated above, only give them in solution (twelve globules to half a teacupful of water), every half hour a teaspoonful until better, or another remedy is indicated.

In such cases hasten immediately to procure a homœopathic physician. Avoid bleeding under any circumstances; the chances of killing the patient by such an act are too frequent.

DIET AND REGIMEN as in "Congestion to the Head."

## 5. INFLAMMATION OF THE BRAIN.

The brain and its two coverings (see Anatomical Part) are subject to inflammation separately, the characteristic symptoms of which show themselves, particularly in the beginning of a brain fever. We will, therefore, give them separately, as it is important, as well as interesting, to be able to distinguish between the different localities of the disease.

DIAGNOSIS. *First species*, called *meningitis*, if the *dura mater*, the skin next to the skull, is inflamed.

The *acute meningitis* attacks a person suddenly, exhibiting, *immediately*, stupor and drowsy symptoms; the patient can not be raised easily, or keep his head erect; it inclines to fall on either side; complains of no pain, except vertigo

and dizziness ; reels when walking ; pupil contracted ; sensitiveness to light, although the eyes are not red as yet ; constipation ; scanty, dark, red urine ; fever, with a soft pulse.

N. B. Old people are most liable to this species of inflammation of the brain ; also, those who have indulged too much in the use of spiritous liquors, particularly after taking cold. It frequently precedes an apoplectic fit.

*Second species.* Called *arachnoiditis*, if the covering next to the brain is inflamed. This species does not commence with stupor, and not without pain, like the first species, but with violent pains over the entire head ; increased by paroxysms of congestion, with violent beating of the arteries of the neck and temples ; sleeplessness ; low muttering ; pulse quick ; if stupor ensues suddenly, the patient dies soon after.

*Third species.* *Encephalitis vera*, or inflammation of the substance of the brain, in its various parts.

It commences with violent congestion and pulsation of the arteries on the neck and head ; face red and bloated ; eyes bloodshot and brilliant ; intense heat of the head, and violent delirium.

The patient tosses about, screams, and evinces great muscular strength and sensitiveness to light ; pupil contracted ; can not hear well. The deeper the interior of the brain is affected, the more the senses are stupified, so that, in some cases, the patient can, finally, neither see nor hear ; dry skin ; violent thirst ; hard, full pulse ; fever accordingly ; frequent vomitings or retchings.

TREATMENT.—It is easily to be conceived that, in such a violent disease, no layman would trust his own judgment, except in cases of absolute necessity ; and, on this account, we give the following treatment of the above three species of brain fever, which is the same as regards the *internal*

remedies, and differs only in the *external applications*, as will be seen below.

EXTERNAL APPLICATIONS. — *First species* — cold-water bandages.

*Second species* — hot-water bandages.

*Third species* — cold-water bandages, or pounded ice.

INTERNAL TREATMENT. — *First species* — *Opium*, *Hyoscyamus*, *Bryonia*.

*Second species* — *Aconite*, *Belladonna*, *Hyoscyamus*.

*Third species* — *Aconite*, *Belladonna*, *Stramonium*, *Bryonia*, *Hyoscyamus*.

Before giving the medicine selected, read the symptoms presented below.

*Aconite* is generally given first, when there is inflammatory fever, delirium, violent burning pains through the whole brain, particularly in the forehead; red face and eyes; hot and dry skin. Six hours after giving *Aconite*, give *Belladonna*, or continue the use of *Aconite* in alternation with *Belladonna*, until better.

ADMINISTRATION. — See “Affections of the Mind,” under *Aconite*.

*Belladonna*. This remedy is almost a specific in inflammation of the brain, particularly when there is *great heat of the head*, red and bloated face, *violent pulsation of the arteries* on the neck and temples; *burying the head in the pillow*; sensitiveness to the *slightest noise and light*; burning and shooting pains in the head; *bloodshot and brilliant eyes*, with a wild expression; contracted or dilated pupils; *violent and furious delirium*; raving; loss of consciousness; sometimes low mutterings, convulsions, vomiting, violent thirst. Administer alone or in alternation with *Aconite* (see above), or with *Stramonium* and *Hyoscyamus* (see below), to be given in solution as above in *Aconite*.

*Hyoscyamus*. Stupor, loss of consciousness, delirium;

sudden starting ; talks about his affairs ; inarticulate speech ; singing, muttering, smiling, picking the bed clothes ; skin dry ; red face ; desire to escape ; involuntary stool and urine.

Administration, see *Belladonna*.

*Stramonium*. Red face ; staring look ; *frightful visions* ; sleep natural, but with twitching, tossing about, and absence of mind when awaking ; desire to escape, and screams.

Administration same as *Belladonna*.

*Bryonia*. Constant stupor, with delirium ; sudden starting from sleep, screams, and cold sweat on the forehead ; burning and shooting pains in the head, or stitches through it.

Administer as *Belladonna*.

*Opium*. Stupor, with heavy breathing ; eyes half open ; confusion and giddiness after waking ; complete listlessness and dullness of sense ; also, no desire for anything, or complaining of anything.

Administer same as *Belladonna*.

DIET AND REGIMEN as in "Common Fever."

N. B. As the brain of children is very tender, its diseases must necessarily be more dangerous and frequent, and we give, therefore, in the following, a more special and somewhat modified treatment for inflammations of the brain in children :

If a *child's head is very hot, feet cold*, skin dry and hot, it *sleeps with eyes half open*, has scanty urine, with or without vomiting or diarrhea, let the cause of the disease be what it may, in such a case give at once *Belladonna* and *Hellebor.*, in solution (twelve globules in half a teacupful of water, each remedy in a separate cup), every two hours a teaspoonful, alternately, until better, or until the symptoms require a change of remedy, generally *Sulphur* or *Mercury*. (See *Materia Medica*, at end of book.)



The above suggestion is *important*, as cases of inflammation of the brain occur so often among children, when, as is said, the disease goes to the head. For instance, in teething; in common cold, with fever; in scarlet fever (see this article); or in other eruptions, when suppressed; also, worm fevers.

If such a state should continue in a child for some time *dropsy of the brain* (see "Hydrocephalus") would speedily ensue; the above treatment will surely prevent this, if applied in time.

N. B. If, at any time during the treatment of inflammation of the brain, the patient is very restless and sleepless, we may give, with great advantage and without injury to the effect of the other medicines (which are discontinued during that time), *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, alternately, every hour a dose (four glob.), until better. This may be done often in the night.

## 6. SUNSTROKE

Is a species of inflammation of the brain, which requires the following

TREATMENT. — *Camphor*. Let it be put on the patient's tongue, at intervals of five or ten minutes, if he is sweating profusely while unconscious, and breathing heavily.

*Belladonna*, after *Camphor* has been given, in alternation with

*Lachesis*, every half hour, in solution (twelve globules of each in separate teacups half full of water), a teaspoonful at a time.

*Externally*, wash the head and face occasionally with brandy, and give, internally, small quantities of brandy until the patient revives. Put neither cold water nor ice on the head.



## 7. HEADACHE.

This disease, so frequent and troublesome, requires to be treated with particular minuteness; the more so, as its treatment falls almost entirely within the legitimate sphere of domestic practice, except where headaches are the concomitant symptoms of severe fevers, as in the typhus, etc.; or the forerunner of dangerous attacks, as in apoplexy, etc.; in which cases, the respective articles must be consulted.

ADMINISTRATION OF MEDICINE IN HEADACHE.—For this purpose, we divide all headaches into *habitual* and *accidental*. This requires two modes of administering the medicine: one for the cure of the *present* attack, if it be accidental, or one of the habitual attacks; and the other, for the eradication of the habitual headache, or the predisposition to it.

First. Administration of medicine *for an attack*.—Having selected a remedy (from those stated below), dissolve of it six globules in three tablespoonfuls of water, and take, according to the severity of the headache, from half an hour to two, three, or six hours, a teaspoonful, until three are taken, when its benefits must be apparent. If beneficial, continue with the same, only at much longer intervals, as long as necessary. If not beneficial, select carefully another remedy, and prepare and take in the same manner.

Secondly. Administration of medicine for *eradicating the disposition* to certain kinds of headache; for instance, sick-headache.—Having selected a remedy, take of it, every third or sixth evening, a dose (four glob.), until four or six doses are taken; when the next best remedy has to be taken in the same manner, or until better.

*a. Headache from Congestion of Blood to the Head.*

*Aconite.* Violent throbbing; heaviness; fullness in the forehead and temples, with a sensation as if the head would

burst ; burning pain through the whole brain ; red, bloated face, and red eyes ; sensitiveness to light and noise ; worse on moving, talking, drinking, rising up ; full and quick pulse. After it, or in alternation with it, *Belladonna* may be given, if necessary. (See “Congestion of Blood to the Head.”)

*Belladonna.* Violent aching pain, as if the head would split, or as if the brain would protrude through the forehead ; heat in the head and coldness of the feet ; undulating feeling, as of water in the forehead ; violent throbbing of the arteries of the neck and temples ; delirium, either with red, bloated face, bloodshot eyes, great sensitiveness to light, noise, and touch, or vertigo and deeply-seated, violent pressing pains, with pale face, drowsy unconsciousness, (compare, in such a case, article on “Apoplexy”) ; worse on moving the eyes, or raising or moving the head. In alternation with *Aconite* or *Rhus*, if the symptoms agree. (See *Aconite* and *Rhus*.)

*Bryonia.* Distending pressure from within, particularly through the forehead, on stooping or moving, with violent beating or single stitches in the head ; desire to lie down ; constipation of the bowels ; bleeding at the nose. Before it, *Belladonna* frequently suits ; after it, *Rhus*. (See these remedies.)

*Rhus.* Fullness of the head, with burning and throbbing pain ; oppressive weight in the back part of the head, with a sensation as if a fluid was rolling inside ; a feeling of looseness, particularly in wet weather. In alternation with *Belladonna* or *Bryonia*.

*Pulsatilla.* Oppressive, dull pains on one side of the head, commencing in front at the root of the nose, and going back, or vice versa ; better in the open air, or by compressing the head, or lying down ; worse toward evening, or when sitting or looking upward ; pale face, agitated mind,

*inclined to weep.* It suits *females* and persons of a *mild* character, and *lymphatic* temperament.

*Nux vomica.* The opposite of the former. Pains; worse in the *morning* and in the *open air*; heaviness of the head, especially when *moving the eyes* or *thinking*; a sensation as if the skull would split; *contusive pain in the brain*, worse when stooping or in motion; *constipation*, with rush of blood to the head; *irritable, rash, or lively* temperament, full habit, and for persons of *sedentary* habits, or who use *coffee* and *ardent spirits* freely. (Compare *Bryonia*, *Belladonna*, and *Chamomile*.)

*Mercury.* Fullness of the head, as if it would fly apart, or was tied up with a bandage; better by *pressing* the head with the hands, but worse at night, in the *warmth of the bed*, and not relieved by profuse sweating.

*Opium.* *Constipation, with rush of blood to the head*; violent tearing, pressing pains through the whole brain, and heaviness with beating in the head; unsteady look. (See "Congestion to the Head.")

*b. Headache from Catarrh and Cold in the Head.*

The principal remedies are, *Nux vomica*, *Mercury*, *Sulphur*, *Aconite*, *Chamomile*. (See, for other remedies, "Catarrh.")

*Nux vomica.* Heaviness in the forehead; obstructions or running of the nose; feverish heat of the head, cheeks, and body, alternately with chilliness, constipation. After it, *Mercury* often suits.

*Mercury* is particularly useful in those headaches accompanying the epidemic catarrh (*influenza*), when there are *pressing* pains in the forehead, over the root of the nose; frequent sneezing and running at the nose, with redness and excoriation; also, with painful itching in the nose;

fever, with chilliness, pains in the limbs, and thirst. It alternates well with *Nux vomica* or *Sulphur*.

*Sulphur*. After *Mercury* has relieved, but not entirely cured the cold, and when there is still left fullness, pressure and heaviness in the forehead, stitches, and painful jerking; especially on the *left side*, with heat and rush of blood to the head; constipation.

*Aconite*. Pressing, dull feeling and heat in the forehead; worse at night; better in the open air; fever, intermixed with chills; running of the nose and eyes.

*Chamomile*. Intolerable tearing and jerking on one side of the head, down to the jaws, with sore throat, hoarseness, and bitter, foul taste in the mouth. Is frequently suitable after *Aconite*.

### c. Headache from Rheumatism.

The principal remedies are *Chamomile*, *Colocynth*, *Ipecac.*, *Pulsatilla*, *Bryonia*, *Mercury*, *China*, *Nux vomica*.

*Chamomile*. Drawing, tearing pains on one side of the head, down to the jaws; very sensitive to touch; hot sweat about the head; one cheek red, with paleness of the other; and if the pain almost drives to despair. Suits well after *Coffea*: after it, *Bryonia*, *Pulsatilla*, and *China*, are suitable.

*Colocynth*. Violent tearing, drawing, cramping, or aching pains, with nausea and vomiting; worse when stooping or lying on the back; pains appear in the afternoon, with restlessness.

*Ipecac*. Drawing in the forehead; worse or excited by touch; headache, with nausea; sensation through the skull, extending to the tongue, as if the brain were bruised; pains better after vomiting; relieved by heat.

*Nux vomica*. Tensive, drawing pains on one side of the head, with a bruised sensation in the head; worse when

stooping or in the *open air*, with nausea and sour vomiting ; constipation.

*Pulsatilla.* *Darting, rending, jerking* pains on one side only, particularly in the temples ; pain as if the *brain were lacerated*. (See “Nervous Headache.”)

*Bryonia.* *Rending and shooting* pain, from the neck up to the sides of the head, with shivering or fever ; worse by *motion* and *at night*, or during changeable weather. In alternation with *Rhus*.

*Mercury.* *Burning, shooting, throbbing*, and *rending* pains, on one side principally, down to the teeth and neck, and in the ears ; worse at night and in the *warmth of the bed*, with perspiration which does *not* relieve.

*China.* Aching pains at night, with sleeplessness, *jerking, tearing* ; boring on top of the head, with a bruised feeling of the brain ; worse, in the open air, by touch, motion, draughts of air and wind ; sensitiveness, even of the *roots* of the hair, to *touch*.

*d. Headache from Constipation and Gastric Derangement.*

1. *Constipation*, which causes congestive headache, is generally cured by *Bryonia*, *Nux vom.*, or *Opium*, according to its concomitant symptoms (see these three remedies, under “Headache from Congestion to the Head,” p. 128), or by *Pulsatilla* and *Mercury*. (See their symptoms, under the same heading.)

But, if the constipation is of a *chronic character*, give

*Silicea*. When there are *beating* pains in the head, mostly from congestion, from the nape of the neck to the top of the head ; pressing in the head, *as if it would split*, with sleeplessness at night, and worse in the morning when awaking, or when reading, writing, or thinking ; constipation, with ineffectual urging. Suits, frequently, after *Lycopodium*, or

*Lachesis* and *Lycopodium*, in alternation, when there is cutting as with knives in the top of the head, or hammering on stooping, with congestion to the head; disposition to faint, and great restlessness; whining mood; yellowish complexion.

*Sepia*. Headache on shaking or moving the head, or walking, with tendency of blood to the head, heaviness, and confusion, with nausea and vomiting; worse in the morning and on looking at bright sunlight.

*Sulphur*. If the former remedies do not cure, or when the headache is more on one side (the *left*), with heat and congestion; roaring in the head; worse by thinking, in the open air, or walking, or when it appears every morning, or night, or every week.

2. *Headache from Gastric Derangement* is, generally, accompanied by a furred tongue, loss of appetite, fullness in the forehead, as if it would split, with beating pain; worse when stooping. It is frequently connected with bad taste in the mouth, and nausea or vomiting, and *sour risings* (*acid stomach*) caused by indigestion. (See these articles.) The principal remedies are,

*Antimonium*. When the pain is worse on going up stairs, and better in the open air. With nausea, loathing, and aversion to food, *Bryonia*, *Pulsatilla*, *Nux vomica*. (See "Headache from Congestion" and "Indigestion.")

#### e. *Headache from External Causes.*

There are many external circumstances that produce headache; we will state the most frequent of them, with their remedies, whose detailed symptoms must be consulted under the headings mentioned within the brackets, and at the end of the book, in the "Materia Medica."

a. Headache from drinking coffee—*Nux vomica*, *Chamomile*, *Ignatia*.



b. From *over heating* — *Aconite*, *Belladonna*, *Bryonia*.  
(See “Congestion to the Head.”)

c. From drinking *ardent spirits* — *Nux vomica*, *Arsenic*,  
*Lachesis*.

d. From *loss of sleep, long watching* — *Nux vomica*, *Cocculus*, *Pulsatilla*, *Lachesis*, *Carbo veg.*

e. From abuse of *tobacco* — *Hepar sulph.*, *Nux vomica*,  
*Antimon. crud.*

f. From excess of *mental or bodily labor* — *Nux vomica*,  
*Sulphur*, *Lachesis*, *Lycopodium*, *Silicea*.

g. From *grief* — *Ignatia*, *Staphysag.* (See “Mental Diseases.”)

h. From *anger or chagrin* — *Chamomile*, *Nux vomica*.  
(See the same.)

i. From *bad, changeable weather* — *Bryonia*, *Rhus*, *Carbo veg.* (See “Headache from Cold.”)

j. From *bathing* — *Pulsat.*, *Antimon crud.*, *Calcarea carb.*

k. From *suppressed eruptions* — *Antimon. crud.*, *Sulphur*.  
(See “Skin Diseases.”)

l. From *mechanical injuries, blows, etc.* — *Arnica*, *Bellad.*,  
*Aconite*. (See “External Injuries.”)

N. B. In regard to the administration of the above medicines, we refer the reader, in all cases of headache, to what is said about it in the beginning of that article, p. 128.

#### SICK HEADACHE.

This form of headache, so well known under the above name, is of a chronic nature, appearing periodically, and depending mostly upon gastric, rheumatic disorders in persons of a psoric constitution, and may be eradicated by a systematic homœopathic treatment. Although this requires the skill of medical attendance, we will try to bring within the reach of every intelligent person sufficient knowledge



for the commencement of a treatment, which, if successful, will encourage him to seek further medical advice.

First, we will give the treatment of the attack itself:

*Ipecac.* is the first remedy, in all cases which commence with nausea, after which a bruised feeling of the whole head manifests itself, followed by vomiting or retching.

ADMINISTRATION.—Of this and the following remedies as directed in the beginning of the article, page 128. If not better within an hour, select one of the following remedies:

*Belladonna.* If with the headache there is great sensitiveness to light, worse in the warmth of the bed, on lying down or in a draught, and from noise, such as shutting the door, etc. (See “Congestion to the Head.”)

*Spigelia.* If there is great sensitiveness to noise, with the pains worse on the *left* side, and insupportable beating in the temples, the pain sometimes descending into the face and teeth, worse by stooping or motion and in the open air. The headache appears generally at a *regular* time in the morning, growing worse with the ascending sun (*sun pain*.) In this case, in alternation with *Belladonna*.

*Sanguinaria.* If there is great sensitiveness to the *walking* of others in the room, with fullness of the head, as if it would burst, worse on the *right* side, with a feeling as if the eyes were pressed outwards, chilliness, nausea, inclination to lie down.

(Compare *Bryonia* in “Congestion to the Head.”)

*Aconite.* (See this remedy in “Congestion to the Head.”) When the patient has a great *sensitiveness to all kinds of odors*, the pain mostly on the *left* side. In this case in alternation with *Sulphur* or *Ignatia*. (See “Nervous Headache.”)

*Sepia.* If there is a great dislike to be *touched*, and sensitiveness to thunder-storms and the cold air. (Com-

pare with *China* in "Headache from Rheumatism.") If the patient is easily vexed, pain mostly above the right eye and worse by shaking or moving the head.

Beside, consult the articles on "Congestion to the Head," "Headache from Rheumatism," and "Neuralgia in the Head."

Second. To eradicate the disposition to sick headache, take *Sepia*, *Silicea*, and *Sulphur*, each remedy for six weeks, commencing with *Sepia*. During the six weeks take of that remedy, the first three weeks, every week, two doses (six glob.) in the evening, on going to bed; for the last three weeks, every week one dose (six glob.) If during that time attacks of sick headache occur, treat them as directed above and resume the other treatment again after they are over.

#### NERVOUS HEADACHE. NEURALGIA IN THE HEAD. (*Megrim.*)

In this kind of headache, to which nervous persons and females are mostly subject, the head is generally cool, the face pale, with a suffering expression; in females great quantities of colorless urine are discharged; if vomiting sets in, it only relieves for a short time; the pain is mostly concentrated in one spot, with the characteristic feeling, as if a *nail were driven through the head* (*clavus hystericus*); the patient is generally very nervous, fickle minded, dejected, hysterical, or hypochondrical.

TREATMENT.—Besides the medicines mentioned under "Sick Headache," which are suitable in this kind of headache, also, (see this article), there are the following:

*Coffea*. Pain as if a nail was driven into the head, or as if the brain were torn or bruised; pain seems to be intolerable; also, noise and music; the patient is very restless, screams, weeps, feels chilly and has an aversion to open air and drinking coffee, though he is, at other times,

fond of it. The headache generally arises from cold, close thinking, or vexation.

*Ignatia.* Aching pain above the nose, relieved by bending forward; sensation as if a nail had been driven into the head; with nausea, dimness of sight, and yet dread of light; pale face, copious and watery urine; *pain is momentarily relieved by a change of position*; tendency to start; fitful mood; taciturn and sad.

*Pulsatilla.* Tearing pains, worse toward evening, with pale face, whining mood, loss of appetite, *no thirst*, chilliness, sometimes with palpitation of the heart; pains are worse during rest, when *sitting*, better in the *open air*, and when the head is bandaged tight; particularly suitable for *mild phlegmatic* persons.

*Aconite.* (See "Sick Headache.") When it arises from cold, with catarrh, buzzing in the ears and pains in the abdomen; also, when there is a sensation as if a ball was rising up into the head, as if of cool air.

*Belladonna.* (See "Sick Headache.") When there are wavering shocks, and a sensation of fluctuation or undulation, as of water. This pain sometimes commences very *gently*, but increases rapidly on one side (mostly the *right*) to piercing pains, which produce agonizing lamentations and temporary delirium, dimness of sight, buzzing in the ears.

*Platina.* Suits often after *Belladonna*, particularly in females, when there is roaring in the head as of water, with coldness in the ears, eyes, and one side of the face; also illusions of sight, as objects appearing smaller than they really are; violent cramp-like pains over the root of the nose, with heat and redness of the face, restlessness, and whining mood.

*Hepar sulph.* Suits after *Belladonna* or *Mercury*, if there is pain as from a nail in the brain, with violent nightly

pains, as if the forehead would be pulled apart, and when painful tumors appear on the head.

*Mercury.* After *Belladonna*, if the pain shoots down into the teeth, neck, and left ear; worse at night, with perspiration, which does not relieve. The pain is better by pressing the head with both hands.

*Chamomile.* Suits often for children and *persons unable to bear the least pain*, and when there are acute, shooting pains in the temples, or a rending, drawing pain on one side extending to the jaw, especially when one cheek is red, the other pale, the eyes painful, sore throat, and bitter taste.

*Colocynth.* Violent, excruciating, tearing, drawing pains on one side, or cramp-like, compressive sensation, aching, with nausea; worse when stooping or lying on the back, with great restlessness; perspiration smells like urine; copious watery urine during the pains, or scanty, offensive discharges of urine between the paroxysms.

*Arsenic.* Pains in the head, sometimes very severe, almost maddening, with nausea, buzzing in the ears, weeping and moaning, tenderness of the scalp; cold applications relieve the pain for the time; suits frequently after *Pulsat.*; when better in open air, worse within doors.

*Veratrum.* Maddening pains one side with great weakness, fainting when rising, cold perspiration, chilliness, diarrhea, tenderness of the scalp; suits well after or before *Arsenic.* If these two remedies will not relieve, give

*China.* Suitable for persons sensitive to pain, feverish, dissatisfied dispositions, talkative and restless at night; the pains are aching, preventing sleep, or are piercing, jerking in the forehead, with a bruised feeling in the brain; worse by *touch*, reflection, conversation, open air, motion, draughts of air; the *scalp* very sensitive to the *touch*.

*Sepia.* (See "Sick Headache.") When the pain in the

temples renders them very sensitive to the touch ; quietness and darkness relieve ; also, a good sleep.

*Sulphur* and *Silicea*. For the chronic treatment, as directed in "Sick Headache," page 135.

ADMINISTRATION. — The above mentioned remedies must be given according to directions in "Sick Headache."

#### FALLING OFF OF THE HAIR. (*Allopecia*.)

This frequently happens after severe fevers, when *Hepar sulph.*, *Silicea*, and *Lycopodium* suit ; or, after *debilitating losses*, such as depletion, excesses, profuse sweats, etc., when we may give *China* and *Mercury* ; or, if it happens to *women when nursing*, *Calcareo carb.* and *Sulphur*. If *long continued grief* is the cause of it, give *Phosphoric acid* and *Staphysag.* If caused by frequent attacks of nervous headache, give *Hepar sulph.* and *Nitric acid*. If caused by having taken too much calomel, give *Hepar* and *Carbo veg.*

ADMINISTRATION. — As it takes a long time to see the effects of a remedy in this disease, do not change it quickly for another, but take every week one dose (six glob.) of a remedy for four weeks, and then discontinue two weeks before taking another medicine. If by that time the loss of the hair seems less, do not change the medicine.

*Externally*, use the same remedy by washing with a solution of twelve globules in half a teacupful of water every other night.

If the *hair is dry* and *splits easily*, cut it often, every month or six weeks a little ; use nothing but pure bear's oil ; all other preparations and pomatums are of little use, and often very injurious.

For the *dandruff*, *scales of the scalp*, take *Calcareo carb.*, *Graphites*, and *Staphysag.* ; if with *itching*, take *Graphites*, *Lycopodium*, and *Sulphur*.

Administer same as above.

## CHAPTER VII.

### AFFECTIONS OF THE EYES.

#### GENERAL REMARKS.

The diseases of the eye ought to claim our most careful attention, as this most important organ is, from its position and complicated and delicate structure, subject to a great variety of diseases, which, heretofore, have been very badly treated under the allopathic system.

All external applications, in the form of strong eye-waters and salves, are injurious, and even if, as in scrofulous sore eyes for instance, these salves seem to have effected a cure, it is only by driving the disease from the external parts to some internal organ, which is, at the time, the weakest—in most cases to the lungs. If external remedies are wanted, pure water is preferable in all cases of inflammation, applied either cold or warm, according to the feelings of the patient. In cases of external injuries to the eye, the arnica lotion (ten drops of *Arnica* tincture to half a teacupful of cold water) must be used externally.

But, when erysipelas affects the eyes, nothing wet ought to be applied, but warm bags filled with bran. (See “Erysipelas.”)

When the inflammation of the eyes and face is caused by the poison of the sumach, or other weeds, apply nothing outwardly, but give the internal remedies recommended for that disease. (See “Poisoning by Sumach,” page 44.)



## 1. INFLAMMATION OF THE EYELID AND ITS MARGINS.

*a. Inflammation and Swelling of the Eyelid.*

*Aconite.* In all cases where the inflammation and the pain is great, attended with fever and restlessness, or where the lids are *swollen, hard, and red*, with *heat* and sensation of *burning* and *dryness*, or when there is a copious secretion of mucous in the eyes and nose, with dread of light, fever, and thirst.

ADMINISTRATION.—Every two hours a dose (four glob.) until three doses are taken or the patient relieved; after it *Belladonna*, *Sulphur*, or *Hepar Sulph.*, are often necessary. (See these.)

*Belladonna.* If *Aconite* is not sufficient; or when there is a feeling of burning and itching in the red and swollen eyelids (which stick together, and bleed easily when opened); also, when the edges are turned to the outside, and feel as if paralyzed (see *Hepar sulph.*)

Administer same as *Aconite*.

*Hepar Sulphuris.* If there is a sensation of *ulceration* or *contusion* in the *red* and *swollen* lids on *touching* them, with sensitiveness to light, and spasmodic contractions of the lids in the morning. In alternation with *Aconite*, *Bellad.*, or *Mercury*, as the symptoms indicate (see these remedies), every three or six hours a dose (four globules), until better.

*Sulphur.* Frequently in alternation with *Aconite*, when the swelling and redness are very bad, with pressing, burning pains in the eye and lids (as if from sand in the eyes), great secretion of mucous, and sensitiveness to light.

ADMINISTRATION.—Alone, or in alternation with *Aconite*, a dose (six globules) every six hours, until three doses of each are taken, or the patient feels better. After *Sulphur*, *Calcareo carb.* is frequently necessary.

*Mercury.* If there is a hard swelling, with cutting pain, and difficulty of opening the lids, as if from contraction, ulcers and scabs on the edges, worse in the night, and in the warmth of the bed, with restlessness and perspiration, or if there is a *burning itching* in the eyes, worse in the open air, with great dread of the light of the fire. After, or in alternation with it, give *Hepar sulph.*, a dose (four glob.) every six hours, until better.

*Euphrasia.* In cases where the eyelids are red and swelled, their margins ulcerated, with itching in the day-time, sticking together in the morning, constant catarrh, headache, heat in the head and dread of light, with *profuse secretion of mucous and tears*. In alternation with *Nux vom.*, if the eyeball is very red, and there is burning with the itching—if the patient is very irritable, and feels *worse* in the open air; but in alternation with *Pulsatilla*, if he is of a mild character, and feels *better* in the open air.

Administration the same as *Sulph.*

*Arsenic.* Inflammation of the inside of the lids, with inability to open the eyes, and *violent burning pain*.

Administer same as above.

*Spigelia.* In inflammation and *ulcerations* of the lids with biting soreness, difficulty of raising the lids (as if from stiffness), with burning in the eyes and pain in the head, worse on the least motion.

Administer same as above.

*Calcarea carb.* Frequently after *Sulph.*, when there are *cutting*, burning pains, especially when *reading*, with red, hard, and great swelling, and copious secretion of mucous, and the lids sticking together in the morning.

Administer same as *Sulphur*.

*Hyosciamus* and *Chamomile* in alternation (administer same as *Bellad.*), when the eyelids are closed spasmodically, with a sensation of heaviness or pressure.

*b. Inflammation of the Margins of the Lids.*

The principal remedies in this disease, which frequently appears without affecting the whole lid, are *Bellad.*, *Euphrasia*, *Pulsat.*, *Nux vom.*, *Mercury*, *Hepar sulph.*, and *Chamomile*.

(See the detailed symptoms of these remedies in the preceding article: "a. Inflammation of the Eyelids.")

2. INFLAMMATION OF THE EYEBALL. (*Ophthalmia*.)

In most cases, where the eyeballs are inflamed, the lids will also be affected, and the remedies stated under "Inflammation of the Eyelids," will be beneficial. This is particularly the case, when *Aconite*, *Bellad.*, and *Euphrasia* are indicated; which are the most important remedies in diseases of the eyes, and to which we refer the reader, in the preceding article. But frequently the eyeballs alone are implicated, especially when cold, together with a scrofulous or rheumatic predisposition, is the cause.

The first remedy in this disease, particularly when it appears suddenly, is *Aconite*. When the eyes are *very red*, with dark redness of the vessels, and intolerable burning, stinging, or aching pains (worse on moving the eyes), dread of light, copious flow of tears, yet with dryness of the eyelids and fever.

ADMINISTRATION. — Same as *Aconite* in "Inflammation of the Eyelids." After which, *Bellad.* is generally sufficient to cure the disease.

*Belladonna*. When the white of the eye is very much inflamed, sensitive to light, either copious flow of burning-hot tears, or great dryness of the eyes, which ache all around, or deep inside, with stitches proceeding from the eyes to the head (worse on moving the eyes), and a severe cold in the head, or a *violent headache*, with stupor and

vertigo, sparks or black spots before the eyes, and a short, dry cough, it suits mostly after *Aconite*; but frequently, too, after *Hepar sulph.*, and *Mercury*.

*Euphrasia*. Same as *Bellad.*, and when this is insufficient; or when the pain in and above the eyes, in the head, is more *pressing*, and the cough not dry, but loose; also, with profuse running at the nose; worse in the evening.

*Nux vomica*. When the eyes are bloodshot, smarting as if from sand, with swelling and softening of the eye, irritable temper, coated tongue, redness of the corners of the lids, with stiffness and itching, acrid tears, dread of light in the morning, headache every night, and stoppage of the nose.

*Ignatia*. If the pain is more like pressure in the eyes, which are not much inflamed, but with copious flow of tears and nasal discharge, great sensitiveness to light, mild, quiet disposition.

*Pulsatilla* is often effective, after *Aconite* has subdued the worst inflammation, and severe pains remain (tearing, stitching, and cutting), in *rheumatic* inflammation of the eyes, where there is sensitiveness to light; worse in the afternoon and evening; the patient is fretful and inclined to weep.

*Bryonia*. After *Pulsat.* when the redness still remains, the lids are swollen, with pain in the head when the eyes are opened; worse at night.

*Rhus*. After *Bryonia*, or in alternation with it, in damp, wet weather, or when the eye continues to burn and stitch, with the running of water, or the eyelids are swollen as if from erysipelas.

*Spigelia*. Aching, stitching, or boring pains, penetrating into the head, and returning regularly at the same time of day, with a sensation as if the eyeballs were too large. The pain is at times insupportable.

Administration of the above remedies, the same as *Aconite*.

*Sulphur* (see "Inflammation of the Eyelid") is also necessary, after *Mercury* and *Pulsatilla*.

DIET AND REGIMEN, in all diseases of the eye must be light, as in fevers generally (see these.) Rest and quiet in the room are necessary. The room itself should be dry, airy, and darkened.

### 3. CHRONIC INFLAMMATION OF THE EYES.

The treatment of these inflammations varies according to their causes, which are either hereditary or acquired.

#### *a. From Gout and Rheumatism.*

For the severest symptoms use *Aconite*, *Bellad.*, and *Spigelia*, as stated in the preceding article; also,

*Colocynthis*. When there is violent burning and cutting, extending far back into the head and nose, with great anguish and restlessness; or violent pressure in the forehead; worse when stooping, or lying on the back; severe drawing pain in the head, with vomiting.

*Hepar sulph.* If *Bellad.*, or *Spigelia*, is insufficient; or when there is redness of the eyes and eyelids, with soreness *when touched*; sensitiveness to light, especially in the evening; sight at times dim and obscured, at others clear; pressure in the eyeball, as if it would *start out of the head*.

Administration of the above remedies, the same as *Aconite* under "Inflammation of the Eyelids."

#### *b. Scrofula, or Scrofulous Sore Eyes.*

Scrofula frequently affects the eyes, after an inflammation of the lids or eyeball has set in, from causes above stated, by which complication such cases are rendered

much worse, and more difficult to cure; for dimness of sight and ulcers on the ball of the eye often remain; they require medical attention, as the consequences of neglect, or faulty treatment, would be too serious; and the necessity for the eradication of scrofula is, of itself, obvious.

In the beginning of such an attack, give *Aconite* and *Belladonna*, as stated under "Inflammation of the Eyelids" and "Eyeball." Also, all the other remedies mentioned there, if they are required by the symptoms. Besides these,

*Dulcamara* is of use, when cold is the cause of the sore eyes, and when they are made worse by reading, with dimness of sight, and a sensation as if sparks of fire were flying out of the eyes; pain over the eyes; worse when at rest; better when walking.

Administration the same as above.

For ulcers on the eyeballs, remaining after the inflammation has subsided, give *Euphrasia*, *Hepar sulph.*, and *Silicea*, each remedy during one week; two or three times every evening a dose (four glob.) When *Silicea* is given, apply externally a wash of twelve globules of *Silicea* in half a teacupful of pure cold water. Make the application several times a day.

For specks, give *Euphrasia*, *Hepar sulph.*, *Silicea*, *Calcare carb.*, *Nitric acid*, in the same manner as stated under "Ulcers on the Eyeball."

DIET AND REGIMEN.—See "Inflammation of the Eyeball."

*c. From Syphilis, or Venereal Disease.*

The eyes may be implicated in this disease, and become inflamed, particularly when a gonorrhea is suppressed and transferred, as it were, to the eyes, or when syphilis has penetrated the whole system. In the latter case, a homœ-



opathic physician must be consulted. In inflammation of the eyes and lids from *suppressed gonorrhea*, administer at first *Aconite*, as stated under "Inflammation of the Eyes;" after it *Pulsatilla*, in the same manner; then wait two or three days, and give, if necessary, *Nitric acid*, *Mercury*, and *Sulphur*; commencing with *Nitric acid*, and taking four doses the first week; four doses of the next remedy, the second week; and so on, until better.

*In the real Syphilitic Eye-affections: Nitric acid, Mercury, and Thuja*, are the principal remedies—to be given as above.

DIET AND REGIMEN.—See "Inflammation of the Eye-ball," and "Syphilis."

*d. From abuse of Mercury.*

This drug, given in allopathic doses, frequently produces severe affections of the eye; for which must be given, *Hepar sulph.*, *Nitric acid*, *Pulsat.*, and *Sulphur*, in the same manner as stated above, under "Syphilitic Eye-Affections." Diet, also, the same.

4. STYE ON THE EYELID.

A styte is a small boil on the eyelid, produced by the swelling of the Meibomian glands, which are imbedded all along the margins of the lid.

The specific remedy in the commencement of this complaint is

*Pulsatilla*. A dose (four glob.) twice a day, until four doses are taken; then discontinue two days; and if not better, or if suppuration appears, take *Mercury* in the same manner, in alternation with *Hepar sulph.*

If no suppuration, but induration takes place, or hard spots remain, and if the styes appear often, give *Staphysag.*, a dose (four glob.) two or three times, during one week;

and afterward, for two or three weeks, give each week two doses of *Calcarea carb.* (each dose four glob.) until better; and, if not, *Sulphur*, for the same length of time, and in the same manner.

Externally, apply nothing; but bathe occasionally in warm water.

DIET AND REGIMEN. — The diet light; no animal food; and regimen cool.

#### 5. WEEPING OR WATERY EYES.

If this complaint arises, as is mostly the case, from an obstruction of the lachrymal duct, or a swelling of its entrance, in the internal corner of the eye, a homœopathic physician should be consulted. In the meantime, use the following remedies: *Bellad.*, *Pulsat.*, *Calcarea carb.*, *Silicea*, *Petroleum*, *Sulphur*; the first two remedies alternately, every day a dose (four glob.) of each, until the acute symptoms subside; the other remedies, each for four or six weeks, every week two doses (six glob. each), until better.

But if the watery eyes arise from a general weakness of the organ, *Euphrasia* and *Spigelia* in alternation, every two or three days a dose (six glob.), will generally be found beneficial. Let this treatment be followed, at the end of two weeks, by two doses of *Sulphur*, on two successive nights (each dose, six glob.)

DIET. — The usual homœopathic diet must be observed during the whole time.

#### 6. CATARACT. (*Glaucoma*.)

This disease, which consists of the gradual darkening of the lens in the eye (thus preventing the light from reaching the optic nerve) is, in its commencement, curable by medicines, if carefully selected and applied. As this, however, ought to be done by a homœopathic physician, we will

here merely notice a few remedies, which may be taken in the beginning, until a physician can be consulted.

By this means, an operation for the cataract may frequently be avoided.

*Pulsatilla, Sulphur, Silicea, Conium, Phosphorus.*

ADMINISTRATION. — Give each remedy for four or six weeks, every three days one dose (six glob.), in the evening, until better, or until the next remedy has to be taken.

DIET AND REGIMEN. — Light, without animal food; otherwise, the usual homœopathic diet.

## 7. WEAKNESS OF SIGHT. (*Amblyopia.*)

*Blindness (sudden and incipient) — Amaurosis.*

Short-sighted and far-sighted persons have not, strictly speaking, *weak* eyes; this latter is a defect in the optic nerve itself, which is weakened or paralyzed, in a greater or less degree. If this nerve is not much paralyzed, so that only dimness of sight exists, we call it *amblyopia*, or *weakness of sight*; if the nerve is more or entirely paralyzed, it is incipient or real blindness—*amaurosis*.

This disease can be cured; but it must be taken in time, and a homœopathic physician ought to be consulted *immediately*, as delay in this case is very dangerous.

In the meantime take, for *simple weakness of sight*, *Pulsatilla, Sulphur, Bellad., Calcarea carb., and Phosphorus.*

Administration the same as stated under “Cataract.”

For *incipient blindness*, give the same remedies as for simple weakness of sight.

*Attacks of sudden blindness*; if caused by congestion to the head, and accompanied with pain in the head and convulsions, give *Bellad.* and *Hyosciamus* in solution (twelve glob. in half a teacupful of water), every fifteen minutes a teaspoonful, until better (see “Congestion of blood to the

Head"); but for sudden blindness, without an apparent cause, give first *Aconite*; then *Mercury*; every two or three hours a dose (four glob.) alternately, until better.

For *night-blindness*; *Belladonna*.

For *day-blindness*, where the patient can only see in the evening; give *Sulphur*, and afterward *Silicea*.

DIET the same as in "Cataract."

#### 8. SHORT SIGHTEDNESS. (*Myopia*.)

This affliction, if not of too long duration, may generally be mitigated or cured by *Pulsatilla*; or, if caused by abuse of calomel, give *Carbo veg.* and *Hepar sulph.*; if from debilitating causes and typhus fevers, *Phosphoric acid*.

These remedies to be administered in the same manner as stated under "Cataract."

DIET AND REGIMEN, also, the same.

#### 9. FAR SIGHTEDNESS.

This frequently occurs after excessive use of ardent spirits; in which case *Nux vom.* must be given; every day a dose (four glob.) for a week; followed by *Sulphur*, in alternation with *Lachesis*; every week two doses of each, for four or six weeks, or until better. If not relieved, consult a physician.

#### 10. SQUINTING. (*Strabismus*.)

A frequent cause of squinting, in young children, is that from the position of their cradle, they constantly have the light on the same side. To obviate this, alter the position of the cradle so that the light will be directly in front.

If, however, squinting continues, notwithstanding, and the heads of the children are hot, give *Belladonna* and *Bryonia*, in alternation, every four or six days a dose (two

glob.) until six doses of each are taken, or the child is better. If not better, consult a homœopathic physician.

### 11. FALLING OF THE EYELIDS. (*Paralysis.*)

If the upper or lower eyelid falls down, and can not be raised easily, give *Spigelia*, *Belladonna*, *Sepia*, *Veratrum*, *Opium*, *Cocculus*, *Nitric acid*.

Administer the same as stated under "Cataract."

DIET AND REGIMEN.—The usual homœopathic diet—bathe the eyelids frequently in cold water.

### 12. FOREIGN SUBSTANCES IN THE EYE.

If dust enters the eye and causes irritation, washing with cold water will be sufficient; but cream or sour milk will answer better, if lime, ashes, or tobacco enters the eye. If a hard substance, or an insect has got into the eye, draw the eyelids apart, and turn the upper one over the lower one a couple of times, until it is felt that the substance is removed. If unable to remove it in this manner, do not rub the eye, but lie down, and bathe it in a solution of *Arnica* (five drops to half a teacupful of water.) Internally, take *Aconite*, and try to sleep.

If particles of iron have entered the eye, and are fastened in the white or pupil, use *Arnica*, as above stated, and have them extracted by a surgeon.

If the eye becomes very much inflamed, take a few doses of *Aconite*, and afterward *Sulphur*; and, if necessary, in a few days after, *Calcarea carb*.

## CHAPTER VIII.

### AFFECTIONS OF THE EARS.

#### 1. EXTERNAL EAR AND ITS SURROUNDING PARTS.

##### *a. Eruptions and Swelling of the External Ear.*

The eruptions which appear on or behind the ear are various, and generally require a chronic treatment to eradicate the disease entirely, which can be done only by a skillful homœopathic physician. To begin the cure, or if severe, acute sufferings, as swelling and inflammation of the ear, appear with the eruption, select from among the remedies given below, and read their symptoms in the "Materia Medica" at the end of the book.

1. For *scurvy eruption* : *Pulsat.*, *Lycopodium*, *Lachesis*, *Hepar sulph.*, *Staphysag.*, *Graphites*.

2. For *soreness* : *Lachesis*, *Mercury*, *Sulphur*, *Graphites*.

3. For *itching in or on the ears* : *Pulsat.*, *Lycopodium*, *Sulphur*.

4. For *swelling of the ear* : *Pulsat.*, *Mercury*, *Lycopodium*, *Sepia*.

5. For *ulcerated eruptions* : *Pulsat.*, *Mercury*, *Hepar sulph.*

6. For *humors* : *Graphites*, *Lycopodium*, *Calcarea carb.*

ADMINISTRATION. — Give each remedy for one week, every other day one dose (four glob.) until better, or until the next remedy has to be given.

DIET AND REGIMEN. — The usual homœopathic diet, and



particular care to be taken not to drive the eruption away by getting cold or by external applications.

*b. Mumps. (Parotitis.)*

This disease consists in an inflammation of the great salivary glands, lying under and in front of the ear; children are frequently subject to it, but seldom more than once in a lifetime.

It is not dangerous in its simple form, and only requires to be kept warm by putting a handkerchief around it, and keeping the patient in the room. Internally give

*Mercury*, which is almost a specific for this disease — every evening a dose (four glob.), until three or four doses are taken.

Sometimes the glands of the neck participate, and remain swelled for three or four days. In this case, also, pursue the above treatment.

But, if the mumps disappears suddenly, by striking in and falling upon the brain, so that the patient becomes delirious and unconscious, give *Belladonna* in solution (twelve globules in half-a-teacupful of water), every hour a teaspoonful. (See, also, “Scarlet Fever,” striking in of.)

But, if in striking in, it falls upon the *stomach*, give *Carbo veg.*, in the same manner as *Belladonna* above, and, if not better in two or three hours, *Arsenic*, and after that *Coccullus*, if necessary, in the same manner.

If fever, restlessness, and erysipelatous redness are combined with it, particularly in damp weather, give *Belladonna* and *Rhus* in alternation, in solution (twelve globules of each in separate teacups, each half full of water), a teaspoonful every three or four hours, till four of each remedy are given, or until better.

If the mumps is very obstinate and very hard, give *Kali*

*carb.*, every evening one dose (six glob.) until three doses are taken.

If a catarrhalic state is connected with it, particularly hoarseness, give *Carbo veg.* in the same manner. This same remedy suits if the patient has previously taken much calomel.

DIET AND REGIMEN. — The diet must be light and farinaceous — baked apples and stewed prunes, if desired, but neither animal food nor soups. The patient must be kept warm, and if feverish must stay in bed.

## 2. INTERNAL EAR.

### *a. Inflammation of the Ear. ( Otitis.)*

This painful disease is generally caused by exposure to cold and damp weather ; sometimes, however, by the striking in of eruptions.

It manifests itself by violent burning, stitching, beating pains, deep in the ear, and finally by swelling and redness of both in and outside of the ear ; frequently, in very severe cases, the brain is implicated, when, in children, convulsions ensue, with vomiting and cold extremities.

TREATMENT. — The principal remedy is, first,

*Pulsatilla*. In solution (twelve globules in half-a-teacupful of water), every hour a teaspoonful until better, or until six teaspoonfuls are taken ; then wait the same length of time and give one dose (four glob.) of *Sulphur*, after which, in two hours, repeat the *Pulsat.* until better. But, if the brain becomes affected, with or without convulsions, give

*Belladonna*. In alternation with *Aconite*, if the fever is very high, every two or three hours a dose (four glob) ; then wait four or six hours and give *Belladonna* <sup>cc.</sup> and *Coffea* <sup>cc.</sup>, in the same manner, if great sensitiveness and

restlessness remain. If an abscess is about to form, which is indicated by the continued beating pain in the ear, give *Mercury*, *Lachesis* and *Hepar sulph.*, in alternation, every six hours a dose (four glob.) until better, or until two or three doses of each are taken.

DIET AND REGIMEN. — The diet similar to that in fever. If the pain is very violent, make an external application of heated flannels, or bags filled with heated bran or scorched flour.

*b. Earache. (Otalgia.)*

Earache, commonly so termed, resembles the inflammation of the ear in the character of the pain, and is only different in regard to its origin, as it mostly appears after taking cold as a neuralgic and rheumatic affection, or in consequence of previous inflammations of the ear. There is little or no fever accompanying the attacks, which are generally of short duration, and relieved speedily by the application of the following remedies :

TREATMENT. — *Pulsatilla* is the first and principal remedy, particularly if the pains are darting and tearing, as if something would *press out through the ears*; the outer ear is *red, hot, and swollen*, or, there is stitching and tearing through the whole side of the face, almost depriving the patient of reason, particularly in females, or in persons inclined to chilliness and disposed to cry.

Administer as *Pulsat.* in "Inflammation of the Ear."

*Chamomile.* The next best remedy, especially after taking cold, or when a perspiration has been suppressed suddenly; tearing, lancinating pains in the ear, which extend to the lobe of the ear; the patient is very cross, cannot bear pain, and is easily offended.

*Mercury.* After *Pulsat.* or *Chamomile.* If these are insufficient, or if the patient perspires a great deal without

relief; *the pains are worse in a warm bed*, and there is a *chilly sensation in the ears*, with moisture; the pains extend to the cheek and teeth.

*Nux vomica*. After *Chamomile*, in persons of a lively, passionate disposition; when the pains are tearing and stinging, tending upward to the forehead and temples, and down into the facial bones; worse in the morning.

*Arnica* is more for sensitive, nervous persons, when the pain returns on the slightest occasion, with sensitiveness to noise, and pressing and stinging in and behind the ears, which feel hot internally.

*China* suits frequently after *Arnica* or *Pulsat.*; if the tearing pains are felt more externally; are worse by *touching the ear*, which is red, stings, and rings.

*Belladonna*. Similar to *China*, or as stated under *Belladonna* in "Inflammation of the Ear."

*Hepar*. After *Belladonna*, when the pain shoots into the ear on blowing the nose, or when the pain is beating, throbbing, or roaring.

*Spigelia*. When there is a painful aching, as if a large nail were sticking in the ear; with aching and tearing in the cheek bones.

*Dulcamara*. When the pains increase during rest, especially at night, with nausea.

ADMINISTRATION of the above remedies the same as *Pulsat.*, in the beginning of this article.

If the above remedies allay the worst symptoms, but there is still left a painfulness in the ear, particularly on the left side, or the pains increase in the evening or before midnight, give *Sulphur*, evening and morning, a dose (four glob.); but if worse on the *right side*, and after midnight, toward morning, or during the forenoon, give *Calcare carb.*, the same as *Sulphur*.

In very small children, after *Pulsat.* and *Chamomile* (two

glob. each) have been given at intervals of one or two hours, give *Coffea* and *Belladonna* in the same manner, and afterwards *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, every hour alternately, a dose (two glob.) until better.

DIET AND REGIMEN. — The diet light; no meat, but fruit and farinaceous substances. Externally use nothing but what is recommended in “Inflammation of the Ear.”

*c. Humming, Buzzing of the Ears.*

As a symptom it invariably accompanies earache, running of the ears and their inflammation, and will disappear with them; but, when it appears alone, it is caused generally by congestion to the head, from catching cold, or other diseases.

*Pulsatilla* relieves it, if worse in the evening.

*Nux vomica*, if worse in the morning.

*Dulcamara*, if worse at night.

*Mercury*, in persons inclined to perspire.

*Chamomile*, in persons who do not perspire easily.

*China*, in persons who have taken much calomel, or where the noise in the ear is more a hissing, singing, or ringing noise.

*Carbo veg.*, if *China* does not relieve, or the noise is worse in damp weather.

*Sulphur*, if *Carbo veg.* does not relieve, or the noise is combined with great sensitiveness to sounds, in which case it ought to alternate with *Aconite*.

ADMINISTRATION. — Either of the above remedies, if selected, must be given, one dose (six glob.), on two successive evenings, and then discontinued three days before selecting another remedy, if no improvement has taken place.

DIET. — The usual homœopathic diet.

*d. Hardness of Hearing. Deafness.*

This complaint is very frequent, and as it is often combined with other chronic diseases of the ear, it ought to be treated by a skillful homœopathic physician. When it is of a recent date it may be cured easily, but not so readily when of long standing. Sometimes it is caused by the accumulation of earwax, in which case have it removed by skillful hands. If, after such an operation, great painfulness remains in the ear, put a few drops of *Arnica* solution (two drops of arnica tincture to three tablespoonfuls of water, well shaken), into the ear and lie down.

If the earwax is very hard to remove, put, before its removal, a few drops of warm milk or sweet oil into the ear.

If there is a want of ear-wax, or too great dryness of the ear, give *Carbo veg.*, particularly after a suppression of fever and ague by quinine, every other night one dose (four glob.) three times; then wait eight days, and, if no better, give *Graphites*, particularly after suppression of an habitual eruption, such as tetter, or where there is a *humming* or *thundering* in the ears, especially at night; give it in the same manner as *Carbo veg.* If no better after *Graphites*, try *Lachesis*, *Nitric acid*, and *Petroleum*, one after the other, in the same manner.

If there is difficulty of hearing, with running of the ears, select among the remedies prescribed for this complaint, especially *Pulsat.*, *Mercury*, *Sulphur*, and *Calcarea carb.*, to be given in the same manner as above.

If it is in consequence of congestion to the head, give, in preference, *Belladonna*, *Coffea*, *Nux vomica*, *Sulphur*, *Silicea*, commencing with the first, evening and morning one dose (four glob.), and wait two days before the next one is given.



If it follows after acute eruptions, such as measles, etc., give,

If after *measles*, *Pulsat.* and *Carbo veg.*;

If after *scarlet fever*, *Belladonna* and *Hepar sulph.*;

If after *small pox*, *Mercury* and *Sulphur* — every other evening a dose (six glob.) alternately, until better.

If it is caused by a *cold in the head*, or a *chill* of the whole body, give the remedies for these complaints (see these articles), particularly, however, *Chamomile*, *Arsenic*, *Lachesis*, *Mercury*, *Sulphur*, and *Pulsat.*, in the same manner as there prescribed.

Sometimes hardness of hearing ensues after intermittent fevers have been suppressed by the use of quinine; in this case give *Carbo veg.*, *Pulsat.*, *Sulphur*, and *Calcarea carb.*, every week one remedy, on two successive evenings, one dose (six glob.) until better.

If caused by the suppression of chronic eruptions, or ulcers healed by salves, give *Sulphur* or *Antimon. crud.* (See “Eruptions.”)

If it is from *swelled tonsils*, give *Mercury*, *Staphysag.*, *Nitric acid*, and *Aurum*, every week one remedy, on two successive evenings one dose (six glob.) until better.

If caused by the suppression of a discharge from the ears or nose, give *Pulsat.*, *Mercury*, *Lachesis*, *Belladonna*, and *Hepar sulph.*, every two or three days one remedy, (once a day a dose, four glob.), until better.

N. B. Do not use any external means, such as oils, however much they may be recommended.

### *e. Running of the Ears. (Otorrhœa.)*

A running of the ears frequently remains after the inflammation has subsided, and is then beneficial in diminishing local tension. In such a case, continue the use of the remedies for “Inflammation of the Ear;” after which it

will generally soon disappear. The principal remedies are *Pulsatilla*, *Mercury*, and *Sulphur*.

If it succeeds the measles, give *Pulsatilla* and *Sulphur*, for four days; each remedy on two successive evenings (six glob. a dose.)

If after scarlet fever, give *Belladonna*, *Mercury*, and *Hepar sulph.*, in the same manner.

If after small-pox, give *Mercury*, *Lachesis*, *Sulphur*, *Calcaria carb.*; each remedy for six or eight days (on two successive evenings one dose; six glob.) until better.

In regard to the quality of the discharged substance, give, if it is

*Purulent matter* : *Pulsat.*, *Mercury*, *Hepar sulph.*, *Calcaria carb.*

*Bloody matter* : *Pulsat.*, *Mercury*, *Lachesis*.

*Mucous liquid* : *Pulsat.*, *Mercury*, *Sulphur*, *Bellad.*, *Calcaria carb.*

*Very offensive matter* : *Mercury*, *Hepar sulph.*, *Lycopodium*, *Pulsat.*, *Sulphur*.

Each of the above remedies, when indicated, for six or eight days (on two successive evenings one dose—six glob.) until better.

The sudden suppression of a running of the ear may produce dangerous consequences, which must be attended to immediately.

If, after such a suppression, the glands of the neck, or in front of the ear, begin to swell or get hard, give *Pulsat.*, *Mercury*, *Bellad.*; if severe headache or fever ensues, give *Bellad.*; if not better in a few hours, *Bryonia*. If the suppression is caused by getting cold in damp weather, give *Dulcamara*; and if not better in three or four hours, *Rhus* and *Bellad.*, in alternation.

If, after a suppression, the testicles begin to swell, give *Pulsat.*, *Arnica*, *Mercury*, *Nux vom.*

ADMINISTRATION of the above remedies, give every two or three hours a dose (four glob.), alternately, until better. If by these directions no relief is obtained, consult a homœopathic physician.

DIET AND REGIMEN. — The same as in "Earaehe."

*f. Polypus of the Ear.*

These exeresenees appear frequently, during a long continued running of the ear; particularly if accidental injuries by ear-picks, etc., have taken place. They either have a small or broad basis, whence they grow to a size, so as to fill up the whole internal ear. They are not painful, but always discharge an offensive mucous, and bleed easily when touched.

They are not so dangerous as they are annoying and obstinate; and require but a proper homœopathic treatment, by a skillful physician. In no case is it advisable to extricate them by force; we have remedies which will remove them. The principal medicine is *Calcarea carb.*, of which every second day, in the evening, a dose (six glob.) may be given, until medical aid can be obtained.

If the polypus is very large, touch it twice a day with a solution of the gum of *Opium*, conveyed into the ear with a sponge. Otherwise, keep the ear clean by frequent injections of milk and water.

*g. Foreign Substances in the Ear.*

Insects sometimes get into the ear; in such cases, drop sweet oil into the ear. This kills the insect, and enables us to take it out easily.

If a bean, or any thing that will swell, gets into the ear, bend the point of a large and stout sewing needle a very little, so as to form a fine hook; enter the ear with it cautiously, without touching the walls of the ear, and press it

into the side of the swollen substance, which is thus caught by the hook, and may be slowly extracted. After such an operation, it is necessary to drop into the ear a few drops of an *Arnica* solution (two drops of the *Arnica* tincture to two tablespoonfuls of water, and well shaken), and internally, *Arnica* and *Pulsat.*, in alternation, every two or three hours a dose (four glob.) If there is great fever and delirium, give *Bellad.*, and *Aconite*, in the same manner.

If painfulness still remains, give, in a couple of days afterward, *Sulphur*, evening and morning a dose (four glob.)

## CHAPTER IX.

### AFFECTIONS OF THE NOSE.

#### 1. INFLAMMATION OR SWELLING OF THE NOSE.

The nose, from its position and function, is exposed to a great many diseases and injuries.

*a. External injuries*, which may befall the nose, from the slightest contusion to the severest blow or cut, require *Arnica* internally, every four or five hours a dose (four glob.), and externally, the same remedy in solution (twelve drops of the tincture to a teacupful of water, well mixed.)

*Arnica* will also stop the bleeding caused by severe blows and other injuries of the nose.

*b. Internal causes* for inflammation and swelling of the nose are *scrofula*, the abuse of *ardent spirits* and *calomel*, or the *congestion* attending a cold.

In cases of *scrofulous* or *erysipalalous* inflammation of the nose, give

*Belladonna*. If the nostrils are swollen and sore ; with heat, redness, and pain, outwardly and inwardly, attended with a burning, pricking, and dryness ; smelling sensitive, or absent ; pains as if bruised, on touching the nose ; give, evening and morning a dose (four glob.) ; and if no better, after two days, give

*Hepar sulph.*, in the same manner, until better ; followed by *Mercury*, in the same manner ; and if this does not re-

lieve, give *Bryonia*, followed by *Rhus*, in the same manner; particularly the latter, in cases of poisoning by weeds.

When caused by excessive use of *ardent spirits*, give *Lachesis*, *Arsenic*, *Pulsat.*, *Sulphur*, *Mercury*, *Bellad.*, and *Calcarea carb.*, one after the other, in the same manner, until better.

If caused by abuse of calomel, give *Aurum*, *Bellad.*, *Lachesis*, *Hepar*, in the same manner, until better.

If caused by *cold*, see "Cold in the Head."

*Black spots* frequently appear on the outside of the nose, for which *Sulphur*, or else *Graphites*, may be taken, on two successive evenings one dose (four glob.)

For *red spots*: *Phosphoric acid*, in the same manner.

When the end of the nose is red, give *Rhus* and *Calcarea carb.*, in the same manner.

If the redness on the end of the nose is *copper-color*, give *Arsenic* and *Veratrum*, in the same manner.

When *warts* appear on the nose, give *Caustic.*, every week two doses (four glob.), and wash with the same externally, until better.

DIET AND REGIMEN must be strictly homœopathie, as laid down in the introduction.

## 2. ERUPTIONS ON THE NOSE. — ITCHING OF THE NOSE.

### a. *Eruptions on the Nose.*

The eruptions on the nose are generally of a herpetic or tettery character, and require, for their radical and safe extinction, a long continued, careful, homœopathie treatment. The use of allopathie remedies, in the form of salves and ointments, is, in the highest degree, dangerous, and ought to be at once rejected; as they may cause the disease to disappear externally, and fall on the eyes or lungs. It is much better to allow the eruption to stay



where it is (as it can do no harm there), than to endanger, with its repression, the life of the patient. To commence its radical cure, however, give

*Rhus*, if the redness is on the point of the nose, with soreness to touch.

*Antimonium*, if it is all over and around the nose, with soreness in the corners.

*Causticum*, if it is on the point of the nose.

*Nitric acid*, if the eruption is tettery, with stinging in the nose, as if from splinters on touch; itching sensation; scabby eruption.

ADMINISTRATION. — These remedies to be given, every three or four days a dose (six glob.), until better, or another one is indicated.

#### *b. Itching of the Nose*

Is merely a symptom of other diseases in the system; particularly in children, where it indicates an irritation near or below the navel, either from worms, inflammation of the bowels, or swelling of the glands in the abdomen.

For its treatment, see these diseases.

### 3. BLEEDING OF THE NOSE. (*Epistaxis*.)

Some persons are subject to this complaint constitutionally, sometimes relieving, often injuring them very much. For weak lungs, with tendency to bleeding at the nose, apply immediately to a homœopathic physician. Meanwhile, take *Aconite* and *Sulphur* alternately, every three or four days a dose (six glob.), until better, or until four doses of each are taken; afterward, if necessary, *Sepia* and *Carbo veg.*, in the same manner. If the bleeding appears in females periodically, every month, and the menses have ceased, restore the latter by the proper treatment (see

“Diseases of Females”), and the former will cease of itself.

If it appears in the beginning of a typhoid or other fever, or irregularly during its period, it is not considered a very favorable sign; but if it appears on the critical days of such a fever (say the third, fifth, seventh, ninth, thirteenth, eighteenth, or twenty-first day), it is thought to be favorable, and ought not to be interfered with immediately (see under “Fevers”), except it becomes too profuse, when *China* will give relief.

If it results from determination of blood to the head, give *Aconite* and *Bellad.*, as directed in “Congestion to the Head.”

If it happens after a blow or cut of the nose, apply *Arnica*, externally and internally, as stated under “Inflammation of the Nose.”

If it arises from the use of spiritous liquors, give *Nux vom.*, every two or three hours a dose (four glob.)

If it is from overheating, lie quietly, with the head elevated, and take *Aconite* and *Bryonia* alternately, every hour or thirty minutes a teaspoonful, in solution (twelve glob. of each in separate teacups half full of water.)

If it is in consequence of great exertion, give *Rhus*, and *Carbo veg.*, in the same manner, until better.

If it occurs mostly during the *night*, give *Rhus*, *Bryonia*, and *Bellad.*, alternately, every evening a dose (six glob.)

If in the *morning*: *Nux vom.*, *Bryonia*, *Carbo veg.*, in the same manner.

If it occurs on the most trivial occasion, after eating, etc., give *Silicea*, *Sulphur*, *Sepia*, *Calcarea carb.*, *Graphites*, *Lycopodium*, at intervals of from eight to ten days, each remedy for four weeks (six globules for a dose), to eradicate the disposition to it.

Bleeding from the nose in children, from worms, requires *Cina* and *Mercury*, in the same manner. During the attack, and to allay the itching in the nose, give *Cina* <sup>cc.</sup>, every two or three hours a dose (four glob.)

If it occurs during a cold in the head, *Pulsat.* and *Arsenic* will be necessary, alternately, morning and evening a dose (six glob.) until better.

If persons are naturally weak, or rendered so by loss of blood from the nose, give *China* in solution (twelve glob. to half a teacupful of water), every half hour or hour a teaspoonful until better, or afterward *Ferrum*, if necessary, in the same manner.

*Externally* use nothing but cold water or ice, on the root of the nose, keep perfectly quiet, and do not remove the clots of blood from the nose, if they form themselves.

Lint pressed tightly under the upper lip frequently stops the severest bleeding of the nose.

The *diet* must be of a cooling nature ; nothing should be used in the least degree stimulating ; no meat.

#### 4. COLD IN THE HEAD. (*Coryza*.)

*Influenza*, *catarrhal fevers*, and colds on the lungs, generally commence with affecting the mucous membrane of the nose, whence the disease descends into the larynx and bronchia. We will here treat only this diseased condition of the nose, which manifests itself mostly by sneezing, obstruction, or running of the nose, with excoriations, headache, feverishness, and affections of the eyes. If the disease runs into one of the above mentioned diseases, see their respective chapters.

TREATMENT. — In all cases where persons are attacked with a cold in the head, first give *Nux vom.* and *Mercury*, in alternation, every four or six hours a dose (four glob.) for twenty-four hours ; and then, if not better, select one

from among the following remedies, each of which may be given three times in twenty-four hours (four glob. in a dose.) If the patient is better, wait two days without giving any medicine, observing the diet, however; and then give a dose of *Sulphur* (four glob.) on two successive evenings, which will complete the cure.

*Nux vomica* must be given for the following symptoms: either *dryness* and *obstruction of the nose*, with *heaviness in the forehead* and impatient humor; or *running of the nose* in the *morning*, but *dryness in the evening* or *during the night*; with *dryness of the mouth*, and constipation of the bowels; the patient feeling better when exercising or warm.

*Mercury*. *Frequent sneezing*, particularly when coming from the cool air into a warm room, *profuse running* of the nose, with *swelling, redness*, and *excoriation* of the nostrils; itching, and aching pains on pressing the nose and cheek-bones; soreness and pains in the limbs, copious perspiration at night; great thirst; patient feels *worse* in the *warmth*, yet *can not bear the cold*.

*Hepar*. For persons who have already taken calomel, or were not relieved entirely by the homœopathic *Mercury*; or when every exposure to cold air renews the complaint, with a headache; worse by movement; or where the cold attacks only one nostril.

*Euphrasia*. When, with profuse running of the nose, there is redness and soreness of the eyelids, with profuse flow of tears, and confused sensation in the head.

*Belladonna*. If *Hepar* does not relieve, or when the smell is alternately too acute or too dull.

*Lachesis*. Frequent sneezing, with profuse running of the nose, with swelling and excoriation of the nostrils and lips: or else, obstruction of the nose, humming in the ears, ill humor, constipation, and unfitness for meditation. Particularly after *Nux vom*.

*Arsenic.* Obstruction of the nose, alternately with profuse running and excoriation of the nostrils, with bleeding; restlessness and sleeplessness in the night; a desire to drink frequently, but little at a time; feels better in a warm air; suits well after *Nux vomica*.

*Chamomile.* When cold is caused by suppressed perspiration, as is often the case in children, with feverishness and drowsiness; stupor; one cheek pale and cool, the other red and hot; soreness of the nostrils; cracked lips and acrid, smarting mucous in the nose.

*Pulsatilla.* When the patient loses his appetite and smell, and the nose secretes a *yellowish, thick*, offensive mucous, with *heaviness* and confusion in the head; particularly when *toward night and in the warmth of a room* the obstruction of the nose increases; feels better in the open air, yet shivering; disposed to shed tears. It is beneficial before *Lycopodium*, and follows well after *Chamomile*.

*Lycopodium.* Obstruction of the nose, especially at night; with the *sense of smell too acute*; tearing pain in the forehead; *dryness of the mouth, without thirst*; constipation of the bowels, with melancholy, sadness, and whining mood. Follows well after *Pulsatilla* or *Lachesis*.

*Sulphur.* Obstruction and dryness of the nose, or profuse secretion of thick, yellowish, puriform mucous; sneezing; blowing of blood from the nose, with excoriation and soreness of the nostrils; worse in the night and in wet weather.

#### *Suppression of the Catarrh, or Cold in the Head.*

If the running stops suddenly and headache ensues, give *Aconite* a few doses, every two or three hours a dose (four glob.); if not better, give *Pulsat.* and *China*, alternately, every six hours a dose (four glob.) If hard breathing or asthma ensues, give *Ipecac.* and *Bryonia*, every two hours

a dose (four glob.), alternately, until better, or until three doses of each are taken; and if not relieved give *Arsenic* and *Sulphur*, alternately, every four or six hours a dose (four glob.) until better. If the eyes are affected, give *Belladonna*, *Lachesis*, and *Hepar*, in alternation, every six hours a dose (four glob.) until better. If inflammation of the windpipe or lungs ensues, give *Aconite* and *Bryonia*, alternately, in solution (twelve globules in half a teacupful of water), every two hours a teaspoonful.

If a person is subject, on the slightest occasion, to cold in the head, bathe the neck and breast frequently in cold water; snuff cold water up into the nose when washing the face, and take *Silicea*, every eight days a dose (six glob.)

N. B. We would remark here that in the treatment of children, where it is difficult to see the symptoms with sufficient clearness for the selection of a remedy, the alternate use of *Chamomile* and *Belladonna* should be adopted in most cases, giving every three or four hours a dose (two glob.), which removes the cold. If drowsiness sets in and sleepiness, with eyes half open, the use of *Belladonna* and *Hellebor.* is preferable, given in solution, as in "Dropsy of the Brain." After a few days, *Sulphur* is frequently necessary to complete the cure.

DIET AND REGIMEN. — If there is fever, see the appropriate diet in "Fever;" if without fever, the usual homœopathic diet. During a cold in the head, abstain from the usual bathing on feeling worse after it.

##### 5. ULCERATION IN THE NOSE.

(*Catarrh. Ozæna narium.*)

This is a chronic disease, where an ulcer has formed in the upper part of the nose (extending sometimes to the



frontal sinus), from which an offensive matter is discharged, frequently mixed with blood; as long as it discharges freely the accompanying pain is inconsiderable, but when the ulcer suddenly stops running, feverishness ensues with headache, swelling and redness of the nose, thirst, etc. All these symptoms cease as soon as the ulcer begins to run again. If this disease is located in the nose, it is called *ozæna narium*; if in the cavity of the cheek bones, *ozæna antri Highmori*.

As both these diseases are difficult to cure, being caused by scrofulous or syphilitic disorders, it is necessary to consult a homœopathic physician; yet we will point out the remedies with which the cure may be commenced.

*a. Ozæna narium. Catarrh.*

TREATMENT. — The cure may be commenced with *Belladonna* every second evening a dose (six glob.), particularly in females; continue this treatment for one week, followed by *Aurum*, in the same manner, and this followed by *Mercury* if necessary, after which take *Sulphur* in the same manner.

If, however, the ulcer does not discharge, and there is fever and heat, give *Belladonna* and *Aconite*, alternately, in solution (twelve globules in half a teacupful of water), every two hours a teaspoonful.

*b. Ozæna antri Highmori.*

The same remedies as above, but if there are severe pains in the cheek bones, throbbing, splitting, or bursting, give *Arsenic* in solution, every two hours a teaspoonful until better.

If the discharge is thick and yellow, give *Lycopodium*, every other evening a dose (four glob.) until better.

For a *chronic ulceration in the lower part of the nose*, and scabs on the nostrils, take *Mercury*, *Lachesis*, *Calcarea*,

*Lycopodium*, *Graphites*, *Sulphur*, each remedy for two weeks, every five days a dose (six glob.); after taking one remedy in this manner for two weeks, wait one week before the next remedy is commenced. Continue this treatment until better.

DIET AND REGIMEN.—The usual homœopathic diet. Frequent applications of cold water are recommended.

#### 6. POLYPUS OF THE NOSE — CANCER OF THE NOSE.

The *polypus of the nose* is of frequent occurrence, and can be cured by the application of the homœopathic medicines, as, also, that variety of this disease which consists in the thickening or enlargement of the cartilages of the nose. As these disorders, however, require medical advice, we merely mention here such remedies as may commence the treatment of both varieties with equal propriety.

TREATMENT.—*Staphysag.*, *Phosphorus*, *Calcarea*, *Silicea*, *Sepia*, each remedy (every five days a dose, six glob.) for three weeks, and then waiting one before the next remedy is commenced, until better.

*Cancer in the Nose*.—The treatment of this disease ought to be confined to the application of the proper internal remedies, as the cause of it lies in the system. By applying early to a homœopathic physician it is sometimes cured, while under the allopathic treatment a cure is never effected.

The cure may be commenced with *Arsenic* and *Sulphur*, in the same manner as above, under “Polypus of the Nose.”

DIET AND REGIMEN.—As usual in homœopathic treatment of chronic diseases.

#### 7. FOREIGN SUBSTANCES IN THE NOSE.

If a child has put a bean into its nose, let it draw in its breath, then close its mouth, and make it force the breath

out through the nose ; or tickle the nose with a feather to make it sneeze. Sometimes the bean can be pushed back so as to fall into the mouth, or it can be extracted with a needle, the point of which is turned into a small hook. If this does not succeed, apply to a surgeon. After the operation apply *Arnica* in water, as usual, and give internally *Arnica* and *Rhus*, alternately, every six hours a dose (four glob.) In a couple of days give one dose (four glob.) of *Sulphur*.

## CHAPTER X.

### AFFECTIONS OF THE FACE, LIPS, AND JAWS.

#### 1. INFLAMMATION AND SWELLING OF THE FACE.

##### *a. Erysipelas of Face and Lips.*

For the treatment of this disease see “Erysipelas,” under the diseases of the skin. The principal remedies are *Belladonna*, *Lachesis*, and *Rhus*, to be given as there indicated.

##### *b. Swelling of the Cheek,*

Caused by toothache, is generally relieved by *Chamomile*, *Belladonna*, and *Mercury*, in alternation, every three hours a dose (four glob.), when the swelling is *red, hot, and hard*.

*Bryonia* and *Nux vomica*, in alternation, in the same manner, when it is *pale*. If the swelling is very painful, put a bread-and-milk poultice on it. (See, also, “Toothache.”)

DIET AND REGIMEN — As in fevers.

#### 2. ERUPTIONS OF THE FACE.

The face is, in common with the nose, liable to eruptions in the shape of pimples, produced by various causes. (See “Inflammation and Eruptions of the Nose,” and “Inflammation of the Face.”)

For the pimples (*acne rosacea*) in *drunkenness* (commonly called *carbuncles*), give *Nux vomica*, *Lachesis*, *Arsenic*, and

*Sulphur*, every eight days one dose (six glob.) of one remedy, and so on in alternation until better. The abstinence from spiritous liquors during this time and afterward is, of course, required.

For *pimples* in the face, as they appear often in young people, give *Belladonna*, *Hepar*, *Carbo veg.* and *Sulphur*, in the same manner.

For all other eruptions, such as tetter, etc., see their respective heads, under "Skin Diseases."

For *crusta lactea*, or *milk crust*, see "Diseases of Children."

### 3. FACE-ACHE — NEURALGIA IN THE FACE.

(*Prosopalgia* — *Tic douloureux*.)

When this disease has been of a long standing or is constitutional, it requires medical advice and careful treatment to eradicate it; but if of recent date the following remedies may be used with benefit. For more information look under the head of "Toothache," as the same remedies there stated will be found to answer.

It appears, generally, in paroxysms of severe pain under the eye and in front of the ear; shooting over the other parts of the face, most generally on one side only, and sometimes accompanied with spasmodic twitchings of the facial muscles.

We give first the principal medicines in the different kinds of face-ache, and afterward their details, which must be read before a selection is made.

a. For face-ache of an *inflammatory* character (heat, redness of the face and thirst): *Aconite*, *Bryonia*, *Belladonna*, *Thuja*, *Chamomile*, *Nux vomica*.

b. For *rheumatic* or face-ache from cold: *Chamomile*, *Bryonia*, *Belladonna*, *Mercury*, *Pulsat.*, *Spigelia*.

c. For *gouty* (arthritic) face-ache : *Colocynth*, *Rhus*, *Spigelia*.

d. For purely *nervous* face-ache (neuralgia in the face, *tic douloureux*) : *Belladonna*, *Phosphorus*, *Capsicum*, *Spigelia*, *Platina*, *Hyoscyamus*.

e. For face-ache from abuse of *mercury* : *Carbo veg.*, *Hepar*.

SYMPTOMATIC DETAIL. — *Aconite*. Redness and heat of the face, with crawling pains on one side, or pains as from ulceration ; swelling of the cheek or jaws ; feverish heat, thirst, great exasperation, with agitation and tossing.

*Belladonna*. Pain following the nerve under the eye, and easily excited by rubbing that part ; or tearing, shooting pains in the bones, jaws, or cheekbones ; rigidity of the nape of the neck ; twitching of the eyelids ; spasmodic jerks of the facial muscles and distortion of the mouth ; heat and redness of the face ; chilliness and feverishness at the same time.

*Bryonia*. *Red face, burning hot*, with swelling on one side, or under the eyes and on the root of the nose ; pressing pains in the face, relieved by external pressure ; pains in the limbs ; chilliness followed by fever ; thirst ; tongue coated yellowish.

*Carbo veg.* Tearing and drawing in the cheek bones ; swelling of the cheeks and face ; night sweats ; great weakness.

*Colocynth*. *Tearing and shooting pains*, more on the left side of the face, extending into the head, temples, nose, ears, and teeth, with swollen face ; or when the pains increase at the slightest touch.

*Chamomile*. Puffed and swollen face, with *hardness, bluish color*, and *beating* of the cheek ; one cheek red, the other pale ; *spasmodic twitchings* of the facial muscles ; greatly excited and sensitive, or drowsy with feverish inquietude ; shivering, with internal heat ; desire to lie down and not be exposed to fresh air, particularly the wind.



*Capsicum.* Pains in the bones of the face, worse by being touched, and when going to sleep ; feeling of *numbness* of the cheek bone.

*Hepar.* Pains in the *bones of the face* ; worse by touch, and extending into the ears and temples ; heat in the face during the night.

*Hyoscyamus.* Pressing or cramp-like pains in the cheek bones.

*Mercury.* Tearing or shooting pains on one side, from the temples to the teeth ; worse at night and in the warmth of the bed ; too much saliva in the mouth and perspiration without bringing relief ; sleeplessness.

*Nux vomica.* Tearing and drawing pains, with swollen cheeks ; redness of one or both cheeks ; *tingling* in the face with beating in the muscles of the face ; worse by meditation and from wine or coffee.

*Phosphorus.* Tearing pains on the left side ; worse by moving the muscles of the face, when eating, etc. ; also, by contact.

*Pulsatilla.* Paleness alternated with redness of the face ; convulsive motions in the face ; painful sensitiveness of the skin of the face, as if sore ; shivering on one side of the face ; no thirst.

*Rhus.* Pressing, cutting pains in the cheek-bones ; with heat and roughness of the skin of the cheeks ; follows well after *Bryonia*.

*Platina.* Tingling, crawling pains, with numbness and coldness of the cheek on one side ; or spasmodic pain and pressure in the cheekbone ; worse in the evening, when it generally begins ; very nervous ; melancholy ; with red face, and thirst.

*Spigelia.* Jerking, tearing, burning, and pressure in the cheek bones ; worse by touch or motion ; with shining

swelling of the diseased part. Appears at a certain time of day; worse on stooping.

*Thuja*. Pain in the cheek bones, as if there was a boring; relieved by touch; spasmodic twitching of the lips; follows well after *Belladonna*; heat in the face, transitory; with perspiration in the face.

ADMINISTRATION.—Dissolve twelve globules of a remedy in half a teacupful of water, and give every hour a teaspoonful, until three teaspoonfuls are given, or the pain is relieved; then discontinue for two hours; after which, give the next remedy, if necessary. If, on giving *Belladonna*, it relieves, but not entirely, give afterward a couple of doses of *Belladonna* <sup>cc.</sup>, until better.

N. B. In very severe cases of neuralgia in the face I have seen the most beneficial effects from tincture of *Aconite* (three drops to half a teacupful of water, every fifteen or twenty minutes a teaspoonful); in such cases its external application on the parts principally affected is also recommended. Otherwise, apply nothing externally but hard pressure of the finger, under the eye, on the cheek bone, where the facial nerve lies, which often deadens the pain; also, coldwater bandages.

DIET AND REGIMEN.—The usual homœopathic diet. Abstain, even in health, from coffee and tea, and bathe frequently.

#### 4. WARTS AND SPOTS ON THE FACE.—CANCER OF THE FACE.

For *warts* on the face, use first the remedies stated in the article on warts, under “Skin Diseases;” and if these do not succeed, use *Caustic.*, *Dulcamara*, and *Sepia*; each remedy internally for four weeks, every week one dose (six glob.), and externally, wet the warts every evening with a solution of the same remedy (three glob. in a few drops of water.)

*Spots*, yellow (*hepatic*), on the face, require principally the use of *Sulphur* and *Sepia*, in the same manner as above.

*Cancer* of the face: see the remarks and treatment of "Cancer of the Neos."

DIET. — The usual homœopathic diet in chronic diseases.

#### 5. PARALYSIS OF THE FACE AND JAW.

In this disease, the muscles of the face have lost their power to act and be moved by the will; consequently, the affected part of the face or jaw hangs down; it mostly results from attacks of apoplexy, rheumatism, loss of fluids, and consequent debility, or suppression of an eruption. For more details, see the article, "Paralysis or Palsy." If it appears in an aggravated form, it must be treated by a physician; however, commence its treatment with the following remedies: *Belladonna*, *Graphites*, and *Caustic.*, every six or eight days one remedy, every other day one dose (six glob.), until better.

In paralysis of the jaw (*hanging jaw*), *Dulcamara*, *Arsenic*, and *Lachesis*, in the same manner. Beside, apply cold water, electricity, or galvanism, at the same time, if the patient is relieved by it.

DIET, as usual in chronic diseases.

#### 6. LOCKJAW. (*Trismus.*) See page 22.

## CHAPTER XI.

### AFFECTIONS OF THE TEETH, GUMS, MOUTH.

#### 1. TOOTHACHE. (*Odontalgia*.)

The proper culture of the teeth is important, in regard to health and comfort; and, as they form a part of the digestive system, their soundness depends a great deal on the healthy state of the stomach and bowels; consequently, whatever may derange these organs, will act more or less injuriously on the teeth, and ought to be guarded against. The extremes of hot and cold drinks should be avoided, and the teeth kept scrupulously clean; the best method for doing this is to brush them gently with cold water and a soft brush. All medicated tooth-powders are more or less injurious. If tartar adheres to the teeth, have it cautiously scraped off, by a competent person; and if any damage is visible, externally, consult immediately a skillful dentist; as he can easily repair it, while an unskillful practitioner makes the evil worse. We can not be too careful in the selection of a dentist.

If *habitually* predisposed to toothache, consult a homœopathic physician, who will be able to eradicate this complaint, which has either a scrofulous or rheumatic basis.

But to commence such a cure: if the toothache is recent, from cold, etc., take some of the following remedies, which may afford speedy relief.

TREATMENT. — In most cases of toothache, try first the following remedies, in this order: *Chamomile*, *Bryonia*, *Mercury*, *Sulphur*, if the patient is a male; and

*Chamomile*, *Belladonna*, *Mercury*, and *Sulphur*, if a female or child.

Give, according to the severity of the pain, from one to six hours a dose (four glob.), alternately, until better. If, however, among the remedies given below, one is found which will suit *exactly*, give this first of all.

In administering the following medicines, be careful not to give another one, when the remedy previously given has already afforded some relief; let it then have time to show its effect; say, from one to six hours. Do not use creosote, and other medicines, such as opiates; as they are very injurious to the general health, and to the teeth especially.

Neither allow the teeth to be extracted so very readily; as frequently it does not relieve, and can be obviated altogether, by the use of one of the following remedies. Select with care, and give as a dose, four globules at a time.

Before taking any medicine, cleanse the mouth of every thing which may have been used previously; particularly of those things which have a smell, such as creosote, etc.

The principal remedies recommended in this article for toothache are the following:

*Belladonna*, *Chamomile*, *Mercury*, *Nux vom.*, *Pulsatilla*, *Sulphur*, *Bryonia*, *China*, *Hyoscyamus*, *Ignatia*, *Rhus*, *Staphysag.*, *Aconite*, *Carbo veg.*, *Coffea*, *Hepar*, *Sepia*, *Silicea*, *Dulcamara*.

Before we give their detailed symptoms, we will state a few general indications, which are of great service in selecting the proper medicine.

First, look under the general indications, and on finding there a head applying to the case, select from among them the proper remedies by reading their detailed symptoms below, and comparing them with the present attack.

GENERAL INDICATIONS.—Toothache in children: *Aconite*, *Belladonna*, *Chamomile*, *Coffea*, *Ignatia*.

In females: *Aconite*, *Belladonna*, *Chamomile*, *China*, *Coffea*, *Bryonia*, *Ignatia*, *Pulsatilla*, *Sepia*.

In young girls of full habit: *Aconite*, *Belladonna*.

At the time of the menses: *Carbo veg.*, *Chamomile*, *Pulsatilla*, *Bryonia*.

During pregnancy: *Belladonna*, *Nux vom.*, *Pulsatilla*, *Staphysag.*, *Rhus*.

During nursing: *China*.

In hysterical females: *Ignatia*, *Sepia*, *Belladonna*.

In sensitive and nervous persons: *Aconite*, *Belladonna*, *Coffea*, *Hyosциamus*, *Ignatia*, *Nux vom.*

When caused by abuse of coffee: *Chamomile*, *Ignatia*, *Nux vomica*.

Of tobacco: *Bryonia*, *China*, *Chamomile*, *Hepar*.

Of mercury or calomel: *Carbo veg.*, *Hepar*, *Pulsat.*, *Sulphur*.

When in consequence of a chill: *Aconite*, *Belladonna*, *Chamomile*, *Coffea*, *Dulcamara*, *Ignatia*, *Mercury*, *Nux vom.*, *Pulsat.*

When caused by damp and cold air: *Nux vom.*, *Pulsat.*

If it is of a rheumatic nature: *Chamomile*, *Mercury*, *Bryonia*, *Bellad.*, *Sulphur*, *Pulsat.*, *China*, *Rhus*.

Of a nervous nature: *Bellad.*, *Coffea*, *Ignatia*, *Chamomile*, *Nux vomica*, *Hyosциamus*, *Sulphur*.

Of a congestive nature: *Aconite*, *Bellad.*, *Chamomile*, *China*, *Pulsat.*, *Hyosциamus*.

In carious teeth: *Mercury*, *China*, *Pulsat.*, *Staphysag.*, *Chamomile*, *Bryonia*, *Silicea*, *Sulphur*.

With affection of the bones of the face: *Mercury*, *Sulphur*.

With pains up to the eyes: *Pulsatilla*.

To the ears: *Chamomile*, *Mercury*, *Pulsat.*, *Sulphur*.



To the head : *Chamomile, Mercury, Nux vomica, Pulsatilla, Sulphur.*

With a swelled face : *Chamomile, Mercury, Nux vomica, Pulsat., Bryonia, Sepia.*

With swelled gums : *Aconite, Bellad., China, Hepar, Mercury, Nux vomica, Sulphur.*

With swelled submaxillary glands : *Carbo veg., Mercury, Nux vom., Staphisag.*

SYMPTOMATIC DETAIL.—*Aconite.* Feverishness, with great anxiety and restlessness, intense beating pains, with congestion to the head and intense burning in the face, which is swollen, and very painful. *Belladonna, Bryonia,* and *Chamomile* suit very well after it; or in alternation with *Coffea*, if the pain is intolerable (see *Coffea*.)

*Belladonna.* Great anguish and restlessness, or excessive sadness, with tendency to shed tears; sensation of ulceration in the teeth and gums; *drawing*, tearing, incisive, or shooting pains in the teeth, face, and ears, *aggravated in the evening, after lying down at night; bleeding of the teeth* on sucking them; swelling of the cheek; *dryness of the mouth and throat*, with *great thirst*; worse in the *open air* and *when eating*; *heat and redness of the face*, with beating in the head and cheeks; burning and redness of the eyes.

After it *Mercury, Hepar, Chamomile, Pulsatilla*, suit.

*Chamomile.* Violent jerking, drawing, and shooting pains, almost *insupportable*; worse at *night*, when the patient is warm *in bed*; with swelling and redness of the cheeks and irritability of mind; disposition to cry, during the pain; swelling of the gums; enlargement of the submaxillary glands; oftener on one side, or without knowing which teeth ache most. Shooting or pulsative pains in the head, ears, and face; worse after drinking any thing *hot* or *cold*; particularly after *drinking coffee*; *one cheek red*, the

other pale ; violent agitation, and tossing about ; yet great weakness, almost to fainting.

After it, *Bryonia*, *Mercury*, *Coffea*, *Sulphur*, *Belladonna*, follow well.

*Coffea*. Violent pains, during which the patient is utterly distracted ; with trembling, tears, tossing and agitation (in alternation with *Aconite*.) Indescribable pains ; worse at night, or after a meal.

After it, *Aconite*, *Belladonna*, *Hyosciamus*, *Sulphur*, suit well.

*Carbo veg*. Drawing, tearing, or beating pains in the teeth, with painful tendency to the touch, especially after eating ; the gums bleed easily ; with a feeling of ulceration and looseness of the teeth ; worse in damp weather, or when hot, cold, or salted things, are put into the mouth ; or when *Mercury* has not relieved the pains.

*China*. When debilitated by loss of fluids, as after having been bled, or during nursing, etc. ; or when the pain is beating, drawing, jerking ; dull pain in carious teeth ; worse after eating ; at night, or when touched ; mitigated by pressing and clinching the teeth.

After it, *Mercury*, *Pulsatilla*, *Sulphur*.

*Bryonia*. Pains in carious teeth, and in the gums, as if sore and excoriated, with the feeling as if the teeth were loose ; jerking and drawing pains, with a sensation as if the teeth were loose or too long, but are not so ; especially when eating, or after a meal ; the pains force one to lie down ; are worse at night ; or on taking any thing hot into the mouth ; also when lying on the side affected. It suits well after *Chamomile*, or before *Mercury* ; also, in alternation with *Rhus*, in those toothaches which are caused either by wet weather, or accompany bilious rheumatic attacks. (See this article.)

*Dulcamara*. In toothache accompanied by diarrhea,

from cold or from a chill ; suits well in such cases after *Chamomile*, when that remedy has proved insufficient ; also, in cases of toothache from cold, where it has befallen the saliva glands, with salivation ; where *Belladonna* and *Mercury* have proved insufficient.

*Hepar*. Suits well for persons who have taken a great deal of mercury ; and when the toothache precedes the formation of gum-boils ; in this case, in alternation with *Silicea* ; besides, it is suitable after *Mercury* or *Belladonna*, when there are painful or erysipelatous swelling of the cheek, or jerking and drawing pain in the teeth ; worse on clenching the teeth, when eating, in a warm room, or at night.

*Hyosciamus*. The pains, when this remedy should be used, are very *violent* ; *tearing* ; *pulsation* from the cheek to the forehead, after *taking cold*, or *in the cold air*, with congestion to the head ; red and hot face ; clenching of the teeth ; convulsive jerkings of the fingers, hands, or arms ; great nervous excitability ; red and brilliant eyes, looking around wildly ; pains are so violent as to almost cause raving.

*Ignatia*. In persons of a sensitive temperament, yet of a mild character ; or, now lively and gay, and soon after, disposed to weep. The toothache for which this remedy should be used, is distinguished by a sensation as if the teeth were broken ; caused by taking cold ; by depressing mental suffering ; or when pains are worse after drinking coffee or smoking tobacco ; after lying down, or in the morning on waking. It suits well after *Chamomile*, *Nux vomica*, or *Pulsatilla*.

*Mercury*. One of the principal remedies for toothache in children, or when there are *shooting* pains in the carious teeth ; and *tearing* pains in the roots of the sound teeth, through the whole side of the face, into the ears ; with swelling of the cheek ; painful on touch ; swelling of the

*submaxillary glands*; *salivation*; worse at night; but particularly so, even insupportable, in the *warmth of the bed*; the pains are excited by *cool and damp air*, or by *eating or drinking any thing cold*. It suits well after *Belladonna* or *Dulcamara*; or before *Hepar*, *Carbo veg.*, or *Sulphur*.

*Nux vomica*. Especially in persons who lead a sedentary life, or indulge habitually in coffee or ardent spirits; pain as from *excoriation*, or *jerking*, *drawing*, particularly in the *carious teeth*; swollen and painful gums, with *pulsation as in an abscess*; worse at night, in the *morning on waking*, *after dinner*, or *during a walk in the open air*; irritable, quarrelsome, and peevish temper, with tendency to constipation and piles. It suits well before *Mercury* or *Sulphur*.

*Pulsatilla*. Especially in persons of a mild, quiet, and timid character, with tendency to shed tears; toothache from cold, with *earache* and *headache* on one side, mostly on the left; tearing, shooting, or *jerking* pain, *as if the nerve were tightened and suddenly relaxed*; pulsation; gnawing pain, with *pricking* in the gums; pains extending to the *face*, *head*, *eyes*, and *ears*, mostly on one side, with *pale-ness of the cheeks*, heat in the head, *shivering in the body* and *difficulty of breathing*; worse in the *afternoon or evening*, or else *after midnight*; aggravated when *warm in bed* or when *seated in a warm room*; mitigated by *cool air* or *when out of doors*. It suits well before or after *Mercury* or before *Sulphur*.

*Rhus*. Particularly in rheumatic toothache in wet weather, with pains in the extremities and head, shivering and restlessness; in such cases in alternation with *Bryonia*. (See this remedy.) Or *tearing*, *jerking*, *shooting*, *tingling* pains, worse in the open air or at night, when it renders the patient very restless; mitigated by *external heat*. It suits well after *Belladonna*.

*Sepia.* Pulsative and shooting pains extending into the ears and arms as far as the fingers; tendency to sick headache; constipation and piles. It suits well after *Belladonna*.

*Silicea.* Shooting pains, with swelling of the jaw-bones, which ache rather worse than the teeth; worse in the night; fistulas or gumboils. It suits well after *Mercury* or *Hepar*.

*Sulphur.* Tearing, jerking, and pulsative pains, particularly in the carious teeth, with swelling of the cheek; congestion and beating in the head; shooting pains in the ears; constipation with tendency to piles; the toothache is worse in the evening or at night, when warm in bed or being exposed to the open air or a current of air; also from cold water. It suits well after *Aconite* or *Mercury*.

*Staphysag.* When the toothache affects carious teeth, or the teeth become so, and break off easily, with paleness, whiteness, ulceration, swelling, and tenderness of the gums; worse immediately after eating or drinking anything cold, or by inhaling the cold air; also in the morning or at night.

## 2. CARIES OF THE TEETH. (*Carious Teeth.*)

For pains in carious teeth the principal remedy is *Antimon.*, *Chamomile*, or else *Staphysag.*, *Nux vomica*, *Mercury*, *China*, *Pulsatilla*, and *Sulphur*. Give of these every three or six hours a dose (four glob.) until relieved.

For a disposition to caries in the teeth, give first *Staphysag.*, then *Sulphur*, then *Calcareo carb.*, every week two doses of one remedy, commencing with the first, each dose six globules.

Beside this treatment consult a good dentist, as teeth may frequently be saved by a simple operation, which without it would decay in a short time.

## 3. OFFENSIVE BREATH.

If it is from carious teeth, consult a good dentist immediately; or, if that is impossible, clean out the hollow teeth yourself, with cotton or rolls of paper, and fill the cavities with wax.

Offensive breath, caused by neglect in cleaning the teeth from tartar or particles of food attached to them after eating, can be removed only by the most careful attention to rinsing the mouth frequently and using the brush freely.

Sometimes offensive breath is a symptom of other diseases, such as swelling and inflammation of the gums and glands from the *abuse of Mercury*. (See this article.) Where it can not be traced to any particular cause, however, try to relieve the patient by giving the following remedies:

If it appears only in the *morning*, give *Arnica*, *Nux vomica*, *Belladonna*, *Silicea*, *Sulphur*.

If *after a meal*, *Chamomile*, *Nux vomica*, *Sulphur*.

If in the *evening* or at *night*, *Pulsatilla*, *Sulphur*.

If in *young girls* at the age of puberty, *Aurum*, *Pulsat.*, *Sulphur*, *Bryonia*, *Belladonna*, *Sepia*.

Administer every evening of one of the above remedies indicated one dose (six glob.) for one or two weeks, when another remedy may be selected, if necessary.

N. B. The smell caused by onions, garlic, or horse-radish, may be destroyed by taking a little wine, or eating a pear or boiled beets.

## 4. BAD TASTE IN THE MOUTH.

A great many diseases, such as fevers and disorders of the stomach, have a bad taste in the mouth as an accompanying symptom, which will disappear with the principal complaint; this must be looked for in the proper chapters; but



sometimes it appears as a solitary symptom, for which the following remedies may be given.

ADMINISTRATION. — Take one dose (four glob.) of the appropriate medicine once a day for a couple of days, which is generally sufficient. If very bad take two or three times a day.

TASTE : *Inspid*—*Bryonia*, *Rhus*, *Dulcam.*; if after drinking, *China*; if after a meal or in the evening, *Thuja*.

*Coppery*—*Cuprum*, *Rhus*, *Cocculus*.

*Earthy*—*China*, *Pulsatilla*, *Sulphur*.

*Metallic*—*Lachesis*, *Mercury*, *Nux vomica*.

*Mucous or slimy*—*Mercury*, *Chamomile*, *Pulsatilla*, *Lycopodium*, *Belladonna*, *Rhus*.

*Putrid*—*Arnica*, *Bryonia*, *Rhus*, *Sulphur*, *Nux vomica*, *Mercury*, *Pulsatilla*, *Sepia*.

If worse in the morning — *Rhus*, *Nux vomica*, and *Sulphur*.

If after a meal—*Rhus*.

*Bitter, bilious*—*Bryonia*, *Pulsatilla*, *Arnica*, *Antimon. crud.*, *Chamomile*.

*Acid*—*Pulsatilla*, *Mercury*, *Nitric acid*, *Sepia*, *Sulphur*.

*Sweetish*—*Mercury*, *Pulsatilla*, *Sulphur*.

*Saltish*—*Mercury*, *Nux vomica*, *Tartar emet.*, *Arsenic*, *Carbo veg.*

For loss of taste—*Belladonna*, *Hepar*, *Bryonia*, *Phosphorus*, *Silicea*, *Natrum muriat.*

## 5. INFLAMMATION AND SWELLING OF THE TONGUE. (*Glossitis*.)

This disease is not of frequent occurrence. When it appears the tongue is swollen, either in part or wholly, often to a great extent, so as to fill the whole cavity of the mouth, thereby rendering it extremely difficult to speak, swallow, or even breathe. The disease, if not checked in time by the proper homœopathic remedies, frequently runs into sup-

puration or hardening of the substance of the tongue, after which the fever, which is generally very intense during the height of the inflammation, gradually subsides, or it terminates in gangrene of the tongue and surrounding parts.

CAUSES. — This disease is mostly caused by local injuries, such as from the points of decayed teeth, the sting of a bee, or by rheumatic and catarrhalic influences.

TREATMENT. — In the commencement of the disease, if the tongue has been injured externally, give *Arnica*, rinsing the mouth with a solution of ten drops of the tincture to a teacupful of water three or four times a day. If injured by *burning* or scalding, give *Urtica urens*, in a similar manner.

But if real inflammation of the substance sets in, and there is great fever, give first *Aconite*, in solution (twelve globules in half a teacupful of water), every hour or two a teaspoonful until better.

*Mercury* is almost a specific after *Aconite*, particularly when there is great swelling and heat of the tongue and surrounding glands; it may be given either alone, every three or four hours a dose (four glob.), or alternately with *Belladonna*, if the mouth is very dry and the pain excessive, with great thirst and congestion to the head.

*Pulsatilla* is beneficial in cases caused by suppression of gouty, or hemorrhoidal affections.

If indications of *gangrene* appear *Arsenic* and *Lachesis* are the principal remedies, alternately to be given as above in *Belladonna* and *Mercury*.

If ulceration of the tongue takes place, see "Ulcers."

DIET AND REGIMEN — As in fevers.

## 6. CANKER OF THE MOUTH. (*Stomacace.*)

This disease consists in an inflammation and ulceration of the inside of the mouth and surrounding parts, preceded by

general lassitude, loss of appetite, gastric derangement and feverishness. The local affection soon makes its appearance with heat, redness, and sensitiveness of the gums, which swell, become spongy, and retract from the teeth; presently small ulcers appear on the gums, spreading to the lips, tongue, palate, and inside of the cheeks, smelling very offensively.

Sometimes the glands of the throat and mouth swell and participate in the inflammation, secreting large quantities of a putrid, acrid saliva, which runs constantly from the mouth. In such a state the patient looks like one who has been deeply salivated by the *abuse of mercury*; we find, also, the same weakness and feverishness.

TREATMENT. — *Mercury* must be given first, as it is almost the *specific* in this disease, but not, of course, in those cases which have been produced by its own abuse; in such cases see “Diseases from the abuse of Mercury.”

ADMINISTRATION. — Dissolve twelve globules in half a teacupful of water, and give every three or four hours a teaspoonful until four or six are taken; if better the patient needs no more medicine; if not better or about the same, select one of the following remedies, preparing and giving it in the same manner as the mercury.

*Capsicum* Particularly in *phlegmatic* persons with a *full habit*, leading a *sedentary life*; or where there are *burning* blisters or vesicles on the tongue, and swelling of the gums.

*Dulcamara*. Where the patient has taken cold, or where the glands of the neck participate, feel sore, and are swollen.

*Arsenic*. Ulceration on the margins of the tongue, with *violent burning* pain; swelling and easy bleeding of the gums; *great debility* and *restlessness*, with a desire to *drink frequently* but *not much at a time*.

*Borax.* Ulceration of the gums with *thrush* in the mouth and on the tongue; bleeding easily, particularly in children.

ADMINISTRATION.—See “Thrush” in “Diseases of Children.”

*Carbo veg.* Ulceration of the gums and tongue, with *profuse bleeding*; burning sensation, and excessive fetidness of the ulcers, and difficult movement of the tongue. It suits well after *Arsenic*, or where *Mercury* has caused salivation.

*Nitric acid.* If *Mercury* has proved inefficacious, or if the gums are swollen, look whitish and bleed easily with looseness of the teeth; excoriations in the mouth; salivation and a *putrid odor* of the mouth.

*Nux vomica.* In the worst cases, and then in alternation with *Arsenic*; principally in lean persons, who lead a sedentary life; or where there is *putrid* and *painful swelling* of the gums; fetid ulcers cover the whole inside of the mouth, which smells very offensively; the face looks discolored, with hollow cheeks, dull eyes; emaciation; constipation and irritability.

*Sulphur.* At the end of the cure, or when there is swelling of the gums with pulsative pains; thrush in the mouth and on the tongue; blisters which burn when eating; offensive and sour smell from the mouth, with a thickly coated, whitish or brownish tongue; constipation, or greenish, slimy evacuations; restlessness at night. It suits well after *Arsenic*, *Mercury*, or *Nux vomica*.

*Externally* use nothing but cold water, with which the mouth should be frequently washed; sometimes lemon juice or sage tea is beneficial, at least is not objectionable, when their use can be borne.

DIET.—Only the mildest nourishment, such as gruels, milk and water, etc., is allowed; no meat or soup of meat until the patient is decidedly convalescent.

## 7. INFLAMMATION OF THE PALATE.

See "Sore Throat."

## 8. ABSCESS IN THE GUMS. — GUMBOILS. — SWELLING OF THE JAW-BONE.

Before an abscess in the gums, or a gumboil, is formed, there is much inflammation and swelling in the parts affected, which requires the use of *Aconite* and *Belladonna* in alternation, every three or four hours a dose (four glob.), until better; or, if not better after twelve hours, *Mercury* and *Hepar* in the same manner. If the boil has formed, give *Silicea*, evening and morning a dose (four glob.), for three or four days; and if not better in eight days, give *Calcarea carb.*, in the same manner, and after this *Staph-asag*.

The same treatment is suitable for swelling and ulceration of the *jaw-bone*, produced by carious teeth, or after the extraction of a tooth.

DIET light, and according to the homœopathic rules.

9. SCURVY. (*Scorbutus*.)

This disease exhibits all the symptoms described in the previous article, on "Canker of the Mouth." It differs, however, as regards its origin and progress, which is slow and chronic.

The real scurvy principally affects persons who are in circumstances which deprive them of fresh provisions (such as sailors), and expose them to damp and cold. If, at the same time, there is want of exercise, impure air, and uncleanness, together with depression of spirits, the disease will progress rapidly.

It begins with swelling of the gums, which become spongy, discolored, and bleed easily; producing fetid

breath; afterward the lower extremities begin to swell, and the skin shows tettery and livid spots; face looks lead-colored and bloated; urine is offensive, and the stools are extremely fetid; hemorrhages from the nose, teeth, tongue, and the urinary organs now ensue; the general debility and melancholia increases, and ulcers make their appearance on various parts; after which, a general dissolution of the system takes place.

TREATMENT. — A great part of the cure of this disease lies in the removal of its causes; and this should be done, if possible; substituting fresh provisions for salt; also, fresh fruits and vegetables, oranges, lemons, cider, vinegar, spruce beer, sour-kROUT, etc. Besides, give the following remedies:

*Mercury.* If this has not already been used to excess in the disease (in which case *Carbo veg.* should commence the cure), give it for the following symptoms:

*Red, spongy, ulcerated, and easily bleeding gums, with burning pains at night; looseness of the teeth;* the whole inside of the mouth covered with aphthæ or thrush; fetid smell of the mouth, and ulcers; profuse discharge of offensive or sanguineous saliva; loose, scalding evacuations; dark red, fetid urine; great exhaustion and debility, with inexpressible feeling of depression of body and mind.

ADMINISTRATION. — Dissolve twelve glob. in half a tea-cupful of water, and give every four or six hours a tea-spoonful, for six times; then discontinue the medicine for twenty-four or thirty-six hours, and, if improving, for a longer time; if not, however, select the next best remedy.

*Carbo veg.* Particularly in persons who have taken much mercury, or when there are burning ulcers in the mouth, which bleed profusely at times, and discharge an acrid, fetid matter; great weakness in body and mind.



*Nux vomica.* For the same symptoms as stated in “Canker of the Mouth.” (See this article.)

*Arsenic.* See “Canker of the Mouth.” Besides these symptoms, it will mitigate the sufferings of the patient, in the most severe forms of this disease; ulcers are either very painful or not sensitive; look dark and livid; so does also the whole skin, which is bluish, cold, and extremely dry; extreme restlessness, particularly at night.

*Nitric acid* and *Sulphur.* See “Canker of the Mouth.”

Administration of these remedies, the same as *Mercury*, above.

DIET. — As in “Canker of the Mouth.”

#### 10. SALIVATION. (*Ptyalism.*)

This disease is not always caused by the abuse of mercury; but frequently occurs after taking cold, or accompanies attacks of fever, such as scarlet, etc. Sometimes it appears in diseases of the reproductive system, as a critical discharge by the action of nature; and is then very beneficial.

*Salivation* (a profuse discharge of saliva or spittle) is only a symptom of that disease to which it has given its name. The disease consists in an irritation, swelling, and inflammation of the glands of the mouth and throat; at times, of the whole mucous membrane, which, in health, secretes the necessary saliva. It is evident and rational that, in order to stop the salivation, we must remove the inflammation of the gland, or mucous membrane, by the appropriate remedies; which, if done, will of itself reduce the discharge of saliva to its legitimate quantity.

All astringents which will directly diminish salivation are injurious; being only harmless when they produce no effect. Mild washes, or gargles, such as milk and water,

may be used with benefit. Depend, however, principally upon the following remedies :

TREATMENT. — If *mercury* or *calomel* is the cause, *Hepar sulph.*, *Lachesis*, *Belladonna*, *Opium*, *Carbo veg.*, *Dulcamara*, *Nitric acid*, and *Sulphur*, in alternation, every six or twelve hours a dose (four glob.), until the patient feels relieved, if not of the quantity of saliva discharged, at least of the ill feelings generally experienced in the whole system. If caused by *cold*, the first remedy to be used is almost always *Mercury* ; and if not better after twelve or twenty-four hours, give the above-named remedies in the same order. If salivation occurs in fevers, as a crisis, do not interrupt its flow, unless it becomes very weakening and dangerous, in which case use the above-named remedies. Besides, compare “ *Stomacace.*”

DIET AND REGIMEN. — Diet of the mildest kind, gruels, milk and water, rusks, crackers soaked in milk, etc.; no meat.

## 11. PARALYSIS OF THE TONGUE.

Paralysis of the tongue is one of the symptoms by which apoplexy shows itself ; and in that case can only be cured by the removal of the whole disease. See, in that case, “ *Apoplexy.*” But where the disease has only palsied the tongue and made it useless, give *Lachesis*, if it is experienced more in the *left side* of the tongue, and *Belladonna*, if more on the *right side*.

Besides, *Caustic*, *Hyosciamus*, *Opium*, *Stramonium*, *Aconite*, *Sulphur*, in rotation, if necessary, one after the other, giving every two or three days a dose (four or six glob.), until improvement begins.

DIET AND REGIMEN. — The usual homœopathic diet in chronic diseases.

## 12. DIFFICULTY OF SPEECH.—STAMMERING.—STUTTERING.

A person may lose his speech in consequence of spasms or paralytic strokes, and if this is the case see their respective chapters. If it is the well-known stammering or stuttering, medicines may be given to decrease the general sensitiveness of the nervous system, and particularly the nerves of speech. Yet the treatment must lie principally in acquiring such habits of speech as will bring it completely within the control of the speaker; reading aloud is of service, *intonating* every word, and *beating time with the fingers* on any thing near. If this is patiently persevered in for a length of time, it will overcome the disease. A kind treatment by parents and teachers is necessary, particularly where lessons are to be repeated, or messages received.

For diminishing the general sensitiveness of the nervous system, give the following remedies, which will have, also, a specific bearing upon the local disorder.

*Lachesis, Belladonna, Mercury, Sulphur, Stramonium, Natrum muriaticum, Graphites.* Give each remedy for six weeks, every week one dose (six glob.), until better.

DIET.—Abstain only from coffee and spices.

## 13. RANULA, OR SWELLING UNDER THE TONGUE.

This disease consists in a swelling of the salivary glands under the tongue, where it appears as a bluish eminence, not painful, but in its final growth impeding speech materially.

The principal remedies are *Thuja, Mercury, Calcareo carb.*, and *Sulphur*—each remedy to be given for six weeks, every week two doses (six glob. each), until better. These remedies may be used at the same time externally, by dis-

solving a few globules in a little water, and keeping it in the mouth for a minute once a day.

DIET AND REGIMEN. — Abstain from coffee and spices.

## CHAPTER XII.

### AFFECTIONS OF THE THROAT.

SORE THROAT.—QUINSY. (*Angina faucium*.)

#### *a. Common Sore Throat.*

This well-known disease consists in an inflammation and swelling of the back part of the throat, palate, and tonsils, which impedes the respiration to a greater or less extent, and occasions fever. It appears in different degrees of intensity, from the slightest irritation of the throat, which vanishes quickly, to the highest degree, the formation of an abscess in the tonsils, or adjacent parts, which is very painful, and obstructs respiration to an alarming extent.

The following list of medicines will exhibit the different degrees of intensity in the symptoms, according to which the remedy must be chosen.

TREATMENT.—I will first remark here, that perhaps in no disease are the beneficial effects of cold water so evident as in sore throat; and advise, therefore, its use in the beginning of this complaint, if it is possible.

In the evening, when going to bed, put a wet bandage around the throat, and cover it with a dry cloth; repeat the same in the day time, if remaining within doors; and at the same time select the proper homœopathic remedies, as the application of the cold water will not interfere with the action of the medicine.

I would also advise sponging the throat and breast with cold water, for persons who are habitually subject to quinsy, in order to strengthen the weak parts. This ought to be done *regularly*, every morning.

Another precaution which such persons ought to take, is to avoid, as much as possible, long-continued or loud talking, particularly at times when the weather is inclement, damp, and cold, producing colds and rheumatism.

They should also strictly avoid any irregularity in diet and habits, and take particular care not to overload the stomach.

ADMINISTRATION OF THE FOLLOWING MEDICINES. — Remember that the patient always ceases taking any medicine, as soon as he feels better, or is, at least, not worse. In the latter case, take the last appropriate medicine at longer intervals, from time to time, until entirely relieved.

Having selected a remedy, dissolve of it twelve globules in half a teacupful of water, and give according to the severity of the case ; every one, two, three, or six hours a teaspoonful, until from three to six teaspoonfuls have been taken, when either the patient is better, or another remedy is needed ; if the latter, prepare and give this one in the same manner as above.

If the patient can not swallow, let him keep the medicine for a minute on the tongue, and then spit it out again, as the effect by that time is fully there ; or give the medicine dry on the tongue ; in which case, do not repeat it as often, say every two, three, four, or six hours a dose (four glob.)

Before giving the list of medicines and their symptoms, I will state that in the commencement of most cases of sore throat, *Chamomile*, *Bryonia*, and *Belladonna*, in alternation, every four or six hours a dose, will be found sufficient, if taken early. If, however, the disease still progresses, select the more homœopathic remedy from among



the following: *Aconite*, *Belladonna*, *Bryonia*, *Rhus*, *Chamomile*, *Lachesis*, *Mercury*, *Pulsat.*, *Nitric acid*, *Dulcamara*, *Ignatia*, *Carbo veg.*, *Nux vom.*, *Capsicum*, *Hepar*, *Silicea*, *Sulphur*; or, if the angina (sore throat) is caused by the abuse of mercury, look principally for *Belladonna*, *Carbo veg.*, *Hepar*, *Lachesis*, *Lycopodium*, *Sulphur*.

SYMPTOMATIC DETAIL.—*Aconite*. Particularly if at the beginning the patient has much fever, heat, thirst, swallows with pain and difficulty; the throat inside exhibits deep redness, causing a pricking sensation, worse when speaking.

N. B. In inflammatory affections of the throat, *Aconite* ought invariably to commence the treatment: after it a suitable remedy is

*Belladonna*. When there are *excoriating* pains, with sensation of scraping and enlargement; generally a *dryness* in the mouth and throat; when swallowing the pain *shoots* into the throat and *up to the ear*; sometimes can not swallow at all; the *drink escapes through the nostrils*; *violent thirst*, with *dread of drinking*; the throat itself inside shows a *bright redness* without much swelling, or a *deep intense redness* with *considerable swelling* of the tonsils, palate, and glands, with tendency to the formation of *abscess* or *ulceration*. Sometimes much slimy, whitish mucous in the mouth and throat; swelling of the muscles and glands of the neck and nape of the neck, with violent fever; hot, red, and swollen face; red eyes, with dread of light.

After it *Mercury*, *Hepar*, or *Lachesis*, is suitable.

*Bryonia*. Sensations of *shooting*, with dryness of the throat and difficulty of speech; after getting cold in damp cold weather, or after over-heating one's self, or drinking ice-water; frequently with pains in limbs and head, chilliness and thirst; in such cases alternate with *Rhus*, every

two or three hours a dose (four glob.) till three doses of each are taken, or until better.

*Rhus.* In most cases where *Bryonia* is indicated (see the preceding "*Bryonia*"), or where there is *obstructed swallowing* as from *contraction of the throat*.

*Chamomile.* Especially in sore throats of children, or where it is caused by *checked perspiration*; the glands of the neck, under the chin, and the tonsils are swollen; in the throat itself a feeling of *enlargement*, with deep redness of the parts affected; inability to swallow *solid food*, especially when lying down; *tingling in the throat*, with *hacking cough*, hoarseness and *rough voice*. If fever is present toward evening; heat and shivering alternate with *redness* (especially of *one*) of the cheeks; *irritability*, with cries and tears; the child wishes to be constantly carried in the arms, and wants things which, upon receiving, it throws angrily away.

*Lachesis.* Swelling, redness, and feeling of excoriation of the tonsils and *palate*; *constant need to swallow*, with a sensation as of a *tumor* or *lump* requiring to be *swallowed*; swallowing is obstructed, particularly of *liquids*, which often escape through the nostrils; worse in the afternoon and morning, after sleeping, or from slight pressure on the neck. Sometimes it is necessary to give *Lachesis* after other remedies, particularly after *Belladonna* and *Mercury* have mitigated the symptoms, but there still remains burning and *dryness in the throat*, *extensive swelling*, especially of the *left tonsil*, with difficulty of breathing, or danger of suffocation; or where there are small ulcers in the throat, or on the left tonsil; in which case, the remedy should be given in alternation with *Lycopodium*; particularly when the bowels are constipated, and on the tonsils in the throat appear very *small yellow eminences* or *tubercles*. (See "*Ignatia*.")

*Mercury.* Suits frequently in the beginning of a sore throat, when caused by taking cold, and, in addition to the throat symptoms, there is a rheumatic drawing pain in the head and nape of the neck; perspiration which affords no relief; the throat symptoms are, *violent shooting in the throat and tonsils, especially when swallowing*, which extend to the ears and glands of the face and neck; or there is a burning in the throat, and pains as from excoriation, swelling and great *inflammatory redness* of the *parts affected*; elongation of the palate, *difficult swallowing*, especially of *drinks*, which escape through the nostrils. In this case give it in alternation with *Belladonna*. If an abscess in the throat can not be avoided, *Mercury*, in alternation with *Lachesis* and *Hepar*, are the principal remedies to promote its maturation, and even after it has broken, *Mercury* is necessary, with *Hepar* and *Silicea*, to promote its healing; particularly when the patient feels worse *at night*, or in the evening, when he feels chilly and hot at the same time; *much saliva in the mouth*, with perspiration, which affords no relief; tongue thickly coated.

*Pulsatilla.* Suits very frequently in females, or persons of a mild phlegmatic character; after having taken cold, and the throat *feels swollen* inside; tonsils and palate look dark and red, and *dark red veins appear in the throat*, which is covered with *tough mucous*; the patient feels chilly toward evening.

*Nitric acid.* Is very suitable after *Mercury* has been given, or in alternation with it where the throat is filled with small, superficial ulcerations, of a whitish appearance, like thrush, with bad mercurial smell; also, in that form of angina caused by abuse of *Mercury* (see this article.)

*Carbo veg.* Suits well after *Mercury* and *Nitric acid*, if these remedies have not been sufficient against the ulcers in the throat; or after *Bryonia* and *Rhus* have been given,

and there yet remain severe *burning* and *pricking* pains, particularly in wet weather.

*Nux vomica*. Suits very well after *Chamomile*, or in persons of a bilious choleric character, when they have a *soreness in the throat*, with a feeling of *rawness*, *hoarseness*, particularly when inhaling cool air, or during the swallowing of the saliva; or when the palate is elongated and swollen, with constipation and thirst. *Nux vomica* is a principal remedy for ulcerated sore throat. (See the next article.)

*Ignatia*. Sensation as of a *plug* in the throat, with *shootings* into the ear, especially when *not swallowing*; greater difficulty in swallowing *liquid* than solid food; the tonsils swollen, indurated, or covered with small ulcers or tubercles. Compare *Chamomile*, *Nux vomica*, *Pulsatilla*, *Lachesis*, *Mercury*.

*Capsicum*. In cases where *Nux vomica* seemed to be indicated, or where there is a *burning* roughness in the throat, with feeling of contraction and stiffness, besides sneezing and running of the nose, also of the eyes. In cases of this kind it relieves very quickly.

*Hepar*. Often after *Belladonna*, *Mercury*, *Lachesis*, or when there are stitching pains in the throat, or in the swollen tonsils; particularly when *swallowing* or *coughing*; violent pressure in the throat, with danger of suffocation; necessary in maturing abscesses. (See *Mercury*.)

*Silicea*. After an abscess has broken, *Silicea*, in alternation with *Hepar*, is recommended, until it has closed and healed.

*Sulphur*. Swelling of the throat and tonsils, with pain as from *excoriation*; besides a feeling of *constriction*; follows well after *Mercury* or *Ignatia*.

DIET *light*. See "Inflammatory Fevers."

EXTERNAL TREATMENT.—Nothing but gargling with warm

water and milk ; or for an abscess of the throat a warm linseed poultice, until it breaks.

*Erysipelas of the Throat*, as it appears mostly in scarlet fever, measles, small pox, etc., sometimes, however, independent of these diseases, requires, in preference, *Bellad.*, *Aconite*, *Nux vomica*, *Mercury*, *Lachesis*, *Sulphur*, *Arsenic*, *Carbo veg.*, *Ignatia*, as the detailed symptoms above may indicate.

For angina caused by *syphilis* or *gonorrhea*, give mostly *Mercury*, *Nitric acid*, *Thuja*, *Lachesis*, or *Sulphur*.

If it arises from *foreign substances* in the throat, such as splinters, etc., give first *Aconite*, as usual ; but if, notwithstanding, swelling and heat make their appearance in the wounded part of the throat, look above, under “*Erysipelas of the Throat*,” for the principal medicines, which in such cases are preferred.

For *constitutional* or *chronic angina*, where a person is liable to these attacks upon the slightest occasion, commence the cure by selecting from *Hepar*, *Lachesis*, *Lycopodium*, *Sulphur*, *Bellad.*, *Nitric acid*, *Nux vomica*, *Thuja* ; and take them as needed, but not oftener than once a day ; and if the case is not too urgent, as far apart as every two or three days one dose (four or six glob.) until better.

*b. Ulcerated and Putrid Sore Throat. (Malignant Quinsy.)*  
(*Angina Gangrænosa.*)

This is one of those dangerous diseases, the treatment of which will seldom be attempted by domestic practice alone ; yet, as persons may be in circumstances where they could not procure the aid of a homœopathic physician, I will give below the symptoms and treatment of this disease, as plainly and fully as it can be done in a work of this kind.

DIAGNOSIS.—This disease, in its most virulent form, ap-

pears mostly as an epidemic among children, when scarlet fever is raging, which it often supplants, or accompanies. In this form, it is highly infectious.

In a milder form, it may occasionally attack any person, child or adult, who, from derangement of the digestive organs, or predisposition to throat diseases, is more subject to it.

It commences with a chill or shivering, followed by fever; great languor and oppression, with or without nausea; eyes suffused, with a reddish hue; cheeks crimson; nostrils inflamed, with a thin acrid discharge, causing excoriation of the nose and lips; pulse weak and very quick, to indistinction; besides, the patient is very restless.

On examination, the palate and visible part of the throat is found to be of a bright redness and swollen; which, together with the tonsils, is soon covered with numerous ulcers, coated with grayish-yellow crusts, which frequently separate in large pieces. At this time, the whole interior of the mouth, and even the lips, are involved in the destruction of the mucous membrane. The fever and restlessness increase, as also the difficulty of swallowing and breathing.

The breath becomes extremely offensive; the glands around the neck swell, and the skin on the neck and face assumes a shining glassy appearance; finally, drowsiness sets in, and the young patient, if not soon relieved, dies with symptoms of apoplexy or suffocation.

The milder form exhibits the above symptoms, with less severity, and terminates, in most cases favorably, on the third or fifth day, by general perspiration; breathing then becomes easier; the pulse less frequent and stronger; quiet sleep ensues, and the expression of the face grows more lively.

TREATMENT. — The typhoid character of this disease requires medicines for that class of fevers, such as *Bellad.*,



*Lachesis*, *Arsenic*, *Carbo veg.*, *Sulphur*; and the peculiar affections of the throat, such as *Mercury*, *Nitric acid*, *Nux vomica*.

*Belladonna*. In the beginning of the disease, when there is dryness of the throat and choking, fear, restlessness, even delirium. For further indications and

ADMINISTRATION. — See "*Belladonna*," under "Scarlet Fever."

*Mercury*. If profuse secretion of saliva makes its appearance, together with ulceration in the throat and mouth, which must be carefully observed, give *Mercury* either alone, in the same manner as *Bellad.*, above referred to, or in alternation, with *Nitric acid*, in the same manner, if there is great sensitiveness in the ulcerated parts.

If these two remedies arrest the disease, causing diminution of saliva and a cleaner appearance of the ulcers, wait patiently twenty-four or thirty-six hours, at which time one dose (four glob.) of *Sulphur* will be sufficient to complete the cure. But if the disease progresses,

*Lachesis* will be necessary to be given, in the same manner as *Bellad.*, above; particularly when the neck becomes much swollen and glossy, and the gangrene of the ulcers still continues.

If no change is perceptible from its use within twelve hours, let it be followed by *Arsenic*; which is the principal remedy, particularly when the *weakness is very great*. These two remedies may be given in alternation, in the same manner, and, for at least twenty-four hours; if then not better, let

*Nux vomica* and *Carbo veg.* follow, in the same manner.

If a drowsy sleepiness, with snoring, appears, give *Opium*, three times every hour a dose (each dose four glob.), and afterward *Cuprum* in the same manner; if the patient

sleeps with eyes half open, give *Bellad.* and *Hellebor.*, as directed under "Scarlet Fever."

After the patient is out of danger, but the ulcers do not heal fast enough, give *Nitric acid*, evening and morning a dose (four glob.), and in two or three days afterward, *Sulphur*, in the same manner.

DIET AND REGIMEN. — Diet as in "Scarlet Fever;" cleanliness and free ventilation are strongly recommended.

N. B. For the treatment of the milder form of this disease, see also the preceding article on "Sore Throat or Quinsy."

#### INFLAMMATION AND SWELLING OF THE TONSILS.

(*Amygdalitis.*)

This disease, in its *acute* form, is already treated of in the article, "Sore Throat or Quinsy," to which we refer the reader. In the *chronic* form, where the tonsils have become enlarged and remain so, often for years, without seriously endangering the life of the patient, it is extremely dangerous to have the tonsils cut out; as frequently, after such an operation, the lungs become affected, and the patient dies of consumption.

The only safe way to cure these enlarged tonsils, is to put the patient under homœopathic treatment; and, even though it takes a long time (the nature of the disease does not admit of a speedy reduction), it is still better than to expose the patient's life to the greatest danger, by an operation.

Commence such a course by giving the patient the following remedies, in their order; of each remedy every four days a dose (four glob.) for six weeks, until an amendment is perceptible:

The remedies are: *Belladonna*, *Calcarea carb.*, *Lachesis*, *Hepar sulph.*, *Ignatia*, *Sulphur*.

---

DIET.—As usual in the homœopathic treatment for chronic diseases.

SWELLING OR ELONGATION OF THE PALATE.

(*Falling of the Palate.*)

Some persons are troubled with this disease after the slightest cold or indigestion; it consists in an inflammation of the palate (*uvula*), which produces the feeling of its falling or elongation. If it continues long, it becomes very annoying, and the cause of a tickling in the throat and an irritating cough. The practice of cutting off the palate, in order to take away its inflammation, is certainly radical enough, but by no means rational, and always pernicious.

TREATMENT.—*Nux vomica* is the best remedy, in all recent attacks; evening and morning a dose (six glob.) If not better in forty-eight hours, take *Mercury*, *Bellad.*, *Calcarea carb.*, and *Sulphur*, one after the other, in the same manner, until better.

DIET.—Avoid meat, coffee, and spices; gargle the throat with cold water, and externally apply cold water bandages. (See "Sore Throat.")

## CHAPTER XIII.

### AFFECTIONS OF THE WINDPIPE AND CHEST.

#### 1. HOARSENESS. (*Raucitas, Aphonia.*)

If this complaint accompanies other diseases, such as measles, croup, influenza, etc., its treatment will be found in the respective articles; but if the hoarseness, which is an affection of the mucous membrane of the larynx, exists alone or only combined with a cough, we will state its treatment here.

Hoarseness differs in regard to time and degree. It is recent (*acute*), or of long standing (*chronic*); also, sometimes very slight, at others very severe, amounting even to a perfect *aphonia*, or loss of voice.

Catarrhs or colds in the throat frequently manifest themselves only as hoarseness, and should be treated as follows:

##### *a. Acute Hoarseness.*

*Nux vomica* and *Mercury* (see "Influenza") are the principal remedies. *Nux vomica* for a rough, dry cough, with soreness and tough mucous in the throat; the patient feels alternately chilly and hot; is impatient and morose.

*Mercury*. Either after *Nux vomica* or *Pulsatilla* (see below), or in alternation with them, particularly when with a hoarse and rough voice there is a sensation of burn-

ing tickling in the windpipe, and disposition to *easy* and *profuse perspiration*, without relief; if with sneezing, etc. (See "Influenza.")

*Capsicum*. Hoarseness with an unpleasant tickling in the nose, which feels dry and obstructed; beside, a severe cough, and pain in the head and bowels.

*Chamomile*. Particularly in children under similar symptoms as *Nux vomica*. Sometimes one cheek red and hot, the other pale and cool; fever in the evening, and great *irritability* of temper.

*Drosera*. Hoarseness with a very deep and hollow voice.

*Sambucus*. Hoarseness, with a deep, hollow cough; difficulty of breathing, which occasions profuse sweating.

*Pulsatilla*. Stinging and soreness in the throat; voice scarcely audible (aphonia), with a loose cough and pain in the chest; running of thick, yellow mucous from the nose. Suits well after *Aconite*, and is followed well by *Sulphur*.

*Sulphur*. Roughness and scraping in the throat; deep, rough voice in damp weather, with difficulty of breathing, particularly at *night*. Follows well after *Mercury*, *Pulsatilla*, *Bryonia*, and *Rhus*, and is necessary in every obstinate case of hoarseness, particularly at the end of the cure.

*Bryonia* and *Rhus*. In alternation, every two or three hours a dose (four glob.), when there is soreness of the throat and chest; worse after talking, with chilliness; pains in the limbs and head; oppressed breathing, frequent *sneezing*, with discharges from the nose. After it *Mercury* and *Sulphur* suit well.

ADMINISTRATION. — Take of the remedies indicated three doses (four glob. each) in twelve hours (every four hours one); then wait twelve hours, and if not better take the next chosen remedy in the same manner.

DIET. — No meat or coffee; drink freely of cold water, and use the cold water bandages around the throat during the night.

*b. Chronic Hoarseness.*

A hoarseness which has continued for some time is a symptom of the beginning of bronchitis, and ought to be attended to immediately. If it can not be cured by the following remedies, a homœopathic physician must be consulted.

*Hepar sulph.* If a person has been used to taking large quantities of calomel or other mercurial preparations, he may commence the cure with *Hepar*, particularly if there is a cough, worse at night, *on lying in bed*, or when *any* part of his body gets *cool*. After or in alternation with *Phosphorus*.

*Carbo veg.* Chronic hoarseness; worse after talking; in wet weather; in the evening and morning; also, after measles.

*Sulphur.* See under "Acute Hoarseness."

*Phosphorus.* Chronic hoarseness, with *roughness* in the throat; voice scarcely audible (aphonia) with dryness in the throat and chest. Follows well after *Hepar sulph.*

*Causticum.* In very obstinate cases, where the hoarseness is worse in the evening and morning; great weakness in the organs of speech, with a *sore feeling in the throat and chest* on coughing.

ADMINISTRATION.—Take of the remedy indicated twice a day one dose (four glob.) for two or three days, and then wait four or six days for its effect; if then not better, take the next remedy chosen in the same manner.

DIET AND REGIMEN.—Very little meat; no coffee or other stimulating drink, but much cold water, and every night cold water bandages around the throat. (See "Hydropathy.") In no disease is hydropathic treatment of more benefit than in the beginning of bronchitis, which is most always prevented by it. Keep the feet warm, and do not converse much.



## 2. ULCERATION OF THE WINDPIPE, commonly called BRONCHITIS. (*Phthisis laryngea*.)

This disease consists in a chronic inflammation and final ulceration of the larynx or windpipe; it is, consequently, dangerous in its results, and ought to be treated by an experienced physician as soon as one can be procured.

It manifests itself by the following symptoms:

DIAGNOSIS. — Pain in the windpipe; difficulty of breathing, and sometimes in swallowing; hoarseness more or less, with frequent attacks of severe coughing; expectoration of a very little mucous or matter, sometimes streaked with blood; finally hectic fever; chilliness; burning in the hands, soles of the feet, and cheeks; thirst during fever.

TREATMENT. — Before a physician can be obtained commence the cure with

*Aconite*. If there is fever and pain in the throat.

*Spongia* and *Hepar sulph*. In alternation, if with soreness in the throat there is difficulty of breathing; sometimes wheezing.

*Lachesis*. If there is a feeling in the throat as of something sticking in it, which can not be removed by swallowing.

*Phosphorus*, *Carbo veg.*, *Causticum*. See “Chronic Hoarseness.”

ADMINISTRATION AND DIET. See “Chronic Hoarseness.”

## 3. CROUP, AND CROUPY COUGH.

(*Angina membranacea*. *Laryngitis Catarrhalis*.)

Although these two diseases are very different in their nature, they are frequently confounded and thought to be the same, only different in degree. The real difference, however, is this:

*Croup* is a spasmodic, inflammatory disease of the wind-

pipe (larynx), and in fatal cases consists in the formation of a membrane, which closes up the windpipe. It occurs mostly in dry, cold weather, with sharp northern winds.

*Croupy cough* is a catarrhalic, inflammatory state of the windpipe, which may end, if not cured, in inflammation and congestion of the lungs. It occurs mostly in damp, wet, and cold weather. Each may precede or follow the other; but as the remedies are entirely different, it is necessary to know the symptoms which distinguish them from each other.

In the *real croup* children are mostly attacked in the fore part of the night, without much previous ailment, after an apparently sound sleep; they are seized with sudden oppression for breath, which appears in hurried, difficult, and hoarse respiration, accompanied by spells of coughing, with a squeaking, shrill, and grating, or metallic sound. These attacks waken the child, which is very restless, and in more or less agony during a spell of the cough, but after it generally sleeps sweetly, and without obstruction in breathing, at least the first evening. Not so in croupy cough, where the breathing is labored, and the cough hollow and hoarse, even during sleep. The child coughs without awaking, showing that no local pain distresses it.

Croup belongs more to a northern climate — the farther north, the more intense the croup; while croupy cough seems to take its place in the more southern countries, where real croup seldom occurs, except during long-continued storms from the north.

In severe cases of croup, the intermission of the disease on the second day, in the morning, is not sufficient to permit the child to sit up and be cheerful for play, which it does in croup of less intensity, where the hoarseness, cough, and fever abate in the morning, to such an extent as to render

the young patient quite cheerful and playful; if, in such a case, the same, or even severer symptoms appear on the second evening, do not be discouraged, but give the same medicine again which gave relief the night previous.

The mildest attack of croup, or croupy cough, is apt to return for three nights, and parents ought to be particularly careful during the first three or four days not to allow their little croup patients to come outside the room, or near the windows, where they could inhale the cold air; keep a flannel around the neck, which must be removed gradually when the patient is convalescent, by tearing a small strip from it every evening.

TREATMENT OF CROUP. — Without going into elaborate details on the treatment of croup, which would only confuse the reader, I will state briefly that course which is pursued invariably at first in every case of croup, coming under the treatment of a homœopathic physician, generally with the *greatest success*, leaving only a few *very severe cases* to be treated with other remedies, to select which requires the experience and skill of a physician.

The first remedy, under all circumstances, is *Aconite*, twelve globules dissolved in half a teacupful of water, to be given every fifteen minutes a teaspoonful until four teaspoonfuls are taken; then administer *Spongia* and *Hepar*, in alternation, each one prepared like *Aconite*, in separate cups, every half hour a teaspoonful until the child is better; when it sleeps do not disturb it to give the medicine; as soon as it is better give only every three or four hours a teaspoonful alternately. If the patient gets *very bad*, breathing very quick and labored, with a constant rattling in the breast and drowsiness (coma) of the head, prepare and give *Tartar emet.*, and *Phosphorus*, in alternation, in the same manner as *Spongia* and *Hepar*, above. If not better in three hours give *Opium* twice, every half hour a

dose (three glob.), and then *Cuprum*, in solution, as *Aconite* above, and if this does not succeed in three hours, give *Camphor* and *Coffea*, in alternation for two hours, prepared as above, and then *Lachesis*, in the same manner as *Aconite* above.

TREATMENT OF CROUPY COUGH.—Give first one dose (four glob.) of *Aconite*, and then in one hour afterward *Chamomile*, *Bryonia*, and *Belladonna*, in alternation, every two hours one dose (four glob.) until better.

DIET AND REGIMEN.—During the attack nothing but the mildest nourishment, such as arrow root, farina, milk-and-water, and water alone, if there is great thirst. Warm foot-and-hand-baths are recommended, with this precaution, that the parts thus bathed should be covered with warmed flannels directly after the bath.

#### 4. INFLUENZA.

This disease appears mostly as an epidemic, sometimes, however, sporadic, and has, in addition to the symptoms of *common cold in the head* (see this article), chilliness, followed by flashes of heat, great debility and headache, which forces one to lie down; rheumatic pains in the whole system; painful and red eyes, with great sensibility to light and difficulty of breathing.

TREATMENT.—If most of the above symptoms are present, give

*Arsenic*. Every three hours a dose (four glob.), until three doses are taken; then discontinue for six hours, awaiting its effect, and, if not better, give one of the following remedies, according to the symptoms:

*Camphor*. If the chilliness still continues, or there is great difficulty of breathing and failing of strength; let the patient either smell camphor or drop three drops of spirits of camphor in half a tumbler of water, and give every fifteen or thirty minutes a teaspoonful until better.

*Mercury.* Particularly in the commencement, when the head symptoms prevail, such as sneezing and running of the nose, with rheumatic pains in the head and body, red, watery eyes, or when a loose cough, with profuse perspiration without relief, sets in; also, when there is a slimy diarrhea, with straining like dysentery. Give it in the same manner as *Arsenic* above. This remedy is considered almost a specific in this disease. (Compare, also, "*Materia Medica*," at the end of the book.)

*Bryonia.* In alternation with *Aconite*, every two or three hours a dose (four glob.), if inflammatory symptoms appear; fever; dry, hot skin, with a hard shaking cough, and stitches in the chest.

*Belladonna* and *Aconite*, in the same manner, if the cough is dry and spasmodic, with severe headache and dryness in the mouth and throat.

*Phosphorus* and *Tartar emet.*, in alternation, in the same manner, if the lungs feel oppressed, weak, and sore; and when coughing or breathing there is rattling in the chest.

*Causticum.* If the feeling of *excoriation* in the chest is prevailing, with a dry cough, so violent that the urine escapes involuntarily; give twice a day a dose (four glob.)

*Pulsatilla.* After *Mercury*, or in alternation with it, every three or four hours a dose, if the cough is loose, and the nose discharges thick, yellow mucous; soreness of the chest in the morning, after expectoration; foul tongue; disagreeable or insipid taste in the mouth; no thirst.

Consult, also, the article on "Coughs."

DIET AND REGIMEN.—No meat or coffee; for drink, water or lemonade, if there is no diarrhea; gruels and toast-water. Keep the face from the direct influence of the fire in the grate or stove; it is preferable to remain in bed, where a gentle warmth may be maintained.

## 5. COUGH.

As cough is, in most cases, only a symptom of some other disease, such as pneumonia, bronchitis, influenza, whooping-cough, consumption, etc., we must refer the reader to their respective chapters. Sometimes, however, a cough appears isolated from any marked disease, and is either caused by cold (*catarrhal cough*), or sympathy, or derangement of other organs (*nervous, spasmodic, and gastric cough*); so that we have to mention here their appropriate treatment.

*a. Catarrhal cough*, caused by a severe cold on the lungs, is, in the beginning of the day, harsh and painful; afterward it becomes loose, with heavy expectoration, relieving the chest from pain and oppression.

TREATMENT.—If such a cold is combined with pains in the head and limbs, chilliness, and thirst (commonly called *bilious*), give *Bryonia* and *Rhus tox* alternately, every two or three hours a dose (four glob.), until better, or until three doses of each are taken; then wait twelve or sixteen hours before the next remedy is taken which will suit the remaining symptoms of the cough.

When the cough is *dry*, without expectoration: *Aconite*, *Chamomile*, *Bryonia*, *Rhus tox.*, *Belladonna*, *Hyosциamus*, *Capsicum*, *Ignatia*, *Nux vomica*, *Phosphorus*, *Hepar*.

When the cough is *loose*: *Pulsatilla*, *Mercury*, *Tartar emetic*, *Phosphorus*, *Dulcamara*, *Stannum*, *Lycopodium*, *Sulphur*.

*b. Nervous and spasmodic cough*, caused by nervous irritation or excitement, either original, or transferred to the lungs from other organs; for instance, pain in the womb (*neuralgia*) may cease suddenly, and, by being transferred to the lungs, creates here the *hysterical cough*. (See article "Hysteria," especially.) All the nervous coughs are



dry; but they may affect the stomach, and cause *vomiting* or *retching*; sometimes *spasms* in the lungs may take place; *fits of suffocation*, which are generally of not so much danger as they appear to be.

TREATMENT.—For *nervous* and *spasmodic cough*: *Bel-lad.*, *Hyosciamus*, *Ipecac.*, *Bryonia*, *Drosera*, *Hepar*, *Cina*, *Sulphur*.

If accompanied by *vomiting*, *retching*: *Ipecac*, *Mercury*, *Bryonia*, *Tartar emetic*, *Pulsatilla*, *Sepia*.

If by *fits of suffocation*: *Ipecac*, *Tartar emetic*, *Opium*, *Chamomile*, *Bryonia*, *Drosera*, *Hepar*, *Lachesis*, *Sulphur*, *Arsenic*, *China*.

*Cough*, caused by *sympathy* from other diseases, such as *worms*, *dentition* (*teething*), *dyspepsia*; *growing too fast* causes in young persons such coughs. Look for these in their respective chapters.

N. B. After having found the kind of cough in the above, select the best remedy from those designated for such kind of cough, comparing their symptoms, as follows:

SYMPTOMATIC DETAIL.—*Aconite*. *Violent, short cough*; sometimes with pains and anxious oppressions in the chest.

*Chamomile*. Dry cough, from constant tickling in the throat and chest; worse after talking *in the night*; with fever and coughing during sleep; particularly in children, when they cough after crying, or a fit of anger or passion. (Compare “*Croupy Cough*,” “*Cold in the Head*,” and “*Hoarseness*.”)

*Bryonia*. Dry cough, from irritation in the throat, or with pain in the chest; worse in the open air; sometimes when from taking cold in damp, frosty weather; with rheumatic pains in the head and limbs; also chilliness, followed by fever and thirst (in alternation with *Rhus tox*); or a loose cough, with yellowish expectoration, or spasmodic

cough after eating or drinking. (For children, see "Croupy Cough.")

*Rhus tox.* Short, dry cough, from a tickling in the chest; worse in the evening, or before midnight; with weakness in the breast, and shortness of breath; restlessness at night; dry tongue (see *Bryonia*); or cough, with a taste of blood in the mouth; bloody saliva runs from the mouth *during the sleep*.

*Belladonna.* Short, dry cough, particularly at night; worse when moving; with dryness and tickling in the throat, and redness of the face, *dry, spasmodic cough*, with retching, *mostly after midnight*; *dry cough* day and night, with soreness of the abdominal muscles, or in the region of the hip, as of falling of the womb. (For children, see "Croupy Cough.")

*Hyosciamus.* Suits well when *Bellad.* has afforded only partial relief, or when the cough at night is temporarily mitigated by sitting up in bed; much rattling in the throat and tickling or dry hacking cough, with weak respiration; worse by muscular action or motion; the patient can hardly walk up stairs. (See "Consumption Beginning or Galloping.")

*Capsicum.* Particularly in persons of *phlegmatical* temperament; easily *chilled after drinking cold water*; cough *worse toward evening and night*; with pains in various parts of the body; bursting headache; painful pressure, and aching in the throat and ears; cough, with offensive breath and disagreeable taste in the mouth.

*Ignatia.* Dry cough, with running at the nose day and night, in the same degree; or short, hacking cough, as if from a stricture in the throat, or the tickling of a feather; aggravated by the continuation of the cough; particularly in persons who are easily affected by grief, or when the

cough is worse after *eating*, or on *lying down at night*, or on *rising in the morning*.

*Nux vomica* is one of the most useful remedies for various kinds of cough from cold. (See "Influenza," "Cold in the Head," etc.) *Tickling cough*; worse *early in the morning*; sometimes with *catarrhal hoarseness*, and scraping around the palate; tough mucous in the throat and lungs, with an itching, tickling sensation in the throat, and chilliness; cough worse by movement, reading, or meditation after meals; and followed sometimes by retching or vomiting; or *fatiguing cough*, with pains in the head, as if it would burst; or pains in the pit of the stomach, as if from a *blow* or *bruise*; worse in the night, or after lying down; cough dry during the after part of the day and night, with oppression on the chest, as if from a weight; expectoration toward morning. Suits well for persons of an energetic, sanguine, choleric temperament, and for those who drink much coffee and ardent spirits.

*Phosphorus*. Dry cough, from tickling in the throat, but more in the chest; worse when laughing, *talking*, and drinking; hoarseness, and pains in the chest, particularly on the *left side*, as if from *excoriation*; *lying on the left side excites the cough*.

*Hepar sulph*. In alternation with *Phosphorus*, in dry, hoarse cough; worse after talking; or excited when *any* part of the body becomes *cool*, particularly at night.

*Tartar emetic*. Loose cough, *with much rattling* on the chest; quick breathing, almost amounting to suffocation; hot and moist hands, with perspiration on the forehead; bluish lips (in alternation with *Phosphorus*, see "Inflammation of the Lungs"); or cough, with *vomiting of food* after eating; *deep, hollow cough*; *expectoration of mucous in the night*.

*Dulcamara*. Loose cough, after taking cold; spitting of

light red blood ; hooping cough, excited by drawing a deep breath ; worse in a room, or when lying still ; better when moving and out of doors.

*Stannum.* Hard, dry cough in the *evening* ; excited by laughing, talking, and singing ; also, a loose cough, with much expectoration of mucous or sweetish matter, with feeling of soreness of the chest after it (see “Consumption”) ; lying on the *right* side excites the dry cough.

*Lycopodium.* Cough worse in the night, or after drinking ; or a tickling cough, excited by drawing a *deep breath*, with a saltish expectoration, and stitches in the *left* side of the chest (see “Consumption.”)

*Sulphur.* Dry cough, worse in the *evening* and *during the night*, or when lying down ; loose cough, with expectoration of thick mucous, and feeling of soreness in the chest (particularly suitable after *Mercury*) ; cough, with pain and stitches in the chest or the head ; also, a sensation of spasmodic constriction in the chest, as if caused by the vapors of brimstone.

*Ipecac.* Nervous or spasmodic cough, sometimes with nausea, retching, and vomiting ; worse at night ; accompanied by oppression in breathing, almost amounting to suffocation, as if *from the accumulation of mucus (congestion to the lungs)*, particularly in children ; when the face looks livid and dark, and the body becomes rigid and stiff. (See “Hooping-Cough.”)

*Drosera.* Dry, *spasmodic cough*, with *retching* ; worse immediately after *lying down*, in the *night* ; chronic cough, and hoarseness after *measles* ; dryness, and a rough, *scrapping* sensation, deep in the throat, which causes coughing. (See “Bronchitis,” and “Hooping-Cough.”)

*Cina.* Dry, *spasmodic cough*, with shortness of breath, pale face and moaning, also in the night, with restlessness and crying. (See “Worms,” and “Hooping-Cough.”)

*Pulsatilla.* Severe shaking, dry cough, mostly in the morning, with retching and inclination to vomit; or loose cough, with much expectoration of a bitter, yellow mucus.

*Mercury.* Cough, with hoarseness (see "Influenza"), or excited by tickling in the throat; shaking, dry cough, mostly at night, or loose cough, with expectoration streaked with blood, and an inclination to perspire easily, but without relief.

*China.* Severe cough, from ulceration of the lungs (see "Galloping Consumption"), after hemorrhage, or other losses of blood; or asthmatic cough in the night, with acute pain in the chest and shoulder-blades.

*Arsenic.* When *China* only ameliorates, without curing; or when there is dry cough after drinking; or cough, with bloody expectoration in the night, with a burning sensation over the whole body; asthmatical cough (see "Asthma.")

*Opium.* Cough when swallowing or breathing, with anxious, heavy, and intermitting respiration; full, red face (see "Congestion of the Lungs.")

*Sepia.* Dry cough in the evening in bed, frequently with sickness at the stomach and bitter vomiting; or loose cough, with much saltish expectoration, particularly in the morning and evening (see "Consumption.")

*Lachesis.* Short, dry, suffocating cough, as if from something sticking in the throat, with ineffectual efforts to expectorate.

ADMINISTRATION.—Of a remedy having been carefully selected, dissolve twelve globules in half a teacupful of water, and let the patient take every two, three, or six hours a teaspoonful (children half a teaspoonful), until three or four doses have been taken; then wait from four to twelve hours, according to the severity of the case; and if not better, select and give the next best remedy in the same manner. This will suit for recent attacks. In coughs of

a longer standing, give the remedy for two days, evening and morning a dose (four glob.) ; then discontinue two or three days ; if then not better, take another remedy in the same manner. To very small children, give the medicine in globules altogether, if they can not take it dissolved in water ; one or two globules at a time is a dose for them.

DIET AND REGIMEN.— If the cough is accompanied by fever or biliousness, abstain from meat and butter ; if not, these may be used : otherwise, keep the diet prescribed in chronic diseases. If persons are habitually liable to take cold, they ought to harden themselves against the changes in the atmosphere, by the systematic use of cold water, externally and internally, as spoken of in the latter part of the book (see “Hydrotherapy”) ; persons, however, with tubercular affections must not use it. Another great dietetic rule is, that persons liable to colds on the lungs ought to abstain totally from *all* spiritous liquors.

The bathing of children in cold water, when they have cold on the lungs, may be suspended for the time (except when allowed by a physician especially) ; as in many cases it might not be admissible at all.

A patient with a cough should not occupy a damp room, facing to the north ; but a dry, airy chamber, exposed to the midday sun, and susceptible of easy ventilation and even temperature. If the cough at any time is very dry and painful, use, as a loosening drink, an infusion of the roots of althea and licorice, in equal parts, as freely as the patient finds it necessary, cold or warm. In chronic coughs, where the expectoration is tinged with blood, no warm drinks are allowed. In fall and winter, until summer, a plaster of Burgundy pitch on the breast is recommended to those who have weak breasts ; but, after taking it off in the summer, they must wash and rub the breast well.



HOOPING-COUGH. (*Tussis Convulsiva.*)

This is one of those diseases which exclusively befalls children, from the infant on the breast up to twelve or fourteen years. It appears mostly as an epidemic, and is, as such, infectious. In some seasons its attacks are milder than in others : sometimes it commences immediately with hooping ; at other times, it is preceded by all the symptoms of an inflammation of the lungs, with great tendency to convulsive and spasmodic affections. The real hooping seldom commences in the earliest stage.

DIAGNOSIS.—We recognise, generally, two stages for this disease :

The *first*, or *febrile stage*, showing all the symptoms of a heavy cold ; running at the nose ; watery, heavy eyes ; sneezing ; irritating and painful cough, with feverishness and difficult respiration. It is in this stage that congestions to the head and lungs often occur, particularly when children are teething at the time, which, of course, adds materially to the danger of the case. The old practice is almost powerless in such cases, while homœopathy offers a safe and successful treatment.

The *second*, or *nervous spasmodic stage*, begins as soon as the feverishness diminishes, and the hooping commences ; the latter increases by degrees, until the paroxysms become, at least in severe cases, really appalling.

The little patients dread the attack, and run, as it were, to the mother for help, or hold fast to something, when they feel the approach of it. In severe paroxysms, the face swells and becomes livid, while quantities of mucus are forced from the mouth, and frequently blood escapes from mouth and nose ; respiration seems to cease for a minute, when a deep inspiration ensues, which ends the attack for the time, and the patient recovers quickly ; being, in

the intervals, almost always perfectly well, except, perhaps, a little fretful and weak.

**DURATION.** — If this disease is not treated at all, or only by allopathic medicines, its duration is left to its own natural limits, and not shortened in the least; in such cases, the first stage may last from three to eight, or even twenty-one days, and the second stage, from the beginning to its height, from five to six weeks; from thence to its total disappearance, the same length of time. Frequently, the cough reappears in those patients in the fall or winter, who had it in the preceding summer.

But under the proper homœopathic treatment, its duration is very much shortened; sometimes the disease is cut off in its progress within five or six days. Very seldom has a person the whooping-cough twice in his lifetime, except as above stated.

**TREATMENT.** — *In the first stage*, where the children apparently have a cold, select from those remedies recommended in the articles on “Cough,” and “Croupy Cough,” by the use of which the cough frequently entirely disappears. These remedies are mostly *Chamomile*, *Belladonna*, *Bryonia*, etc.

*Aconite.* If the cough does not cease in a few days, but becomes very dry and whistling; the child has fever, and complains of a burning pain in the windpipe (*administration*, see “Croup”), until better, or some of the following remedies are indicated.

*Dulcamara.* If the cough is loose and moist, with copious expectoration of a thin mucus, evening and morning a dose (three or four glob.) for two days.

*Pulsatilla.* In the same manner, if there is a *loose cough*, with vomiting of mucus or food; also, a slimy diarrhea.

*Ipecac* and *Nux vomica*, alternately, if the cough is dry, accompanied by great anguish, strangling, and bluish face;

worse after midnight, until morning; *with vomiting* and bleeding at the nose. Give alternately every hour or two hours a dose (three glob.) until better.

*Tartar emetic* and *Phosphorus*, in alternation, in the same manner, is frequently necessary in this period, if there is rattling in the lungs, short breathing, great debility, drowsiness, and thirst; also, retching, with diarrhea.

*Carbo veg.*, is the principal remedy, when the cough exhibits the first signs of hooping; particularly, when it is worse in the *evening*, the patient complaining at the same time of sore throat when swallowing; shooting pains in head and chest; especially useful, when the weather is damp and cold. Give it the same as *Dulcamara*, above.

*In the second stage*, when the hooping or spasms during coughing have really commenced, the principal medicines are: *Drosera*, *Veratrum*, *Cuprum*, *Tartar emetic*, *Phosphorus*, *Cina*, *Bryonia*, *Bellad.*, *Mercury*, *Iodium*, *Opium*, *Hepar sulph.*, *Arnica*, *Sulphur*, *Hellebor*.

*Drosera*. After the *Carbo veg.*, when the paroxysms become more violent, with real hooping, vomiting of food and mucus, without or with fever, which is characterized by chilliness, heat with thirst, hot perspiration in the night; the patient feels better when moving about.

*Veratrum*. After *Drosera*, or in alternation with it, for similar symptoms, only accompanied with great weakness; small and quick pulse; slow fever, with cold perspiration on the forehead; child is unable to hold up its head; miliary eruption (prickly heat) over the whole body; apathy and drowsiness, from weakness; child dislikes to move or speak.

*Cina*. If the child, during the hooping-cough, exhibits symptoms of worms—as picking at the nose, sudden pains or gripings in the bowels, or itching at the anus—or if it becomes *stiff* during a paroxysm of cough, after which a

gurgling noise is sometimes heard, descending from the throat into the stomach. This remedy can be given in alternation with *Mercury* (see "*Mercury*"), if there is bleeding from the nose and mouth at the same time.

*Cuprum*. If the paroxysms appear *very often* during twenty-four hours, rendering the little patients *rigid* and *unconscious*; if with drowsiness and rattling of mucus in the chest between the paroxysms; in alternation with *Tartar emetic* (see "*Tartar emetic*," compare, also, "Congestion to the Head.") This remedy is particularly useful, when convulsions appear instead of the cough, and cease when the paroxysms return; as if the paroxysm of the lungs had been temporarily transferred to the brain. (See "Convulsions.")

*Tartar emetic* and *Phosphorus*. See page 227.

*Belladonna*. Before or after *Cuprum*, when the brain becomes affected; patient can not bear the light, noise, or motion; head seems to ache; burning fever; thirst; restlessness; delirium; convulsions (see "Inflammation of the Brain"); in such cases, in alternation with *Hellebor.*; paroxysms terminate in sneezing.

*Hellebor*. At any time during the hooping-cough, when symptoms of congestion to the head appear, particularly when the patient sleeps with the eyes half open (see *Belladonna*, above); it is of great service when children are teething, or otherwise weakly and delicate.

*Bryonia*. Sometimes in alternation with *Hellebor.*, when *Belladonna* was insufficient, for similar symptoms; or when the paroxysms of suffocating cough appear more in the evening or at night; mostly, however, after eating or drinking, which is vomited up.

*Mercury*. The principal remedy, when the child *bleeds profusely* at the nose and mouth when vomiting, with co-

pious sweat at night, and great nervousness; also, if there are worm symptoms (see "*Cina*." )

*Sulphur*. At any period, if the paroxysms are accompanied by vomiting, which will not yield to other remedies.

*Iodium*. If the patient pines away, or has a chronic, watery diarrhea, of lightish color.

*Opium*. If the best indicated remedies seem not to have the desired effect, or if there is stupor (see "Congestion to the Head"), with hot perspiration; *irregular breathing*, with great anguish; constipation.

*Hepar sulph*. When the cough is better, but yet hollow, hoarse, and dry, with retching and *crying* after a paroxysm.

*Arnica*. When the child cries much after coughing, or coughs after crying (in alternation with *Cina*<sup>cc</sup>.)

ADMINISTRATION.—Whenever no reference is given to other chapters, give the remedies as follows: Dissolve twelve globules in half a teacupful of water, and give for two days a teaspoonful, morning, noon, and night; then wait two or three days for the effect; if the cough is slightly improved, wait still longer; because the whooping-cough needs, besides the right medicine, a little time, to disappear by degrees; but, if no better, give the next remedy, in the same manner.

DIET AND REGIMEN.—In cases with fever, see the diet in "Croupy Cough;" where there is no fever, give the usual diet, without coffee and spices. A great deal of fresh air is good for the young patients, if they have no feverish symptoms. The same holds good in relation to their usual bathing, which must be discontinued, if fever is present.

#### INFLAMMATION OF THE LUNGS. (*Pneumonia*.)

This disease generally commences with chills, followed by fever, with difficulty of breathing, and a short, dry cough; pulse soft in the beginning, afterward hard, but

always very quick ; dull pains in the chest at every deep respiration, with great oppression ; expectoration of a tough, lumpy mucus, afterward mixed with blood ; the patient generally prefers to lie on his back ; is not inclined to speak or when speaking, pauses after every articulation ; is sometimes sullen, and wishes to be quiet ; in some cases, the face turns to a bluish *purple*, with red cheeks ; tongue dry, parched ; great thirst ; skin dry, hot. As soon as the skin becomes moist and natural, the oppression in breathing diminishes, and a thick mucus is coughed up abundantly ; the danger is over, and the patient needs only good nursing and rest in bed during convalescence. If this is not the case, however, the inflammation runs either into the nervous or typhoid stage (see "Typhoid Fever"), or in *hepatization* of the lungs (the air-cells are filled with lymphatic matter) ; both very dangerous and critical situations for the patient.

*Causes of Pneumonia* are, mostly, exposures to cold, north or northeast winds ; particularly when at the time one is over-excited by fast running, etc. ; also, suppression of hemorrhages or habitual secretions ; inhalation of obnoxious gases ; irritating treatment of catarrh (I have seen inflammation of the lungs follow the use of hoarhound candy, in a simple cold, and whooping-cough.) Pneumonia appears in its highest inflammatory degree in the north, while toward the south it gradually decreases in violence, but not in danger ; because here the bilious complication occurs more frequently (*bilious pneumonia*), where bleeding is *particularly injurious*.

TREATMENT. — This disease requires quick and energetic action ; and the first remedy to be given, under almost all circumstances, is

*Aconite*. In the chill, as well as the highest fever, in the commencement (either alone, in water dissolved, twelve



globules to half a teacupful), every half hour a teaspoonful, until the fever is broken, and the pain and oppression in the chest is ameliorated; or in alternation with

*Bryonia*, when oppression and pain in the chest is very severe (resembling pleurisy—see that article); worse by every movement or cough; expectoration bloody, of a brickdust color; also, pains in extremities; tongue is coated, often dry, with great thirst; constipation.

Administer the same as *Aconite*, only every hour or two hours a teaspoonful.

If after these two most important remedies the disease still seems unchecked, give, in six or eight hours afterward, if not better,

*Phosphorus*, in alternation with *Tartar emetic*, dissolved in water, the same as *Aconite* and *Bryonia*, every hour, or two hours, a teaspoonful, until three or four teaspoonfuls of each are taken; wait for the effect of these remedies from eight to twelve hours, as frequently their effects are not seen immediately. If necessary, repeat both the above prescriptions, or choose among the following remedies, if the patient, in place of being better, has run into the *typhoid* or *hepatized* stage of the pneumonia.

The *typhoid* stage of pneumonia commences when the patient is restless, particularly at night; throws himself about and becomes quite delirious, or lies motionless on his back in a stupor; breathing quick and irregular; snoring with his eyes half closed, and rattling of mucus in his lungs; tongue dry, pulse quick and thread-like; skin dry, or covered with clammy sweat (see “Typhus Fever.”)

TREATMENT.—In the beginning of this state give

*China*. Particularly if the patient, in the early part of the disease, under allopathic advice, had been bled, or lost blood by other means, such as hemorrhage, or a fall or blow (in the latter case, in alternation with *Arnica*); or

when the patient complains mostly of a *pressure* in the chest, stitches in the *breast* and *sides*, when breathing and coughing, together with palpitation of the heart; besides, great weakness, fine, quick pulse and yellowish coated tongue. Administer two or three times, every two or three hours a dose (four glob.); then wait from four to six hours, after which recourse may be had to *Bryonia* and *Rhus tox*, as prescribed in "Typhus Fever" (see this article.) If the disease still progresses, it will be necessary to give

*Opium* (see this remedy in "Typhus Fever"), or *Arnica*, under similar symptoms, but without delirium, or *Hyoscyamus* (see "Typhus Fever"), if the cough is spasmodic or very irritating.

*Belladonna*. (See "Typhus Fever") when the oppression is very great.

*Veratrum*. If the pulse is very small and weak; extremities cold; delirium; vomiting and diarrhea; sinking; no sleep whatever.

*Camphor* and *Coffea*, in alternate doses, in a similar state, particularly if the patient is covered with cold perspiration.

*Phosphoric acid*. If, after the patient has been relieved by the above remedies, there is yet diarrhea present. (See "Typhus Fever.")

*Sulphur* may be given once or twice, if the amelioration brought about by previous remedies seems not to be permanent; then return to those medicines again, which previously were the most efficacious.

Administration the same as in "Typhus Fever."

*Hepatisation*, or the *second* stage in pneumonia, when infiltration of the lungs with coagulated lymph takes place, is indicated by greater difficulty of breathing, and a dull sound by percussion of those parts of the breast where the pain was the severest.

TREATMENT. — *Sulphur* has first to be given (twelve

glob. dissolved in half a teacupful of water, every two or three hours a teaspoonful) four times; then wait from eight to twelve hours, and, if not better, give

*Lachesis*, *Lycopodium*, *Phosphorus*, one after the other, in the same manner as *Sulphur*, until better.

*Arsenic* and *China*, in alternation, in the same manner, when fetid expectoration, of a dirty, green color, appears.

DIET AND REGIMEN.—As in fevers generally; but particularly as in “Typhus.”

*After-Diseases*.—If pneumonia threatens to become chronic, or consumption ensues, see this article; when the patient expectorates clear matter, *China*, *Mercury*, *Sulphur*, *Stannum*, *Lachesis*, *Hepar*, *Lycopodium*, are the principal medicines to be given, as stated in “Consumption.”

*Pneumonia in infants and children* up to two or three years of age is very frequent; in fact, every cold on their lungs affects them more or less in an inflammatory manner, and ought not to be neglected for a moment.

Besides heat, fever, thirst, dry mouth, there is *quick, oppressive breathing*, with *short cough*, after which the child always cries; these latter symptoms are a *sure sign* of inflammation of the lungs in children, and unless they are relieved or disappear (particularly the respiration is slower and regular), we must not feel secure in regard to the safety of the young patient.

TREATMENT.—The principal remedy is

*Aconite*. In solution or dry; half a teaspoonful, or two glob., every two or three hours, three or four times, or until better; after which, if necessary, give

*Bryonia*. In the same manner, if the cough is very bad; or

*Belladonna*. In the same manner, if the head is very hot at the time. If these remedies do not relieve in from twelve to twenty-four hours, give

*Tartar emetic* and *Phosphorus*, in alternation (ten glob. of each, dissolved in two teacups, each half full of water), every hour, alternately, half a teaspoonful, for twelve or twenty-four hours, until better ; discontinue for the same length of time, and repeat afterward, if necessary ; if, during that time, the child is very sleepless and restless, give *Coffea* <sup>cc.</sup>, and *Belladonna* <sup>cc.</sup>, in alternation, a dose (three glob.) every hour, until better. Continue with this treatment for three or four days, and in the most severe cases it will prove successful.

If convulsions, or congestions to the head occur, see these articles, and treat accordingly.

DIET AND REGIMEN. — Keep the patient as quiet as possible ; in a darkened chamber, if the head suffers ; not too warm ; if the child does not take the breast, give cold water freely, if thirsty ; of solid food, crackers soaked in milk or water ; neither broth nor meat ; gruels are good.

*Pneumonia in Old People. (Pneumonia notha.)*

This disease shows itself in old people as follows : They become at once prostrated, having symptoms of a common cold, with a cough, chills, and fever ; the cough is loose, and the expectoration white, yellow, slimy, and streaked with blood ; the respiration quick and labored ; pain in the breast, when drawing a long breath, mostly on a small spot ; fever is not high ; pulse soft but quick ; cheeks slightly flushed ; skin generally moist at night, without relief ; during the day, the patient feels tolerably comfortable ; the voice, however, is weak, almost in a whisper. As these symptoms do not *seem* to indicate any danger, the disease is frequently left to itself, but very often with the most fatal results ; because paralysis of the lungs soon ensues, and death is inevitable.

TREATMENT. — *Aconite*. Every three hours a dose (six

glob.) will be first necessary, three or four times, during the fever, after which give

*Mercury*. In the same manner; wait twelve hours and then, if not better, in alternation with

*Belladonna*. Particularly when the cough becomes short and dry, and the breathing difficult and suffocating; if not better, give

*Sulphur* once or twice; afterwards repeat *Mercury* and *Belladonna*.

*Veratrum* and *Arsenic*. If they sink very low; cold extremities, etc. (See "Typhoid Pneumonia.")

*Tartar emet.* and *Phosphorus*. If, with constant rattling in the lungs, paralysis threatens; alternately every half hour a dose (three or four glob.)

Compare, also, the remedies under "Pneumonia."

DIET AND REGIMEN.—The same as in "Pneumonia," only more nourishing as soon as the inflammatory symptoms have left in the least; the yolk of an egg with sugar; broth of beef with rice, etc. After-diseases, see "Consumption."

There is another disease, called *galloping consumption*, in which the inflammation of the lungs forms the most important part. This species of pneumonia occurs mostly in young people of florid complexion, at the time of development, and needs a separate article. (See "Galloping Consumption.")

It is evident, from the nature of pneumonia and the dignity of the organ affected, that a good homœopathic physician ought to be consulted, if he can be procured; but if this can not be done rely on the above rules and prescriptions.

#### REMARKS ON PNEUMONIA.

This disease may frequently be complicated with other diseases, such as inflammation of the pleura, heart, and

brain; also, with bilious, rheumatic, or gastric symptoms; the treatment, however, does not vary from the above; beside, consult the respective articles.

PLEURISY. (*Pleuritis.*)

- a. *Real Pleurisy.* (*Pleuritis serosa.*)
- b. *False Pleurisy.* (*Pleuritis muscularis.*)

a. *Real Pleurisy.*

If the sac around the lungs (called pleura) becomes inflamed, we call it *real pleurisy*; this can be the case with or without inflammation of the lungs, at the same time. If complicated with pneumonia or bronchitis, these articles must be consulted together with this chapter.

DIAGNOSIS. — In real pleurisy the pain is preceded mostly by a chill and fever, as in pneumonia, but it does not produce so much general oppression in breathing as pneumonia; however, in pleurisy the pain is more severe on a fixed spot (mostly in one of the sides of the chest), from which it radiates; increased often to intolerance by deep inspiration or hard pressure between the ribs, where the pain is located; the breathing is, on this account, short, and the lying on the affected side impossible; if the back part of the pleura towards the spine is inflamed, the pains increase by moving the spine, and running up into the neck impede frequently the swallowing of food, with inclination to vomit, or produce real vomiting.

Fever is never wanting, but not so evident as in pneumonia; neither is the cough so distressing, but more a short and dry one, very seldom with much expectoration of mucus.

This disease terminates frequently in the formation of lymphatic effusion, which result may be known by the following change of symptoms, different from the above



real pleuritic ones: Respiration becomes more anxious and short; the patient can not now lie on the well side either, or, if he does, he feels in danger of suffocation, because the water presses over toward the well lung; by turning, which is very difficult, he feels something moving within the chest, like water; the fever becomes hectic; the urine is offensive, and has a sediment.

This description will be sufficient to distinguish pleurisy from pneumonia; but there is another form of pleurisy, the so-called

*b. False Pleurisy. (Stitch in the Side.)*

Which resembles the true or real pleurisy very much, yet is different in origin and location, as well as in termination.

DIAGNOSIS.—This false or spurious pleurisy is not an inflammatory, but a rheumatic affection of the intercostal muscles, and occurs mostly in persons subject to rheumatism; it is preceded generally by pains in the neck, shoulders, and throat, as if from cold, without much of a chill, fever, or thirst; the pain in the side is worse by *exhaling*; in real pleurisy, worse by *inhaling*; *slight* pressure on the affected side increases the pain in real pleurisy, while only *hard* pressure with the fingers between the ribs increases it in false pleurisy, which has a *slight* cough only; in real pleurisy the cough is *dry* and *painful*.

TREATMENT. *a. Of Real Pleurisy.*—*Aconite* and *Bryonia*. First of all, in the same manner as described in “Pneumonia,” for so long a time as the pain is very intense; if not better in thirty-six or forty hours, give

*Sulphur*. Twelve globules dissolved in half a teacupful of water, every two or three hours a teaspoonful, until four or six are taken, particularly if the fever continues with the painful oppression on the chest, yet the skin has

become a little moist; this is the principal remedy, also, when effusion of plastic lymph is feared, or has taken place; it is well enough to give the patient even a dose (six glob.) of *Sulphur*, if *Aconite* or *Bryonia*, or both, have restored him so far as to be able to be out of bed. If *Aconite* ameliorates the pains, do not be in haste to give *Bryonia* immediately, but wait until the disease increases again, and the same in regard to *Bryonia*. In general, cease to give another remedy as soon as the former has shown a decided beneficial effect.

If the patient is sleepless and restless at night, give at any time *Coffea* <sup>cc.</sup>, and *Belladonna* <sup>cc.</sup>, in alternation, every hour a dose (four glob.) until better.

If the patient complains of laying hard on his side, in consequence of which he changes his position often; has cold hands and feet, yet feels hot inwardly; or if a short, dry cough troubles him very much, particularly by increasing the stitching pain in his side, give *Arnica* once or twice, every two hours a dose (four glob.)

If the real pleurisy is complicated with pneumonia, (*pleuro-pneumonia*), see and give the remedies described under "Pneumonia," such as, *Belladonna*, *Rhus tox.*, *Lachesis*, *Sulphur*, *Phosphorus*, *Lycopodium*, *China*, *Sepia*,\* according to the symptoms indicating their use.

N. B. As these pleuro-pneumonias are frequently of a *bilious* character, the remedies for *bilious* or *bilious rheumatic* fever must be consulted.

TREATMENT. *b. Of False Pleurisy, (Stitch in the Side.)*  
If there is any fever, give first once or twice a dose (four

\* NOTE. — *Sepia* once cured an old man of a most violent attack of *pleuro-pneumonia* on the left side, caused by being transferred from the liver, where it had been subdued, but not cured, a few days previous, by excessive allopathic blood-letting; *China* had preceded the *Sepia*, without decided effect.

glob.) of *Aconite*, every hour and a half one dose ; after which administer the principal remedy :

*Arnica*. Every two hours a dose (six glob.), and in alternation with

*Pulsatilla*. If the patient is not better in two hours ; but if he is relieved, scarcely any thing more is needed, but rest and perspiration in bed. In two or three days afterward one dose of *Sulphur* (four glob.) may be given to complete the cure.

If at any time the patient becomes feverish, with pains in limbs, side, and head, give *Bryonia* and *Rhus tox.* (See "Rheumatic Bilious Fever.")

DIET AND REGIMEN.—As in fever generally. In false pleurisy frequently a mustard draught, or some other heating substance—for instance, a bag of heated salt—will relieve, and may be used with advantage.

#### DISEASES OF THE HEART.

##### *a. Inflammation of the Heart. (Carditis.)*

It is well that inflammation of the heart does not occur frequently, as its course is very rapid and dangerous, and its diagnosis difficult.

DIAGNOSIS.—The prominent symptoms are *violent* pains, mostly *burning* or *cutting*, in the region of the heart and toward the pit of the stomach ; great oppression in breathing, which shows itself in the distorted and anxious features of the patient ; the pulsation of the heart is *violent* and *irregular*, while the pulse on the wrist is small, frequent, and trembling ; the patient prefers lying *quietly on the back* ; sometimes there is delirium, difficulty of swallowing, fainting spells, vomiting, starting in the sleep.

CAUSES.—This disease is caused mostly by taking cold in drinking cold water when the body was heated ; also,

by wounds, blows, etc.; the most frequent cause, however, is acute rheumatism, particularly in the knees. See "Rheumatism of the Heart."

It must be well understood that not every palpitation of the heart is inflammation of the heart; the latter has always an agonizing feeling of oppression and pain.

TREATMENT AND ADMINISTRATION. — In every case give first *Aconite*, as directed in "Pneumonia;" afterward *Bryonia*, *Nux vomica*, *Cocculus*, *Arsenic*, *Pulsatilla*, and *Cannabis sativa*, at intervals of from two to four or six hours, until better; each remedy, when used, may be dissolved in water (twelve globules to half a teacupful), and every half hour a teaspoonful given; but, if possible, procure the advice of a homœopathic physician *immediately*.

*b. Palpitation of the Heart. (Palpitatio cordis.)*

Young people in their growing years, as well as old persons of high age, are troubled with this disease; the former from developement and congestion, the latter from contraction, ossification, and want of blood. Between these two extremes of age, persons sometimes are taken with it by violent emotion of the mind, by the use of ardent spirits, or by constitutional predisposition, particularly when inclined to be affected by rheumatism, which latter mostly causes the chronic palpitation of the heart. In pregnancy this disease is frequent, as, also, in persons of a plethoric habit, and at the time of change of life, or during floodings, in consequence of loss of blood.

If it is dependent on organic disorders of the heart, as polypus, ossification, etc.—as, also, in the chronic form—we advise strongly to apply in time to a homœopathic physician, as frequently a permanent cure may be effected.

TREATMENT. — If caused by *fright* — *Opium* or *Coffea*.  
By *sudden joy* — *Coffea*.

By *fear or anguish* — *Veratrum*.

By *disappointment* — *Aconite*, *Chamomile*, *Ignatia*, *Nux vomica*.

By *congestion of blood, or plethora* — *Aconite*, *Belladonna*, *Coffea*, *Ferrum*, *Lachesis*, *Aurum*, *Phosphorus*, *Opium*, *Sulphur*.

By *loss of blood, or other debilitating losses* — *China*, *Phosphoric acid*, *Nux vomica*, *Veratrum*.

If it is present in *nervous persons, hysterical females, etc.*  
— *Coffea*, *Ignatia*, *Chamomile*, *Cocculus*, *Nux vomica*, *Lachesis*, *Pulsatilla*, *Veratrum*.

In *young, growing people* — *Aconite*, *Pulsatilla*.

In *old, decrepit people* — *Arsenic*, *Lachesis*, particularly when with temporary dimness of sight.

After the suppression of an eruption — *Arsenic*, *Lachesis*, *Sulphur*.

For ameliorating the frequent attacks in *chronic palpitation* of the heart, the most suitable medicines are, *Pulsatilla*, *Arsenic*, *Lachesis*, *Aconite*, *Sulphur*, *Phosphorus*, *Aurum*.

ADMINISTRATION. — Dissolve of the selected remedy twelve globules in half a teacupful of water, and take from one to twelve hours a teaspoonful, according to the severity of the case; or, if traveling, take of the medicine a dose (four glob.) from one to six or twelve hours; if not better after from one to twelve hours, select another remedy.

DIET AND REGIMEN. — Diet as usual in chronic diseases. Sleep as quiet as possible during the administration of the medicine; be careful always to have dry, warm feet; avoid getting wet by exposure to rains, etc. Young people must abstain from eating much in the evening, but drink freely of cold water.

CONGESTION OF THE CHEST. (*Plethora pectoris*.)

*Congestion or determination of blood* to the chest is a frequent complaint in young people, during the time of the development of their systems, and in persons of a phthisical habit. It ought never to be treated by blood-letting, even if this means relieves for the time, as it does not remove the disposition or the bad consequences in the system, but has a tendency, if once resorted to for palliation, to augment the difficulties for a final cure, rendering the lungs weaker by each successive attack. The homœopathic remedies, on the contrary, relieve in a different manner, and in a much shorter time.

**DIAGNOSIS.** — Great fullness, weight, and pressure in the chest; more or less palpitation of the heart, with oppression, short and sighing breathing; cold hands and feet, and anxious look; sometimes a hacking, short cough.

**TREATMENT.** — *Aconite*. Particularly in plethoric females during menstruation; *violent oppression*, with palpitation of the heart; shortness of breath; dry cough; heat and thirst.

*Belladonna*. If *Aconite* does not suffice, and if the head is congested at the same time. See “Congestion to the Head.”

*Nux vomica*. Heat and burning in the chest, more at night, with sleeplessness and agitation; or fullness and throbbing in the chest, with palpitation; worse in the open air; clothes feel too tight around the chest.

*China*. In consequence of debilitating losses, as of blood, etc.; *violent oppression*, with great anguish; loss of sight; palpitation of the heart; can not breathe with the head low.

*Phosphorus*. Violent oppression, with heaviness, fullness, and tension in the chest; palpitation of the heart, which is felt in the throat.



*Mercury.* After *Aconite*, if there is anxious oppression for breath, with a desire to take a long breath; heat and burning in the chest; cough, with expectoration of mucus streaked with blood.

*Pulsatilla* and *Bryonia*. In alternation, before menstruation, or when this has stopped suddenly; also, when the piles have been suppressed; or when there is ebullition of blood in the chest, with external heat.

*Spongia.* When movement increases the anguish, and produces danger of suffocation, nausea, fainting, and prostration; particularly in persons subject to sick-headache, or other nervous affections of the head, which frequently affect the chest in this manner.

*Sulphur.* After *Nux vomica* or *Pulsatilla*, if necessary.

ADMINISTRATION, DIET, AND REGIMEN, same as in "Palpitation of the Heart."

#### HEMORRHAGE OF THE LUNGS. (*Hæmoptysis*, *Pneumorrhagia*.)

*Spitting* or *coughing* up of blood is generally considered by persons such a dangerous and, for the life of the patient, fatal symptom, that they lose, when it occurs, all presence of mind, frequently incapacitating them for the right action in the case; yet not all the hemorrhages of the lungs are alike in their character; some are dangerous, it is true, but most of them yield readily to the proper remedies, and some even cease of themselves. It will be necessary to distinguish well between them, and act accordingly, not in too great haste, but in the right manner; in this way, time is saved. Moreover, we will see hereafter that the most dangerous kinds of bleedings from the lungs leave ample time to provide means, while the milder forms take more by surprise, thereby causing more fear than injury, with the exception of one form, the so-called "Apoplexy of the Lungs," which is easily dis-

tinguished by marked symptoms. Be careful to examine whether the blood coughed or hawked up is from the lungs, or merely from the nose or palate.

DIAGNOSIS. — Without going into the distinction as to the different places in the lungs and windpipe from whence the blood may issue, as this would be without practical advantage, we will give now the qualitative difference of hæmoptysis, which has the highest practical bearing. We distinguish two varieties :

1. The *active* or *idiopathic* hæmoptysis, caused by congestion of the lungs, or mechanical injury ; this is the most frequent and least dangerous.

2. The *passive* or *symptomatic* hæmoptysis, caused by the ulcerative process in consumption, which gradually consumes the substance of the lungs (*if not arrested*), and causes the rupture of bloodvessels ; although this is the most dangerous form, as it is caused by a destructive and sometimes incurable disease, yet it does not occur frequently, and even then its appearance may be expected and measures taken accordingly.

First, *Active hemorrhage of the lungs* depends always on congestion of blood to the lungs, which is sometimes constitutional, sometimes caused by external circumstances, such as quick running, violent exercise of any kind, lifting, etc., the frequent inhaling of injurious dust, such as from lime, gypsum, or plaster of Paris, tobacco, flour, or metal filings, or of obnoxious gases, such as from nitric and sulphuric acid.

Besides these causes, it sometimes appears epidemically, mostly at the time of the equinoxes, on account of the rapid changes in the temperature of the atmosphere ; the abuse of spiritous liquors is a frequent cause of hemorrhage of the lungs ; also, suppression of discharges of blood from other organs, as stoppage of the menses, sud-

den disappearance of the piles, etc.; excessive growing and consequent weakness of the chest predisposes to it, as in general young or middle-aged persons are more liable to it. It occurs in attacks, accompanied with more or less fever, frequently of an intermittent type, worse toward night.

This form of hemorrhage of the lungs presents itself in the most various degrees, from the least spitting of blood to a violent effusion; the highest degree, which really endangers life immediately, is the *apoplectic hemorrhage* (apoplexia pulmon.) of the lungs, where all the symptoms of an apoplectic fit are present; the patient loses his consciousness, looks as if he was suffocated, eyes protrude out of their sockets, bloody mucus issues from the mouth.

TREATMENT. — The first thing necessary in severe cases is to let the patient keep as quiet as possible, in a half-sitting, half-lying posture; he must remain perfectly quiet, without speaking or being spoken to, except what is indispensably necessary; if no homœopathic medicine is at hand, give every five or ten minutes a teaspoonful of table salt in water, but cautiously, that it may not make him cough; or, if *sulphuric acid* is convenient to be had, drop five or ten drops into a tumbler full of water, and give a teaspoonful every five or ten minutes, until the severest bleeding ceases. If homœopathic medicine is at hand, give directly

*Aconite*. Twelve globules dissolved in half a teacupful of water, every ten or twenty minutes a teaspoonful, until the severest bleeding ceases; afterward administer it at longer intervals, from three to six hours. This remedy suits always at first, but particularly when there is ebullition of blood in the chest, with a fullness and a burning pain; palpitation of the heart; anxious looks, with paleness of the face; profuse expectoration of blood at intervals,

provoked by a slight cough. Try this remedy at least for two hours ; if not better, give

*Ipecac.* When there yet remains a taste of blood in the mouth ; frequent, short cough ; expectoration of mucus, streaked with blood ; *nausea* and weakness ; or,

*Arsenic.* When there is yet palpitation of the heart, with great anguish, dry heat, and desire to leave the bed ; it is particularly applicable for drunkards after *Hyoscyamus* has been given.

*Opium.* Often in the most serious cases, especially in persons addicted to spiritous liquors ; or when there is expectoration of thick, frothy blood ; cough worse after swallowing ; oppression and anguish ; weak voice, drowsiness, and anxious starts ; cold extremities, and heat in the chest.

*Nux vomica.* Suits well after *Opium*, *Ipecac.*, or *Arsenic* ; particularly when there is cough, affecting the head, caused by tickling in the chest ; or when the hemorrhage was caused by a fit of passion, by taking cold, or suppression of piles. *Sulphur* suits well after it.

*China.* If the patient has already lost much blood ; or when the cough is violent, hollow, dry, and painful, with taste of blood in the mouth ; shivering and flushes of heat ; weakness, faintness, cloudiness of sight, roaring in the head.

*Ferrum.* After *China*, if this has in severe cases relieved, yet there is great fatigue after talking ; slight cough, with expectoration of scanty, bright red blood, with pain between the shoulder-blades ; difficulty of breathing.

*Hyoscyamus.* Dry cough at night, with expectoration of blood ; frequent waking with a start ; particularly in drunkards, after *Opium* and *Nux vomica*, and followed by *Arsenic*.

*Dulcamara.* Tickling cough, from cold ; the hemorrhage was preceded for some time by a loose cough.

*Carbo veg.* Hemorrhage, with violent *burning* pain in the chest, in persons susceptible to changes of weather, or who have taken much calomel.

*Pulsatilla.* In obstinate cases, when *black* and *coagulated* blood is expectorated, more in the morning or night, particularly in timid persons, or in females when caused by suppression of the menses.

*Arnica.* In slight hemorrhages of *black* and coagulated blood, with stitching, burning pains and heat in the chest, palpitation of the heart, difficulty of breathing; or caused by *mechanical injuries, fall, blow, lifting*, leaning against a table by studying, sewing; or when the expectoration is clear, frothy, mixed with pure lumps of matter; hacking cough, from tickling under the breast-bone; pains in the limbs, as if they had been beaten. It is often administered in alternation with *Aconite*.

*Belladonna.* Cough from tickling in the throat, with hemorrhage; sensation of fullness, as if from blood in the chest, with pressing or shooting pains, worse when moving; is especially suitable for hemorrhages of the lungs in females of full habit, at the change of life, and then in alternation with *Aconite*.

In very serious cases, *Aconite, China, Ipecac.,* and *Opium* will be found most efficacious.

For the bad consequences of pulmonary hemorrhage, such as weakness, cough, etc., give *Carbo veg.* and *China*. If possible, procure the advice of a physician immediately.

ADMINISTRATION of the above remedies, in very severe cases, the same as in *Aconite*, stated above; in less severe cases, or mere spitting of blood, give every six or twelve hours a dose (four glob.) until better, or until another remedy is necessary.

DIET AND REGIMEN. — The patient must be kept cool; his drinks, for two weeks, at least, must be cool, not spiritous or

exciting, but more slimy, such as rice and barley water, water with raspberry syrup, lemonade, etc. If the feet are cold, put them in warm water, mixed with ashes or mustard.

In *apoplectic* hemorrhage of the lungs, give *Aconite* immediately, same as above; if not better in twenty or thirty minutes, *Opium*, and then *Ipecac.*; afterward, if necessary, *Belladonna* or *Lachesis*, in the same manner. Put the feet in hot water, with mustard; also the hands, if they are cold. Send immediately for medical aid.

Second. *Passive* hemorrhage of the lungs, or rupture of a bloodvessel in tubercular consumption, requires principally *Aconite*, *China*, *Pulsatilla*, *Arsenic*, *Lycopodium*, to be given in the same manner as described under active hemorrhage. See, also, "Pulmonary Consumption."

DIET AND REGIMEN the same as above.

#### ASTHMA.

As this disease is well known in regard to its symptoms, we will only mention here that distinction in its character which has a practical bearing. There are two kinds of asthma, as regards the cough and expectoration of mucus in an attack; it is either a *dry* or *humid*. The former is more sudden and violent in its attacks—has, however, slight cough and expectoration, but great difficulty of breathing. The attack of the latter, or humid asthma, proceeds slowly—has a severe cough, with gradually increasing and afterward copious expectoration of mucus, which relieves greatly. Asthma, in itself, is not a dangerous, but a very distressing disease, sometimes hereditary, but oftener acquired; its different causes will be enumerated below, with their respective remedies attached. In general, it is curable, although it takes a comparatively longer time than most other chronic diseases require, par-



ticularly when it is constitutional. A radical cure must only be undertaken by a skillful homœopathic physician. We subjoin below the treatment for the attack itself, to alleviate temporarily, advising the patient, at the same time, not to omit any favorable opportunity by which he may be allowed to place himself under the care of a physician, for the final cure of the disease.

TREATMENT.—When caused by *congestion of blood* to the chest (see this article): *Aconite, Belladonna, Nux vomica, Phosphorus, Pulsatilla, Spongia, Sulphur.*

By *derangement of the menstruation*: *Belladonna, Cocculus, Bryonia, Pulsatilla, Veratrum, Sulphur.*

By *vapors of Sulphur* (brimstone): *Pulsatilla.*

By *vapors of Arsenic* or *copper*: *Ipecac., Hepar, Mercury.*

By a *chill*: *Aconite, Ipecac., Bryonia, Belladonna.*

By a *fit of passion*: *Nux vomica, Chamomile.*

By *fright*: *Opium, Ignatia, Aconite, Belladonna, Veratrum.*

By *sudden joy*: *Coffea, Aconite.*

By a *suppressed catarrh*, or *cold in the head*: *Ipecac., Nux vomica, Arsenic.*

By *dust from stone or flour*: *Calcarea, Silicea, Sulphur.*

By *suppression of piles*: *Nux vomica, Sulphur, Arsenic.*

By *flatulency*: *China, Chamomile, Nux vomica, Sulphur.*

If it is present in *children*: *Aconite, Belladonna, Ipecac., Chamomile, Bryonia, Tartar emet., Sambucus.*

In *hysterical women*: *Coffea, Ignatia, Pulsatilla, Ipecac., Chamomile, Belladonna.*

In *aged persons*: *Arsenic, Lachesis, Opium.*

When it is *constitutional*: *Ipecac., Arsenic, Sulphur.*

When *dry*, or without much expectoration (similar to *cramps* in the chest): *Cocculus, Cuprum, Belladonna, Nux vomica, Sambucus.*

When *humid* or *moist*, with considerable expectoration of mucus: *Ipecac.*, *Arsenic*, *China*, *Pulsatilla*, *Sulphur*.

In the above tabular view, we have arranged the different kinds of asthma, as regards their origin and appearance; so that, in search for a remedy, the number of remedies to be looked for is reduced to very few. Their distinguishing symptoms are given in detail below, and their administration at the end of the article. The principal and first remedy in most all attacks is *Ipecac.*, after the administration of which one may proceed with more calmness to find out the next best remedy, by comparing carefully its symptoms with those of the case present.

SYMPTOMATIC DETAIL.—*Ipecac.* Difficulty of breathing; nocturnal paroxysms of suffocation, spasmodic constriction of the throat, rattling in the chest, from an accumulation of mucus; short, dry cough, great anguish, cries, agitation, fear of death; alternately either redness and heat, or paleness, coldness, and ghastliness of the face; sometimes nausea, with cold perspiration on the forehead; breathing at the same time *anxiously* and *rapidly*, with moaning and rattling. After it follows well *Arsenic*, *Bryonia*, or *Nux vomica*.

*Arsenic.* In all cases of acute or chronic asthma, with an accumulation of thick phlegm; difficult breathing after a meal; oppression of the chest and want of breath, rendered almost intolerable by any motion, such as walking, going up stairs, laughing, etc.; constriction of the chest and throat; *suffocating fits*, more at *night*; panting, wheezing, as if *dying*, with *cold perspiration*; these paroxysms grow lighter on the appearance of a cough, with expectoration of mucus, or of viscid saliva; the attacks render the patient *very weak*; burning pain in the chest. Suits well after *Ipecac.*, and before *Sulphur*.

*Nux vomica.* Asthma, resembling that of *Arsenic*, only the spasmodic constriction is felt worse *in the lower part of*

*the chest*, where even loose clothes seem to be too tight ; *short, dry cough*, sometimes with a little blood ; congestion of the chest (*see this article*) ; asthma, better by *lying on the back* ; occasionally turning, or sitting up ; suits well for persons addicted to the use of spiritous liquors, or after *Arsenic*, or before *Sulphur*.

*Pulsatilla*. Asthma in children, after suppression of a miliary eruption ; in weak, timid females, after cessation of menstruation, or after having taken cold ; *choking, as from the vapors of Sulphur* ; *paroxysm of suffocation, with deadly anguish, palpitation of the heart*, and spasmodic constriction of the throat or chest ; worse at night, or when lying horizontally, or when moving ; better in the open air ; expectoration, with much mucus, streaked with blood, which sometimes is coagulated ; with a feeling of fullness and pressure of the chest.

*China*. Under similar symptoms as *Pulsatilla*, after which it suits well, if there is, with the asthma, *great weakness*, and *easy perspiration in sleep*.

*Sambucus*. Asthma, similar to *China*, but more in children (*see "Asthma in Children"*) ; when there is *much perspiration* ; suits well after *Ipecac*.

*Sulphur*. Almost in any acute or chronic attack of asthma, if several other remedies were insufficient. After *Sulphur*, if necessary, select carefully another remedy. It suits principally when there are paroxysms of asthma, almost *suffocating*, mostly at *night*, with *fullness and weariness, burning, or spasms* in the chest ; expectoration of white mucus, detached with difficulty, sometimes bloody ; constriction and pain in the breast-bone ; bluish redness of the face ; short respiration, and inability to speak.

*Phosphorus*. Asthma worse in the evening or during *movement* ; attack of *suffocation* in the night, as if from

paralysis of the lungs (in alternation with *Tartar emetic*); congestion of blood to the chest, with stitches, and fullness or sensation of heat to the throat, and palpitation of the heart (in alternation with *Belladonna*); phthisical habit.

*Tartar emetic.* Asthma in *old persons* and *children*, particularly when there is *choking*, *retching*, and paroxysms of *suffocation* in the evening, with *rattling* in the chest, and wheezing (in alternation with *Phosphorus*; see "Inflammation of the Lungs.")

*Aconite.* Asthma in sensitive persons, young, and of plethoric habit, easily excited by mental emotions. In children, especially, when they cough *at night*, *spasmodically*, as if they would *suffocate*; or with congestion to the head, with vertigo; cough, with expectoration of blood.

*Belladonna.* Asthma in children (see "Croupy Cough"), or in *plethoric women* at the critical period; congestion to the lungs, with stitches under the breast-bone, and fullness in the chest; *dry cough at night*; anxious moaning; respiration, sometimes *deep*, at other times *short* and *rapid*; constriction of the throat, as if to suffocate, with loss of consciousness and relaxation of all the muscles.

*Bryonia.* Asthma, worse by *motion* and in the *night*, with pain in chest; breathing like *Belladonna*.

*Coffea.* Asthmatical breathing from sudden joy; short, dry cough.

*Opium.* Suffocating cough, with bluish-red face; deep, rattling, respiration, especially when fright was the cause.

*Chamomile.* Asthma in children, after taking cold (see "Croupy Cough"), or when the hypochondriacal region is swollen, painful to the touch; crying, and drawing up of the thighs; asthma caused *by passion*.

*Cocculus.* Asthma of hysterical women, especially when they complain of constriction of the throat and chest, with

oppression ; worse at night ; congestion of the lungs in nervous persons (in alternation with *Belladonna*.)

*Ignatia*. Asthma after fright or indignation, especially in women ; choking and constriction in the pit of the neck, as if from the vapors of *Sulphur* ; *want of air in walking*, and *cough when resting* ; short, spasmodic cough.

*Spongia*. Want of breath, and paroxysms of suffocation after every movement, with fatigue ; rush of blood to the head and chest ; heat in the face ; respiration wheezing, deep, or slow, as if from weakness (*nervous asthma* in persons who are habitually addicted to neuralgic affections.)

*Cuprum*. In hysterical women, after fright or anger, or before or during the menses ; *spasms* and *oppression* in the chest ; worse on speaking or going up stairs ; short, spasmodic cough.

*Veratrum*. Paroxysm of suffocation when rising up, during movement, especially before the menses ; coldness of the face and extremities ; faint pulse and cold perspiration. Suits well after *Ipecac.*, *Arsenic*, or *China*.

*Lachesis*. Slow and wheezing, or short respiration ; worse after eating ; paroxysms of *suffocation when lying down*, especially in persons suffering from water on the chest. (Suits well after *Arsenic*.)

*Calcareæ carb.* In chronic asthma, with *frequent dry cough* ; worse *at night* ; or caused by the habitual inhalation of dust.

*Silicea*. In chronic asthma, caused by the inhalation of stone-dust ; oppression worse when lying on the back, when running, or coughing.

ADMINISTRATION. — Dissolve twelve glob. of the selected remedy in half a teacupful of water, and give, in severe cases, every half hour ; in less severe cases, every two or three hours, and in chronic cases, twice a day a teaspoonful, until three or four are given. If better, discontinue

until the symptoms are worse again; if not better, prepare and give the next remedy in the same manner. In very small children, give one or two glob. of the medicine at the same intervals, dry, on the tongue.

DIET AND REGIMEN.—Persons suffering from chronic asthma, ought to abstain from coffee, meat, and greasy substances almost altogether, and live mostly on farinaceous diet, such as gruels, etc.

Sometimes, in very severe attacks, the burning of a piece of paper, soaked in a solution of saltpetre, and filling the chamber of the patient with its smoke, will relieve; or smoking tobacco, or gypsum-weed (*stramonium*.)

*Asthma of Millar.*

See “Children’s Diseases.”

DROPSY IN THE CHEST. (*Hydrothorax*.)

This disease appears mostly in consequence of previous diseases, either acute or chronic, and is too dangerous in its course and results to be managed by domestic treatment alone. Skillful medical aid is indispensable; yet we will briefly notice its symptoms, and some few remedies which mitigate its immediate distresses.

DIAGNOSIS.—We distinguish an acute and chronic dropsy of the chest; the principal distinguishing feature of both kinds is, the difficulty of breathing, when moving, particularly going up stairs, or lying with the head low; swelling of the feet in the evening, and of the eyelids in the morning; thirst for water, yet scanty urine.

The *acute dropsy* of the chest occurs mostly in young people after eruptive fevers, such as scarlet fever, measles (see *these diseases*), or after the suppression of other eruptions by salves, ointments, etc.



The *chronic dropsy* of the chest occurs mostly in old age, or in persons addicted to the habitual use of ardent spirits.

TREATMENT. — To alleviate the difficulty of breathing, and render the patient more comfortable, give him *Arsenic* (twelve glob. dissolved in half a teacupful of water), every hour a teaspoonful, until the judicious advice of a homœopathic physician can be obtained.

#### PULMONARY CONSUMPTION. (*Phthisis Pulmonalis.*)

We distinguish several forms of this disease—the most frightful destroyer of human life and happiness; but, as its treatment would be too complicated for a work of this kind, we would advise our readers to apply, in time, to a homœopathic physician; as consumption, in its beginning, is in most cases curable.

DIAGNOSIS. — If some of the following symptoms appear, we advise to seek medical aid: constant hacking cough, either dry, or with an expectoration of a frothy mucus; shortness of breath, pain in the chest, derangement of the stomach, hectic fever, chilliness followed by flashes of heat, burning in the hands and soles of the feet, night-sweats; circumscribed redness of the cheeks during the fever, but pale cheeks at other times. The so-called *tubercular* consumption consists in the softening of the small tubercles, which are in the lungs frequently for years, without disturbing health, when they remain in their compact state; their softening, however, produces irritation and ulceration of the lungs. In the beginning, this process can be repressed, and the patient saved.

The *galloping* consumption is peculiar to young persons, in their growing years, or after debilitating diseases, and requires the most careful attention of a physician. It is very rapid, and is rightly called *galloping*; as it is most acute, sometimes even infectious, on that account. *Aconite*, *Hyos-*

*ciamus*, are remedies, frequently useful in this species of consumption, to allay the high fever and restlessness, and particularly for the dry, short, irritating cough and shortness of breath, with great weakness in the night.

Under *scrofulous* consumption, we understand an ulceration of the lungs, caused by the transfer of scrofulous affections from other parts of the system, such as swellings of the glands of the neck, etc., and all other more specific diseases, such as syphilis, cancer, etc.; also, when eruptive fevers fall with their fatal decision on the lungs, and there produce ulceration.

As this subject is so extensive and important to all, we refer our readers to a separate work, entitled "Consumption treated homœopathically," by A. C. Becker, M. D.

#### CONTUSION OF THE CHEST BY A FALL OR BLOW.

When the chest, externally or internally, is injured by a fall or blow, apply tincture of *Arnica* externally, as directed under "*Sprains*," and internally give *Aconite* and *Arnica* alternately, in solution, as directed under "*Hemorrhage of the Lungs*," until the patient is better; if, after two or three days, the fever still continues, with an ulcerative pain in the chest, give *Pulsatilla*; and in alternation with *Mercury*, if there is a heavy, thick, yellow mucus expectorated; the latter remedy, however, in exchange with *China*, if night-sweats occur, and derangement of the stomach, bad appetite still continues, with sleeplessness and fever.

DIET AND REGIMEN. — The same as under "External Injuries."

## CHAPTER XIV.

### AFFECTIONS OF THE STOMACH AND BOWELS.

#### 1. WANT OF APPETITE. (*Anorexia.*)

Most diseases, particularly those of the bowels and those of an inflammatory character, produce loss of appetite, which, in such cases, is more beneficial than otherwise; as it prevents the suffering system from being burdened with nourishment, at a time when the organs for its digestion and use are out of order. This applies to most of the acute, and to a great many of the chronic diseases. When these have disappeared by the use of proper remedies, the appetite returns of itself. But sometimes persons complain of loss of appetite without any apparent cause; in which case, it mostly proceeds from a derangement of the nerves of the stomach. For such patients, it is of no permanent advantage to use stimulating drinks or food, such as bitters, mustard, or coffee. These things only create an artificial appetite, never removing the cause of the disease. The best remedy, which is applicable in almost all cases of loss of appetite, is *cold water*, freely drank at any time in the day and evening; in large quantities, before, during, and after meals, etc., with the precaution, not to be overheated at the time. Besides this dietetic remedy, as it were, the following medicines are recommended: to be used according to directions given in "*Dyspepsia.*" These remedies are,

*Nux vomica, Chamomile, China, Antimon., Ipecac., Bryonia, Lachesis, Hepar, Arnica, Sulphur.*

## 2. MORBID APPETITE (*Bulimy.*)

A morbid, craving, or voracious appetite, manifests itself in consequence of another disease present in the system, such as worm affections, dyspepsia, etc.; or it frequently is the necessary result of severe loss of strength, after violent, debilitating diseases. In every case, it consists in a weakness and derangement of the nerves of the stomach, and is treated in the respective diseases, of which it is generally a concomitant symptom.

We will here only indicate some of these affections, with the remedies attached, which arrangement will be particularly useful. For their application and administration, see the respective chapters of these diseases.

If there exists a craving for food in

*Worm affections: Hyosciamus, Cina, Mercury, Silicea, Spigelia.*

*Pregnant women: Nux vomica, Sepia, Petroleum, Natrum muriaticum.*

*Convalescence after violent diseases, or loss of fluids: China, Veratrum.*

*Dyspepsia, where unnatural hunger is present: China, Veratrum, Nux vomica, Sulphur, Bryonia, Ignatia, Mercury, Pulsatilla, Lachesis, Lycopodium.*

In satisfying an unnatural appetite, care ought to be taken not to overload the stomach. Frequent, but moderate meals, are preferable.

## 3. DYSPEPSIA, OR WEAK STOMACH.

DIAGNOSIS.—Dyspepsia is generally characterized by weakness of digestion, with loss of appetite, slight or irregular appetite, painful and disordered stomach, flatulency,

sour risings, ill humor, drowsiness after a meal, sometimes vomiting of acid, or mucus, secreted in great quantities in the digestive organs.

CAUSES. — The causes of this disease are as numerous as the symptoms vary, under which it shows itself in different constitutions and periods of life. A principal cause lies in the irregularity of the diet; eating too much, too rich, or too indigestible food; using stimulating drinks, tea, coffee, liquors, to an excess, and at an unreasonable time; eating too quickly, or fasting too long between meals; studying or laboring mentally too intensely, or after a full meal—this is particularly the case in our country, where merchants are in the habit of confining themselves to their arduous business too closely after full dinners, when a few hours, complete rest, bodily and mentally, would be of the greatest benefit to them; lastly, the immoderate use of tobacco, especially in the way of chewing, the use of such deleterious medicines as calomel, etc., by which weak stomach, or dyspepsia, is produced.

TREATMENT. — Avoiding the above-named causes of this disease is the first indispensable requisite in its successful cure; at the same time, the following remedies will be of the greatest benefit, to be applied according to the symptoms.

The principal remedies for recent dyspepsia are *Nux vomica* and *Pulsatilla*; for chronic dyspepsia, *Hepar*, *Calcarea*, *Sulphur*. (See their symptoms below.)

To facilitate the choice of the remedies, we classify the disease according to its appearance and causes, with the remedies attached to each kind; before selecting a remedy, consult its details below.

If dyspepsia is present in *children*: *Ipecac.*, *Bryonia*, *Calcarea*, *Nux vomica*, *Sulphur*.

In *old persons*: *Antimon.*, *Nux vomica*, *China*, *Carbo veg.*

In *hypochondrical persons* : *Nux vomica*, *Sulphur*.

In *hysterical females* : *Ignatia*, *Pulsatilla*, *Sepia*.

If the *dyspepsia* is caused by *sedentary* or *confined* habits : *Nux vomica*, *Sulphur*, *Sepia*.

By *prolonged watching* : *Nux vomica*, *Arnica*, *Pulsatilla*, *Veratrum*, *Carbo veg.*

By *excessive study* : *Nux vomica*, *Sulphur*, *Lachesis*, *Pulsatilla*, *Calcarea*.

By *debilitating losses*, such as *bleeding*, *purgings*, *vomiting*, etc. : *China*, *Carbo veg.*, *Lachesis*, *Nux vomica*, *Sulphur*, *Calcarea*.

By *sexual excesses* : *Phosphoric acid*, *Nux vomica*, *Mercury*, *Staphysag.*

By the *abuse of ardent spirits* : *Nux vomica*, *Sulphur*, *Lachesis*, *Arsenic*, *Carbo veg.*

Of *coffee* : *Nux vomica*, *Ignatia*.

Of *tea* : *Thuja*.

Of *tobacco* : *Nux vomica*, *Hepar*, *Cocculus*, *Staphysag.*

By *over-eating* : *China*, *Pulsatilla*, *Antimon.*

By *distressing emotion*, such as *grief*, *anger*, etc. : *Chamomile*, *Nux vomica*, *Phosphoric acid*, *China*, *Staphysag.*

By *mechanical injuries*, *blow*, *fall on the stomach*, etc. : *Arnica*, *Bryonia*, *Rhus tox.*, *Sulphur*.

If the *dyspepsia* is worse after *partaking of almost any thing* : *Nux vomica*, *Sulphur*, *Lachesis*, *Carbo veg.*, *China*.

After *drinking cold water* : *Arsenic*, *China*, *Pulsatilla*, *Veratrum*.

After *drinking beer* : *Arsenic*, *Calcarea*, *Rhus tox.*, *Sepia*, *Sulphur*.

After *drinking milk* : *Bryonia*, *Nux vomica*, *Calcarea*, *Sulphur*.

After *using bread* : *Nux vomica*, *Sulphur*, *Pulsatilla*, *Bryonia*, *Mercury*.



After using *acids* : *Nux vomica*, *Sepia*, *Sulphur*, *Arsenic*, *Lachesis*.

After partaking of *meat* : *China*, *Sulphur*.

After partaking of *fat substances* : *Pulsatilla*, *Sulphur*, *China*, *Carbo veg.*

If the dyspepsia is combined with *diarrhea* : *Pulsatilla*, *China*, *Phosphoric acid*, *Carbo veg.*, *Veratrum*, *Arsenic*, *Mercury*.

With *constipation* : *Nux vomica*, *Sulphur*, *Bryonia*, *Lachesis*.

With *sour stomach* : *Pulsatilla*, *Sepia*, *Phosphoric acid*, *Nux vomica*, *Calcareo*.

With *headache* : See this article.

With *piles* : *Nux vomica*, *Sulphur*, *Sepia*.

REMEDIES ACCORDING TO THEIR SYMPTOMS. — *Nux vomica*.

In most cases of dyspepsia at the commencement, particularly in persons with a tendency to piles (see this article) and habitual costiveness ; or when dissipation and late hours, the abuse of ardent spirits, or exposure to cold, was the immediate cause of this disease ; this remedy is especially indicated for *sour or bitter taste in the mouth*, and if *food*, particularly *bread*, tastes sour, bitter, or insipid ; patient has not much appetite, but a *craving for beer*, wine, or spirits, or he has great hunger, but a little satisfies him ; after eating he is troubled with *nausea*, *eructation*, *vomiting of food*, *vertigo*, hypochondriacal humor, or, after a meal, lassitude, sleepiness, drowsiness, distension, fullness and tension in the stomach, tender to the touch ; around the waist a *sensation of tightness of the clothes* ; sour risings and belchings ; *water-brash* ; sour stomach ; makes the patient unfit for intellectual labor ; with constipation, and *frequent heat and redness of the face*, *restlessness*, and *irascibility*. After it, *Sulphur* suits well.

*Sulphur.* Is in cases of long standing the principal remedy after *Nux vomica*, or when there is no appetite for meat and bread, but a craving for acids or wine; milk, acids, and sweet substances disagree; after a meal, difficulty of breathing, nausea, pain in the stomach, belching or vomiting of food, shivering, acidity and water-brash, with flatulency and constipation; sad, morose humor, with irascibility. After it *Calcarea carb.* is often suitable.

*Pulsatilla.* In recent cases of dyspepsia, particularly those caused by over-eating, or the use of fat or pork, or any greasy substance, causing flatulency and indigestion; (in such cases in alternation with *China*); no thirst; bitter or sour eructations; the taste of the food comes up again, water-brash, sadness, melancholy, inclined to loose bowels. *China* or *Sulphur* suit well after it.

*China.* Dyspepsia from loss of blood or other humors, or caused by exposure to miasms in marshy countries, near canals, etc. (see "Fever and Ague"); particularly if there is indifference to food and drink, as if from satiety; craving wine or acids; food tastes acid or bitter; after a meal uneasiness, drowsiness, fullness, distention; patient wants to lie down, sensitive to draughts of wind, ill-natured and dislikes everything. After it suits *Bryonia*, *Rhus tox.*, *Carbo veg.*

*Bryonia.* Dyspepsia especially in summer or damp weather; or when there is headache, chilliness, pains in limbs, arms, and small of the back, aversion to food after a meal; eructation of wind, pressure and pain in the stomach, can not bear tight clothes; vomiting of food, constipation, with restlessness and irascibility. In alternation with *Rhus* or *China*.

*Rhus tox.* Always in alternation with *Bryonia*, (see "Bilious Rheumatic Fever"), especially when there is great restlessness, thirst, and dry tongue during the night.

*Carbo veg.* Bitter taste in the mouth, aversion to food, milk, or fat, with acidity in the mouth; *frequent flatulency* and belching up, tasting the fat and food which had been eaten; *nausea in the morning*, water-brash during the night; heart-burn; hiccough during motion; pains under the short-ribs, particularly on the left side, in the form of stitches, with pressing and fullness; can not bear the clothes around him tight; worse in damp weather; *offensive diarrhea*.

*Ipecac.* Almost the same as *Carbo veg.*, but especially suitable for children after indigestion, when they have vomiting of food, drink, mucus or bile; retching, vomiting, with coldness of the face and extremities; sinking at the stomach. (See "*Cholera Morbus*.")

*Veratrum.* After *Ipecac.*, if this should be insufficient, or there is still diarrhea, attended with griping in the bowels; great thirst.

*Antimon. crud.* After *Ipecac.*, if this has not removed the nausea and vomiting, with a sensation of great fullness in the stomach, and much flatulency.

*Ignatia* and *Staphysag.* See "*Diseases of the Mind*."

*Hepar.* One of the most useful remedies in this disease, where a person has taken much calomel, and is easily taken with indigestion, with a longing for stimulating things, wine, or acids; nausea in the morning, with eructation or vomiting of sour, bilious, or mucus substances; hard, light-colored, difficult evacuations, or else whitish diarrhea. (In this case alternate with *Nux vomica*; otherwise with *Mercury* or *Lachesis*.)

*Calcarea.* Sensation of coldness in the head, with inclination to sick-headache; *acid stomach*; *water-brash*; *fullness and swelling in the region of the stomach*, with great *tenderness to touch*; evacuations similar to those of *Hepar*,

but the diarrhea in children has a sour smell. Follows well after *Sulphur*.

*Sepia*. In persons subject to habitual sick-headache, who frequently suffer from dyspepsia when their heads feel comfortable; aversion to food, or else great craving; taste sour, and after a meal acidity in the mouth, and swelling of the abdomen, or else pressure, as if from a stone, in the stomach, with much belching, mostly sour or painful; inclined to constipation; suits well for nervous, hysterical persons, or those who are subject to congestions of the head or bowels.

*Arnica*. Frequent eructations, sometimes with a taste of rotten eggs; great sensitiveness and nervous excitement; tongue dry, thick, yellowish coated; putrid, sour taste, hypochondriacal humor; inclined to diarrhea.

*Phosphoric acid*. Great mental and physical debility, caused by grief, unrequited love, or loss of bodily strength by losing too much and too constantly animal liquid. In diarrhea, alternate with

*China*. After every meal pressure in the stomach, as if from a load; perception of the taste of food, particularly of bread, long after it is taken.

*Lachesis*. Irregular appetite; repugnance to bread, with craving for wine and milk; frequent nausea and eructation, vomiting of food; uneasiness, indolence, heaviness, fullness and pains in the stomach after every meal; constipation. Suits well before or after *Mercury*.

*Mercury*. Putrid, sweetish or bitter taste in the morning; repugnance to solid food and meat, with craving for cooling things; inclined to diarrhea, with straining, or to perspiration; has much saliva in the mouth.

*Cocculus*. Sensation of emptiness and hollowness in the stomach; sour taste, with aversion to acids; dryness of the throat; nausea, even to fainting; constipation.

*Phosphorus.* Vomiting after a meal; burning in the stomach; acidity and sour taste in the mouth.

*Thuja.* Weakness of the stomach after the abuse of tea; rancid or bitter eructation, and throwing up of that which has been eaten.

ADMINISTRATION. — In severe and painful cases of dyspepsia take of the selected remedy from one to six hours a dose (four glob.), until better; but in chronic cases take only every evening or every other evening a dose (six glob.) until four doses are taken; then discontinue four days, and, if not better, take the next medicine in the same manner.

DIET AND REGIMEN. — Besides avoiding those excesses in the diet and regimen which we have mentioned in the “Diagnosis,” as so many causes of this disease, it must be borne in mind that every thing even allowed under other circumstances must not be indulged in, as soon as it disagrees with the patient. Keep busy and cheerful; avoid fatigue and exposure; the application of cold water, in a systematical way, externally and internally, is highly recommended. (See “Hydrophathy.”) Exercise freely in the open air; travel and mingle in cheerful society.

#### 4. WATER-BRASH. HEART-BURN. (*Pyrosis.*)

(*Indigestion. Flatulency. Sour Stomach.*)

These ailments are mostly comprehended under the name of dyspepsia, of which they form some of the most distressing symptoms; but where they occur more isolated their remedies may be found in the following tabular view; reference to the details of these medicines may be had in the article “Dyspesia.”

TREATMENT. — For *water-brash*: *Nux vomica, Arsenic,*

*Carbo veg.*, *Sepia*, *Calcarea carb.*, *Phosphorus*, *Lycopodium*, *Sulphur*.

For heart-burn : *Nux vomica*, *Pulsatilla*, *China*, *Phosphoric acid*, *Sepia*, *Sulphur*.

For vomiting or throwing up,

a. Of food : *Arnica*, *Bryonia*, *Carbo veg.*, *Nux vomica*, *Phosphorus*, *Pulsatilla*, *Sulphur*.

b. Of indigested food (indigestion) : *China*, *Bryonia*, *Pulsatilla*, *Ignatia*, *Lachesis*, *Phosphorus*. (See end of this article.)

c. Of acid : *Nux vomica*, *Chamomile*, *Calcarea carb.*, *China*, *Phosphorus*, *Sulphur*.

For flatulency, or frequent rising of wind : *Arnica*, *Nux vomica*, *Pulsatilla*, *Bryonia*, *Rhus tox.*, *Sulphur*, *Carbo veg.*, *Veratrum*.

a. After eating flatulent food : *China*.

b. After eating pork or fat meat : *Pulsatilla*, *China*.

c. After drinking : *Nux vomica*.

d. Colic from flatulency : See "Colic."

In regard to indigestion and its consequences, we will particularly remark here that if it is caused by an overloaded stomach, a cup of coffee, without milk or sugar, will remedy most of its bad consequences; the remaining symptoms require *Pulsatilla*, *Nux vomica*, *Ipecac*.

Indigestion from fat food, pork, pastry, etc., requires *Pulsatilla*, *China*, *Carbo veg.*, *Ipecac*.

That which is caused by ice cream, fruit, or other things which chill the stomach, requires *Pulsatilla*, *Carbo veg.*, *Arsenic*.

If caused by the abuse of wine : *Nux vomica*, *Lachesis*, *Carbo veg.*, *Pulsatilla*, *Antimon. crud*.

By acid wines : *Antimon.*, *Pulsatilla*.

By sulphurated wines : *Pulsatilla*.

By vinegar, sour beer, and other acids : *Aconite*, *Carbo veg.*, *Arsenic*, *Hepar*.



If *tainted meat* or *fish* causes indigestion, give first a little *pulverized charcoal* in brandy; afterward *China* or *Pulsatilla*.

If *salt food* causes indigestion: *Carbo veg.*

Indigestion frequently produces the following disorders, which require the remedies attached to them:

*Fever*: *Aconite*, *Bryonia*, *Antimon.* (See “Gastric Fevers.”)

*Eruptions*, like *nettle-rash*: *Ipecac.*, *Pulsatilla*, *Bryonia*. (See “Nettle-rash.”)

*Diarrhea*: *Pulsatilla*, *Coffea*, *Ipecac.*, *Nux vomica*. (See “Diarrhea.”)

*Colic*: *Pulsatilla*, *Nux vomica*. (See “Colic.”)

*Flatulency*: *China*, *Pulsatilla*, *Carbo veg.* (See “Flatulency” above.)

In regard to *heart-burn* or *sour stomach*, we would remark that it will often be relieved by taking of a mixture of one drop of *Sulphuric acid* in a teacupful of water, every hour or two hours a teaspoonful; or by drinking lemon-water in moderate quantities. This is particularly recommended to pregnant women, who frequently suffer from acidity of the stomach from constitutional causes.

ADMINISTRATION, DIET, AND REGIMEN, the same as in “Dyspepsia.”

## 5. BILIOUSNESS. GASTRIC DERANGEMENT.

Biliousness, or bilious complaints, as a popular term, comprises different disorders of the digestive organs, which mostly have nothing to do with the bile or liver, but with the stomach alone. For practical purposes we divide these diseases:

In *gastric affections*, where the stomach alone is implicated; and

In *gastric bilious affections*, where the derangement of the liver reflects its morbid consequences on the stomach.

These attacks occur without much fever.

1. *Gastric affections (gastroataxia.)* This is a general name for all those disorders of the stomach, which we have already treated in the articles on "Dyspepsia," "Indigestion," "Heart-burn," "Water-brash," and "Flatulency," to which we refer the reader.

2. *Gastric bilious affections (gastrodynia biliosa.)* This disease is characterized by an *oppression* and *swelling* in the pit of the stomach, with inclination to vomit, and eructation of wind, of an offensive smell, like rotten eggs or meat; no appetite, and particularly a dislike for meat; thirst great, especially for acids; tongue thickly yellowish coated; either constipation or looseness of the bowels, of dark (black) offensive operations, with a great deal of wind. With these really bilious symptoms are combined a yellowish color in the face, around the nose and mouth; headache in the evening, with a full pulse and a feeling of heat without much fever.

CAUSES. — Persons removing from the north to the south are subject to this disease very often, which sometimes runs into bilious fever. Otherwise, it is caused more by the imprudent use of too much meat and alcoholic drinks in warm climates; also, by mental derangement, such as anger, grief, etc. It is, frequently, the forerunner of bilious fever.

TREATMENT. — Similar to "Bilious Fever," in its early stage. See below.

DIET AND REGIMEN. — Oatmeal gruel, with sugar and lemon-juice; lemonade, in small but often-repeated quantities; oranges; dry or milk toast; cold water. Meat, meat-soups, and eggs, are strictly forbidden. When recovering, a few oysters are allowed. If the bowels are very loose, the acids must not be used too freely, but very cautiously, if at all.

Cold baths are beneficial, particularly mornings and evenings. Exercise can be taken, on hills or dry places.

### NAUSEA. VOMITING.

Nausea and vomiting seldom occur as solitary symptoms, except when caused by errors in the diet; in which case, the ejection of the noxious substances by vomiting is most beneficial, and ought rather to be fostered, by drinking freely of lukewarm water, than checked by medicine. But when nausea or vomiting is caused by, or connected with, other diseases, their respective chapters must be consulted; where, generally, these two, as prominent symptoms, are mentioned and prescribed for.

We will, however, furnish the reader here with a list of the most frequent diseases in which nausea and vomiting exist, and attach the principal medicines thereto.

The principal remedies, useful in most cases of nausea and vomiting, are: *Ipecac.*, *Nuxvom.*, *Arsenic*, *Veratrum*, *Tartar emetic*, *Bryonia*, *Pulsatilla*, *Arnica*, *Cuprum*, *Sulphur*.

If nausea and vomiting is present

In *pregnant females*: *Nux vomica*, *Ipecac*, *Arsenic*, *Veratrum*. (See "Diseases of Females.")

In *drunkards*: *Nux vomica*, *Arsenic*, *Opium*, *Lachesis*, *Sulphur*.

In consequence of *riding* in a carriage, *sailing*, etc., *Cocculus*, *Petroleum*, *Arsenic*, *Ferrum*, *Secale*, *Belladonna*. (See "Sea-Sickness.")

In consequence of *worms*: *Cina*, *Aconite*, *Ipecac.*, *Nux vomica*, *Mercury*, *Sulphur*, *Lachesis*, *Carbo veg.* (See "Worm Disease.")

In consequence of *overloaded stomach*: *Pulsatilla*, *Ipecac.*, *Antimon.*, *Nux vom.*, *Arsenic*, *Bryonia*, *Rhus tox*, *Sulphur*. (See "Indigestion.")

After drinking : *Arsenic, Ferrum, China, Bryonia, Chamomile, Aconite, Arnica, Silicea.*

After eating : *Pulsatilla, Nux vomica, Bryonia, Arsenic, Ferrum, Sulphur.* (See "Dyspepsia.")

In the morning : *Nux vomica, Arsenic, Drosera, Veratrum, Hepar, Lycopodium, Silicea.*

In the evening, or at night : *China, Arsenic, Nux vomica, Ferrum, Silicea, Sulphur.*

After a fall on the head : *Arnica*, internally and externally. (See "External Injuries.")

In connection with a cough : See "Hooping-Cough."

In connection with headache : See "Sick Headache."

In connection with diarrhea : See "Cholera."

Or, in regard to ejected substances,

If blood is vomited (*hæmatemesis*) : *Aconite, Arnica, Ipecac., Nux vomica, Ferrum, Hyosciamus, Bryonia, China.* (See "Inflammation of the Spleen," "Cessation of Menses," and "Hemorrhoids.") Generally, medical aid is immediately required, when blood is vomited. If the blood is very dark,

*Black vomit (melaena)*, with discharge of similar blood by the stool : *Veratrum, Arsenic, China, Ipecac.* (See "Diarrhea, Typhus.")

If fecal matter is ejected : *Opium, Nux vomica.* (See "Constipation," and "Ileus.")

If bile is vomited, of greenish look and bitter taste : *Chamomile, Ipecac., Nux vomica, Pulsatilla, Antimon., Cuprum, Veratrum.* (See "Biliousness," etc.)

If mucus of a sour taste or smell : *China, Calcarea carb., Nux vomica, Pulsatilla, Sulphur.* (See "Dyspepsia.")

If watery substances : *Bryonia, Ipecac., Bellad., Pulsatilla, Sulphur.* (See "Water-brash," etc.)

ADMINISTRATION. — Give the medicine either dry, every hour or two hours, from three to six glob.; or, dissolved in

water (twelve glob. in half a teacupful), every half hour a teaspoonful, until better, or until three or four doses of a medicine are taken, when another remedy must be selected, if the patient is not better.

DIET AND REGIMEN. — In feverish attacks, see “Fever.” In all others, see their respective chapters.

#### SEA-SICKNESS.

This distressing complaint depends greatly on constitutional peculiarities, to which the remedy must be adapted. As the cause (the motion of the vessel) continues, it is necessary to repeat the medicine frequently. It will be of service to be careful in the diet just before embarkation; especially to avoid rich, fat food.

TREATMENT. — The principal remedies are :

*Cocculus*. In giddiness, headache, nausea; the patient feels better when lying down.

*Nux vomica*. The same. The patient feels better when *not* in the open air.

*Pulsatilla*. The patient feels better *in the* open air, on deck.

*Colchicum*. After *Pulsatilla*, or when the smell of food or the scent of the vessel sickens.

*Tubacum*. If *Pulsatilla* or *Colchicum* are insufficient, or when the patient sickens at the *slightest* movement of head or body, with *deadly paleness of the face*.

*Petroleum*. In cases of great debility.

*Arsenic*. If the patient becomes *very weak*, and there is violent retching.

*Ipecac*. If there is frequent vomiting, without feeling very weak.

For *constipation*, during a voyage on sea, give *Opium*, *Cocculus*, *Nux vomica*, *Lachesis*, *Sulphur*.

When constipation is attended with a putrid taste and bloody gums, give *Staphysag.*

When patients are constipated, and have a longing for acids and something piquant, give *Sepia*.

N. B. It is well enough to provide one-self on sea voyages with Scotch ale and herrings; also, lemons; as those things are mostly desired by sea-sick patients.

ADMINISTRATION. — The same as in “Nausea and Vomiting.” It is beneficial to take as much exercise on deck as possible, whatever the weather may be, although the patient may not incline to do so at first.

#### SPASMS, AND PAIN IN THE STOMACH.

(*Cardialgia, Gastralgia, Neuralgia of the Stomach.*)

DIAGNOSIS. Spasmodic pains and contractions in the pit of the stomach, sometimes slight, sometimes of insupportable violence, and spreading to the chest and back, exciting nausea and vomiting; anguish; coldness of the extremities, and even faintness; belching up of wind often relieves the patient for a time, as, also, vomiting of an acid, limpid fluid, if acidity of stomach or a gouty disposition was connected with it.

CAUSES. — Any derangement of the nerves of the stomach, either sympathetic, from diseases of the liver, spleen, or pancreas, or more directly from long fasting, the habitual use of *ardent spirits, indigestible food* (new bread, nuts, sweetmeats, chesnuts, cherries, figs, cheese, etc.), *coffee* or *tea*, may cause this disease; especially if, in addition to this, the person exposes himself to cold, damp weather, or if he is of a gouty disposition, which then develops itself on the stomach in this form (*rheumatism* and *gout* of the stomach.) Females are particularly liable to this disease, which, in them, appears mostly with derangement of the



menses, especially after their cessation<sup>1</sup> (*neuralgia* of the stomach), when it is sometimes associated with fits of fainting, and may end in vomiting of blood. In such cases, the periodical appearance of the disease is frequently observed.

**TREATMENT.** — Persons liable to spasms of the stomach must avoid every thing which could cause them directly, as mentioned above. Besides this, however, they should keep their feet warm, and wear a woollen shirt or bandage, or, in obstinate chronic cases, a Burgundy pitch plaster on the stomach. Above all, however, we recommend to apply to a homœopathic physician, who, in most cases, is able to eradicate the complaint. For recent cases, or in the attack itself, the following remedies will be found beneficial.

We give, first, a tabular view, to facilitate the choice of the proper remedy.

*Spasm in the stomach (gastralgia), from*

Abuse of *Chamomile tea*: *Nux vomica*, *Ignatia*, *Pulsatilla*.

Abuse of *coffee*: *Nux vomica*, *Ignatia*, *Cocculus*, *Chamomile*.

Abuse of *ardent spirits*: *Nux vom.*, *Lachesis*, *Arsenic*, *Sulphur*.

*Indigestion*: *Pulsat.*, *Antimon. crud.*, *Bryonia*, *China*, *Carbo veg.*, *Nux vom.*

*Want of exercise, torpid bowels*: *Nux vom.*, *Sulphur*.

*Excess of mental emotions, as anger, indignation, etc.*: *Chamomile*, *Colocynth*, *Nux vomica*, *Staphysag.*

*Loss of animal fluids, causing excessive debility, such as blood-letting, nursing, perspirations, excessive diarrheas*: *China*, *Carbo veg.*, *Phosphoric acid*, *Coculus*, *Nux vomica*.

*Rheumatism, and exposure to cold and damp weather*: *Chamomile*, *Bryonia*, *Rhus tox.*, *Carbo veg.*

*Derangement of the menses*, if they are too slight : *Pulsatilla*, *Cocculus*. If too profuse : *China*, *Bellad.*, *Platina*, *Calcarea carb.*

**SYMPTOMATIC DETAIL.** — *Nux vomica*. This is one of the most important remedies in this disease, and it is efficacious almost in every spasm of the stomach, in the beginning ; at least, there is no risk in giving directly one dose (two, three, or four glob.), as it will procure time for cooler observation and a more correct choice, according to the symptoms. If *Nux vom.* does not give any relief, *Chamomile* or *Cocculus* alternated, every half hour, if necessary, will be of great benefit. The pains requiring *Nux vom.* especially, are : *contracting*, *pressing*, and *spasmodic*, with sensation of *retraction or clawing* in the stomach, with oppression of the chest, as if it were compressed by a band, with pain extending into the back and loins ; nauseous accumulation of water in the mouth, or vomiting of sour liquid or food during the pain, *worse* after a *meal*, *from using coffee* and *ardent spirits* ; *from over-study*, creating thereby constipation, piles, hypochondria, with *irascible* humor, hasty and passionate character. Such persons generally are subject to nervous, pressing headache on one side, with unfitness for exertion, palpitation of the heart, with anxiety.

Administration, see at the end of this article.

*Cocculus*. Often very beneficial, when *Nux vom.* or *Chamomile* has given but temporary relief, or when there are spasms in the stomach, mitigated by belching up of wind ; otherwise, similar to *Nux vom.*

*Chamomile*. Distention of the stomach, with *pressure*, as from a stone, or as if the heart were being crushed with oppression and shortness of breath ; pains are worse at night, with *great anguish* and *tossing* ; *momentary mitigation by drinking coffee*. The patient is *peevish* and *irritable* ; pains drive him out of bed. This remedy suits well to be

given alternately with *Coffea*, and if then no improvement takes place, try *Bellad.*

*Belladonna.* Especially where *Chamomile* was of no avail, although well indicated, mostly in women of a delicate, sensitive constitution, at the time of too *profuse menstruation*, and then alternately with *Platina*, until better. *Bellad. cc.*, alternately with *Coffea cc.*, will be very beneficial in this disease, if the patient is decidedly better, but can not sleep nor lie quiet. *Belladonna* also suits in such violent pains as to *take away consciousness*, and cause *falling down*; also, great thirst, but the pain gets worse after drinking.

*Pulsatilla.* Is a principal remedy in mild, sad, tearful dispositions, or where the menses are tardy, producing those spasms. The pains are *shooting*; worse by walking and making a false step; spasmodic pains in the stomach, caused by *fasting*, or *overloading* the stomach; with nausea, *vomiting of food*; not much thirst; pains are *worse in the evening*, with feeling of *chilliness*; frequently loose bowels.

*Ignatia.* Frequently after *Pulsatilla*, and when there are *pressing pains*, as *from a stone*, especially after a meal or at night; weakness, burning, emptiness in the pit of the stomach, sensitive to touch; aversion to food and drink; can not bear tobacco; also, pains from mental causes, indignation, etc.; or suits well for persons who have the pains from having fasted too long.

*China.* Spasms from indigestion, with swelling of, and painful pressure in, the stomach, with rumbling of wind and flatulency; spasms after loss of strength by blood-letting, nursing, diarrheas, etc.

*Carbo veg.* Follows well after *China*, if the spasms were caused after eating *flatulent* or *spoiled* food; also, spasms occurring in damp and wet weather; particularly if there is a *painful, burning pressure*, with *anxiety*, worse on touch or at night; or contracting, spasmodic pain, forcing the

patient to bend double, with shaking; worse when lying down; heart-burn; nausea; the very thought of food sickens him.

*Lachesis.* *Pressing pains*, ameliorated soon after a meal, but renewed afterward, with *constipation*, dyspepsia, and flatulency.

*Lycopodium.* Compressive pains, ameliorated in bed during the night; worse in the morning, after a meal, and especially in the *open air*.

*Arsenic.* *Violent pains* and *anxiety* in the cardia and the stomach; *burning* and spasmodic pains in cardia and stomach; *vomiting after drinking the smallest quantity of liquid*; suitable for spasms in old people, where cancer of the stomach may be supposed to exist, and in drunkards.

*Sulphur.* *Pressing pains*, as from a stone, mostly after a meal, with nausea, water-brash, or vomiting; *acidity*; *throwing up of food*; disposition to piles, or accumulation of mucus in the intestines; hypochondriacal, whining mood.

*Antimon. crud.* See "Indigestion."

*Bryonia.* *Pressure as from a stone* in the cardia, when eating, or after a meal, with a sensation of *swelling in the pit of the stomach*; pains worse by any motion; constipation; rheumatic pains in head and limbs. In alternation with

*Rhus tox.* When the *pressure in the stomach and cardia obstructs the breathing*, particularly after taking cold in damp, cold weather. (See "Bilious Rheumatic Fever.")

*Staphysag.* Spasms in the stomach after anger or indignation.

*Calcarea carb.* In females who have their menses very profusely; or after *Belladonna* has given partial relief; especially when there is a sensation of *clawing* and retraction in the stomach, worse after a meal, *frequently with vomiting of food*; disposition to piles, constipation, and palpitation of the heart.

ADMINISTRATION. — Dissolve of the remedy selected twelve globules in half a teacupful of water, and give, every thirty or forty minutes, a teaspoonful, until three or four are taken ; then wait thirty minutes, and, if not better, select another remedy, and give it in the same manner, until better. This is the mode of administration in the most urgent cases. When the pains are not so severe, give the medicine at longer intervals—say, from one to two, three, or six hours, until amelioration takes place. In real chronic cases of this disease, give the chosen remedy only evening and morning a dose (four glob.), for two days, and then wait the same length of time, before another remedy is given ; if better during that time, no more medicine is necessary.

DIET AND REGIMEN. — The same as in “Biliousness,” and “Gastric Derangement.”

Externally, a mustard plaster, or other heating substances, such as scorched flour, heated bran in small sacks, may be laid on the stomach. The patient is advised to remain as quiet as possible during the paroxysms.

GRIPING COLIC. (*Colica ; enteralgia.*) (*Stomach-ache.*)

DIAGNOSIS. — Pains in the intestines, more or less violent ; griping, pinching, tearing, or burning, mostly in the region of the navel, yet often spread all over the bowels ; severe cases of colic are accompanied by anxiety and cold sweat ; stools either loose or confined ; when the colic pain is seated in *one* place of the abdomen for a length of time without moving, danger of inflammation of the intestines ensues, which is indicated by the pain, hitherto griping and pinching, becoming *burning*, and the fever appearing (see “Inflammation of the Intestines”) ; vomiting of sour, bilious phlegm, or distention of the abdomen, painful to the touch, often attend the severest cases of colic.

CAUSES.—These are constitutional or acquired by circumstances; the latter are very many, such as colds, indigestion, worms, constipation, mental emotions, and occupations which require persons to work in deleterious metals, such as lead, etc. According to these exciting causes, the three principal varieties of colic are,

1. *Bilious colic*, with bilious vomiting, severe griping, or shooting or twisting pain in the abdomen; at first relieved by pressure, afterward tender to the touch, with painful distention, cold extremities, restlessness, yellowish cast of the skin and eyes, the face expressing great anguish.

2. *Flatulent or wind-colic*, with frequent belching up of wind, with or without relief; swelling of the abdomen, which is distended and painful in different places, as the wind accumulates and moves in different parts of the intestines; the pains come in paroxysms, and are sometimes very severe.

3. *Painter's or lead colic*, which commences sometimes before an attack with general lassitude, wandering pains in the bowels and extremities, heaviness in the lower limbs, chilliness, and depression of spirits; the attack itself exhibits the same symptoms as bilious colic, except there is not often vomiting of bile, but greater pain and restlessness, during which the patient bends forward, pressing firmly against the abdomen, his extremities feeling cold, and his pulse being small and suppressed.

TREATMENT.—The first thing to be done in case of any colic, if it is at all severe, is the application of a general warm bath, the patient sitting in a tub, the warm water reaching up to the pit of the stomach, and covered with blankets to confine the hot steam closely around him; after ten minutes, he ascends from the bath and lays down with the blankets, without being dried off, and is covered with sufficient clothes; the warmth then may be kept up by putting heated bricks or hot water flasks around the



patient, on the spots where he complains the most ; a profuse perspiration will ensue, which relieves the cramp and makes the patient feel easier at once ; it must be kept up until every vestige of pain has left for some time. The medicines to be prescribed below may be given, however, from the commencement of the pain, and continued until relief is obtained. We will now give in a tabular view the principal kinds of colic, with their remedies, the symptomatic details of which may be consulted below.

a. *Bilious colic* : *Nux vomica*, *Colocynth*, *Chamomile*, *Bryonia*.

b. *Flatulent, or windy colic* : *Pulsatilla*, *China*, *Carbo veg.*, *Cocculus*, *Nux vomica*, *Colocynth*, *Lycopodium*, *Sulphur*.

c. *Painter's or lead colic* : *Opium*, *Belladonna*, *Platina*.

d. *Colic with obstinate constipation* (see "Miserere") : *Opium*, *Nux vomica*.

e. *Colic caused by piles or hemorrhoids* : *Nux vomica*, *Lachesis*, *Pulsatilla*, *Colocynth*, *Carbo veg.*, *Sulphur*.

f. *Colic from an inflammatory state of the intestines* (inflammatory colic) : *Aconite*, *Belladonna*, *Hyosciamus*, *Mercury*, *Lachesis*, *Nux vomica*, *Arsenic*, *Sulphur*. (See "Enteritis.")

g. *Colic from worms* : *Mercury*, *Cina*, *Sulphur*. (See "Worms.")

h. *Colic from indigestion* (gastric colic) : *Pulsatilla*, *China*, *Nux vomica*, *Belladonna*, *Bryonia*, *Carbo veg.*,

i. *Colic from indignation or rage* : *Chamomile*, *Colocynth*, *Staphysag.*, *Sulphur*.

k. *Colic from a chill* : *Aconite*, *Chamomile*, *Colocynth*, *Mercury*, *Nux vomica*.

l. *Colic from exposure to cold, damp weather* : *Pulsatilla*, *Bryonia*, *Rhus tox.*

m. *From bathing* : *Nux vomica*.

n. From *external injuries*, such as *strains* or *blows* : *Arnica*, *Bryonia*, *Rhus tox.*, *Carbo veg.*, *Lachesis*.

o. Colic in *children* principally : *Chamomile*, *Rheum*, *Coffea*, *Aconite*, *Belladonna*, *Cina*, *Sulphur*. (See "Diseases of Children.")

p. Colic in *pregnant* or *lying-in women* : *Chamomile*, *Nux vomica*, *Arnica*, *Bryonia*, *Pulsatilla*, *Sepia*. (See "Diseases of Women.")

q. In *hysterical women* (nervous colic of women) : *Ignatia*, *Nux vomica*, *Cocculus*. (See the same.)

r. Colic at the time of the *menses* (menstrual colic) : *Pulsatilla*, *Coffea*, *Veratrum*, *Cocculus*. (See the same.)

s. Colic in persons with *low spirits* : *China*, *Nux vomica*, *Sulphur*, *Calcarea carb.*, *Stannum*.

Select first, according to the character of the disease, one of the above remedies, and then read its symptoms in the following symptomatic detail. It will be seen above that *Nux vomica* and *Colocynth* are the principal remedies in almost all kinds of colic, and consequently mostly to be used ; we recommend, therefore, the careful study of their distinguishing symptoms. Beside these two medicines, *Pulsatilla* and *Belladonna* will be found of frequent application.

Administration of these remedies, see at the end of this article.

SYMPTOMATIC DETAIL. — *Nux vomica*. Constipation, *hard* or *difficult stool*, with *pressure in the abdomen as from a stone*; the colic pain is *contractive* or *compressive*, sometimes pinching and drawing; *pressure at the pit of the stomach*, with distention and tenderness of the abdomen when touched; feeling as if the clothes around the waist were too tight; flatulency and severe *griping*, deeply seated in the abdomen, with cold hands and feet, and sometimes sharp and severe *pressure on the bladder and rectum*, as if wind were

about to escape violently, forcing the patient to bend double; beneficial in almost all kinds of colic, where *constipation* predominates, or the patient likes to be quiet, to sit doubly bent, or to lie down on account of violent pains in the loins and head.

*Colocynth.* This is the principal medicine for colic ; *violent*, even the *most violent cutting, pinching clawing* pains and *stitches*, as from *knives*, with *excessive restlessness, agitation* and *tossing*, sometimes *cramp* in the *limbs*, *shivering*, either no inclination to stools or *violent diarrhea* and *vomiting of bile* or food, immediately after eating; abdomen either swollen with wind, *flatulency*, or feels *very empty*; however, very tender, as from a bruise. After it are suitable *Chamomile, Mercury, Belladonna*.

*Belladonna.* The colic pains are: swelling and *protrusion* like a pod of the colon between the pit of the stomach and the navel, relieved by pressing upon it or by bending double; pain at the same place, as if the intestines were *grasped by the finger nails*; or *spasmodic constriction* in the abdomen in females, with a feeling of *bearing downward* as if something would fall out, similar to the falling of the womb, or pains connected with this disorder; general characteristic for *Belladonna*: congested hot head, red face, severe pain in the head, aggravated by movement, can not bear the light, feels dry in the throat, etc. (See “*Materia Medica*.”) After it *Mercury, Lachesis*, or *Hyosciamus*, is suitable.

*Pulsatilla.* Shooting, beating pains in the pit of the stomach; farther down in the abdomen a feeling of heaviness and fullness, with *disagreeable tension* and *distension*, tenderness and pain as if from a bruise when touched; suffering from flatulency (see this article), and consequent colic after indigestion; worse when *sitting, lying*, or in the *evening*, with *shivering*; patient feels better *out doors*, than

within; is inclined to, or has diarrhea; *pale face*, with livid circle around the eyes; *pressure*, tension, and aching in the head. After it are suitable, *China*, *Lycopodium*, *Mercury*, *Sulphur*.

*China*. Excessive distension of the abdomen, as in *tympanitis*, with *fullness* and *pressure*, as if from hard bodies; or spasmodic constriction; pain, with *stoppage of wind* and *bearing toward the hypochondriacal region*; worse at night, after errors in diet, and in persons weakened by severe loss of animal fluids, such as bleeding, too heavy perspiration, nursing, diarrhea, etc. (See "*Pulsatilla*."')

*Cocculus*. One of the principal medicines, particularly in spasmodic and windy colic, and in those colics of women before or at the time of their menses (see "*Female Diseases*"); or when there are *constrictive, spasmodic* pains in the hypogastrium, nausea, difficulty of breathing, flatulency, and full, distended stomach; or a feeling of emptiness in the abdomen, with squeezing and tearing in the stomach; nausea; *constipation*; nervous excitability.

*Coffea*. *Excess in pain, which drives to despair*; great agitation, tossing, grinding of the teeth; coldness of the limbs; moaning, and fits of suffocation.

*Ignatia*. Colic in the night, waking out of a sleep, with shooting into the sides and chest; difficult, but relieving discharge of wind; suitable for sensitive and delicate women.

*Hyosciamus*. Spasmodic and griping pain, with vomiting; cries; pain in the head; hardness and distension of the abdomen, and tenderness when touched.

*Chamomile*. *Tearing, drawing pains*, with great restlessness; nausea; *bitter vomiting* or *bilious diarrhea*; incarcerated wind, with anguish, *tension, pressure, fullness in the pit of the stomach*; one cheek red, the other pale; the colic

appears mostly at *night*, toward sunrise, or after a meal. (*Pulsatilla* suits after it.)

*Bryonia*. Fullness and pressure in the abdomen after eating; cutting and *stitching* pains in the bowels, particularly after drinking warm milk; colic pains with diarrhea after taking cold, or in the heat of summer. Alternately with *Rhus tox.*, in bilious rheumatic attacks. (See this article.)

*Rhus tox.* Pressure in the stomach and pit of the stomach; *pain in the abdomen at night*; restlessness and diarrhea of watery and slimy substances. (See *Bryonia*.)

*Sulphur*. Against *hemorrhoidal colic*, when *Carbo veg.* or *Mercury* has been given without effect; against *bilious colic*, when *Chamomile* or *Colocynth* was insufficient; against *flatulent colic*, after *Chamomile*, *Calcarea carb.*, *Mercury*, or *Carbo veg.* was administered; or against *worm colic*, after *Mercury* or *Cina* had been given.

*Carbo veg.* Distension and fullness of the abdomen; difficult belching up of wind; colic after riding in a carriage; flatulent colic, with rumbling in the bowels; discharge of wind of a putrid smell; either constipation, or diarrhea of a putrid smell.

*Arsenic*. Excessive pain, with great anguish in the abdomen; intolerable burning, or sensation of cold in the abdomen; pains are worse at *night*, after eating or drinking; watery or bilious vomiting; diarrhea, with thirst, shivering, and excessive debility.

*Veratrum*. Cutting as from knives in the abdomen, which is very tender to touch; burning in the whole abdomen; flatulent colic, with noisy rumbling of wind, which is discharged with difficulty. (Alternately with *Arsenic* in severe cases, or with *Coffea* at the time of menstruation.)

*Opium*. Abdomen hard and distended; tympanitis, with heaviness in the abdomen, as from a weight (*lead* or

*painter's colic*); *obstinate constipation*, with vomiting even of fæces and urine; or involuntary stools, of a dark color and fetid.

*Lachesis*. In *spasmodic* and *inflammatory* colic, when *Colocynth*, *Belladonna*, *Chamomile*, *Nux vom.* have been administered without effect; in *hemorrhoidal* colic, and after *mechanical* injuries; after *Carbo veg.*, when there is great weakness, sinking feeling, small, quick pulse; or alternately with *Lycopodium*, if there is great constipation left. (See this article.)

*Lycopodium*. In *flatulent colic*; *swelling*, *fullness*, and *distension* of the pit of the stomach, with colic pain, incarceration of wind, which is not discharged. (Is suitable after *Pulsat.* or *Lachesis*; see these.)

*Platina*. In consequence of *fear* or *anger*, or *poisoning by lead*, after *Opium* and *Belladonna*; suitable principally for females, or when the bodily sufferings disappear as soon as the mental distresses begin, or *vice versa*; is afraid of dying; can not refrain from weeping; *contracting pain in the abdomen*; *pressure in the stomach after eating*; also a contracting pain, as if laced too tight; with a sensation of constant bearing down.

*Aconite*. *Extreme tenderness of the abdomen*; much anguish, restlessness and tossing, particularly, when the pains are located in the region of the bladder; when the bowels seem to retract, with frequent but unsuccessful want to urinate.

*Arnica*. *Contusive pain in the sides of the abdomen* in pregnant females; pain from lifting in the abdomen; *fullness in the stomach*, as if one had eaten too much; *stitches in the pit of the stomach*, with *pressure back to the spine*, and *an oppression on the chest*, worse after eating, drinking, and touch; stitches in the left *hypochondria*, *taking away the breath*; *distension and hardness of the abdomen*, with sore-



ness in the sides, better after discharge of urine ; diarrhea, watery, or from indigestion.

*Mercury.* 'Shooting, or violent contracting pains in the abdomen, especially around the navel, with nausea ; frequent desire to evacuate, or *slimy diarrhea* ; profuse quantity of saliva in the mouth ; shivering, with heat, especially in the face ; abdomen tender to touch ; colic pains, worse in the night ; great lassitude.

*Cina.* Colic pains from worms, especially around the navel, which is tender to the touch. *Cina* <sup>cc.</sup> in such cases, every half hour repeated (the dose from two to four glob.), is more effectual than the other preparation of *Cina* (see "Worms.")

*Rheum.* Particularly in infants, when the colic is accompanied by diarrhea of a *sour smell* and pressing down before and after the stool ; or when there are in adults *cutting pains* in the abdomen, forcing one to *bend over*, frequently soon after eating ; worse when standing.

*Sepia.* Colic pains in *pregnant or lying-in women*, or even *spasms*, as if the intestines were turning over ; *cutting pains after motion* ; *burning and stitches* in the abdomen, which is *hard and distended* ; *pressure and sensation of heaviness* in the abdomen, with *bearing down* ; *rumbling of the bowels* after eating ; constipation, or greenish diarrhea, especially in children.

*Calcareæ carb.* Contractive, gnawing spasms, or cutting pains in the abdomen ; difficulty of discharging the wind, with noisy rumbling of the bowels ; can not bear the clothes tight around the hypochondria ; *acidity of the stomach*, with vomiting of food, especially in children when teething ; lightish diarrhea.

*Stannum.* Spasms in the region of the *diaphragm*, as in *hysterical or hypochondriacal persons*, with difficulty of eject-

ing the wind; constipation, or slimy diarrhea, and great weakness. (Compare *China*.)

ADMINISTRATION. — Dissolve twelve glob. of the selected remedy in half a teacupful of water, and give every fifteen or thirty minutes a teaspoonful, for two or three times; then wait the same length of time before giving another remedy, if not better; but if the patient is better, do not repeat the medicine as often. This is particularly the case with *Colocynth*, which is the most frequently needed remedy. In the most agonizing pains, the alternate use of *Coffea* and *Camphor*, in solution, given as above stated, affords temporary relief, until other remedies may complete the cure. In cases where the pains are not so excessive, the medicines must not be repeated as frequently; say, only from two, three, to six hours a dose (four glob.) until better, or until another remedy is needed.

DIET AND REGIMEN. — Persons subject to this complaint must avoid all substances which produce it, such as kale (kohl) and other green vegetables, acidulated drinks, veal, etc. In an attack of colic, give only the mildest nourishment—gruels, toast-water, toasted bread, etc.—no milk, except in worm colic, when *sweet, cold* milk, will pacify the worms. Keep the feet dry and warm, also in wet, cold weather a flannel bandage around the bowels. In colic from indigestion, a cup of black coffee *immediately* taken is strongly recommended.

#### CONGESTION AND STAGNATION OF BLOOD IN THE ABDOMEN.

That process, by which disease is generated and grows, depends upon certain laws, which in no case appear in more constant regularity, than in producing the different kinds of congestions. The three great divisions of the body—head, breast, and abdomen—are, at certain periods of man's life, successively, the places for congestions and stagnation.

Thus the head inclines to be congested in childhood, the breast in youth up to thirty or thirty-five years (hence, in this period most all lung-diseases occur), and, finally, the abdomen in middle age, from thirty-five to old age (hence in this period so many bowel-diseases, such as hypochondria, piles, dyspepsia, etc., take place.) One of the most natural causes for these various diseases is *the determination of blood to the abdomen*. We must not increase, by our mode of living, this natural tendency to congestion; as, if neglected or mistreated, its cure will be very difficult.

DIAGNOSIS. — Persons troubled with this disease have a sensation of heat and burning in the stomach, together with the feeling of heaviness, hardness, and tension in the lower part of the abdomen.

TREATMENT. — *Nux vomica* and *Sulphur* are the principal remedies.

*Nux vomica*. Especially for persons who lead a sedentary life, are *constipated*, have pains in the loins, hips, and back, as if broken and very weak; hardness and tension of the abdomen; can not bear the pressure of the clothing around the waist.

*Sulphur*. After *Nux vomica* has relieved partially, in all cases, even the most obstinate, where constipation, hypochondriacal and hemorrhoidal symptoms are present; *Nux vomica* alternately with *Sulphur*, every second or third day a dose (four to six glob) until four doses of each are taken, will generally be sufficient to relieve; if not, repeat it, after an interval of from eight to fourteen days, during which time no medicine must be taken.

*Arsenic* and *Carbo veg*. Administered in the same manner, if there is great weakness, with tendency to diarrhea and flatulency.

*Capsicum*. If the evacuations are soft, trifling, slimy,

watery, give twice or three times a dose (four glob.), until better.

Other remedies to be used successively, but slowly, are : *Pulsatilla*, *Sepia*, *Belladonna*, *Mercury*, *Lachesis*, *Lycopodium*.

DIET AND REGIMEN. — Exercise on foot and horseback ; the systematic use of cold water (see “Hydrotherapy”) ; regularity of all habits and cheerfulness of mind are necessary.

DIET. — See “Piles.”

#### INFLAMMATION OF THE DIAPHRAGM. (*Diaphragmitis*.)

This disease resembles pleurisy very much ; sharp, shooting pains under the short-ribs ; worse on pressing between or below the ribs, on sneezing, coughing, inhaling, or striking the body ; relieved by bending the body forward ; frequently there is hiccongh present.

CAUSES. — The same as in pleurisy. We have only to add, that external injuries, such as blows, falls on the ribs directly or a jarring of the diaphragm by a jump on the heels and feet, or (as in little children it is often the case) a twisting or unnatural turning of the body, are the most frequent causes of this disease.

TREATMENT. — The same as in “Pleurisy.”

The principal remedies are : *Bryonia*, *Aconite*, *Nux vom.*, *Chamomile*. This latter, particularly in infants and children, when they are swollen around the short-ribs, where the slightest touch can not be borne without screaming or stoppage of breath (see, in “Diseases of Children,” “Liver-grown.”) *Nux vomica* when *Chamomile* seems to be insufficient, and there is extreme constipation.

If caused by external injuries, *Arnica* externally, in solution, and internally, in alternation with *Aconite*, every hour or two hours a dose (six glob.) must be used until better.

ADMINISTRATION, DIET AND REGIMEN, the same as in "Pleurisy."

### ABDOMINAL INFLAMMATION.

We now proceed to the inflammations of those organs which lay between the peritoneum, or that serous membrane which envelops, as within a sack, all the abdominal viscera. The inflammations of these organs have this one peculiar feature in regard to the pulse, that the higher the inflammation runs, the smaller and more suppressed, often wiry, the pulse appears; and that the pulse itself, as an indication of inflammation, is frequently inconsiderable, at least, only discoverable by hard pressure.

#### 1. *Inflammation of the Stomach. (Gastritis.)*

DIAGNOSIS.—This is a dangerous disease, and ought not to be treated by laymen; but, on account of its rapid course, we feel the necessity of introducing it here, in order to enable every one to diminish, or take away entirely its danger, by applying those remedies which have a specific bearing upon it.

Gastritis is characterized by a constant, violent, burning, and stitching pain in the gastric regions, increased by pressure, respiration, or taking any kind of food; besides this, there is heat, swelling, and tension, with pulsation in the region of the stomach; nothing will stay on the stomach, not even pure water; although there is great thirst, yet water is frequently rejected, because sometimes, besides other violent nervous fits and convulsions, hydrophobic symptoms occur, with tetanic spasms, great and sudden prostration; fainting; the pulse gets smaller, more wiry; extremities cooler, and the restless anxiety greater, the higher the inflammation is. Death takes place, either by gangrene (which is announced by sudden cessation of all pain, small, remitting

pulse, scarcely perceptible, hands and feet icy cold), or by nervous paralysis, with spasms and fainting fits, together with symptoms of the utmost debility. Signs of improvement are, when the pulse becomes larger and broader, the pains less severe, without ceasing entirely, and the hands and feet warmer.

This disease may, after an improper treatment, pass into chronic inflammation, induration, or ulceration of the stomach.

CAUSES. — Bilious diarrheas, if suddenly checked; acrid poisons taken into the stomach; abuse of emetics and ardent spirits; transfer of gout; suppressed secretions; external injuries; or drinking of ice water when heated.

TREATMENT. — *Aconite* is the first remedy to be given, if the pains are severe, and the heat great in the stomach, or when a chill preceded the affection, or when it was caused by taking cold drinks when overheated.

ADMINISTRATION. — Dissolve twelve glob. in half a tea-cupful of water; give every twenty or thirty minutes a tea-spoonful until ten spoonfuls are taken; if then not better, select another remedy, which must be prepared and given in the same manner, until better, or another remedy is necessary.

*Bryonia*. Generally after *Aconite* or *Ipecac*.

*Ipecac*. After *Aconite*, if vomiting predominates, and the disease was caused by indigestion, but the tongue is clear.

*Antimon*. Under similar symptoms as *Ipecac*.; but the tongue has a thick coating of white or yellowish mucus.

*Pulsatilla*. After *Ipecac*. or *Bryonia*, or when the disease was caused by indigestion, or a chill in the stomach from having taken ice.

*Arsenic*. For a similar condition, but there is a *rapid failure of strength*, pale, sunken countenance, cold extremities; alternately with *Veratrum*.



*Veratrum*. For similar symptoms (see “Asiatic Cholera.”)

*Belladonna* and *Hyoscyamus*, alternately, if nervous, spasmodic, hydrophobic symptoms appear; drowsiness, delirium, loss of consciousness.

*Arnica*. In case external injuries were the cause of the disease.

*Nux vomica* and *Lachesis*, in alternation, if abuse of ardent spirits caused the gastritis; after which,

*Lachesis* and *Arsenic* may be alternated with benefit.

*Cantharis*. In the most violent cases, particularly when the burning pain is intolerable; alternately with *Coffea*.

*Opium* and *Camphor*, alternately, if no remedy seems to ameliorate the case; drowsiness and stupor, particularly in drunkards; afterward, give other suitable remedies which will have a better effect.

For gastritis from acrid poisons, see “Poisoning.” After the poison is removed, treat as above.

For the chronic form of the disease, see “Dyspepsia, Cardialgia, Spasms of the Stomach.”

DIET AND REGIMEN.—As during the height of the disease nothing will stay on the stomach, it is better to offer the patient nothing but water; afterward the mildest gruel, made of rice or flour, milk and water, buttermilk, gum-arabic water, milk toast. Continue this diet for some time before it is changed for a stronger one; and then only soups and mild vegetables.

Externally apply, if necessary, on the stomach, fomentations of hops, mustard draughts on the feet, and, if possible, a lukewarm bath.

## 2. *Inflammation of the Bowels. (Enteritis.)*

This disease occurs in two varieties; either *idiopathic*, where no other disease preceded it, or *symptomatic*, when

it appears in consequence of other diseases, such as “Typhus,” etc. The former is very acute, but of rare occurrence, and involves in its sphere the mucous membrane, as well as the submucous tissue and peritoneal coat—a very painful and dangerous disease; while the latter, not so acute in its course, occurs more frequently, attacking the mucous membrane of the intestines, mostly in portions; being, on that account, not so painful and dangerous, but sometimes leading to ulcerations of the bowels.

DIAGNOSIS of the *idiopathic* or *very acute form*.—Violent burning or pungent pain, permanently on one spot of the abdomen, generally in the region of the navel; increased by the slightest touch or movement; with tension, heat, and bloatedness of the abdomen; obstinate constipation; vomiting of slime or bile, finally of fæces (*ileus miserere*); violent thirst, yet cold drinks do not agree; sobbing, anxiety; restlessness; pulse small, contracted as in gastritis; urine frequently suppressed.

The fatal termination of this disease occurs under similar symptoms as in gastritis by gangrene; the most violent pain suddenly disappearing, pulse sinking, remittent, imperceptible; involuntary, cadaverously fetid stools.

The symptoms of recovery are the same as in gastritis. In regard to the obstinate constipation which is such a constant symptom in this disease, we would remark that this does not constitute or cause the disease, but is its natural consequence, and disappears as soon as the inflammation, by the proper remedies, is reduced. Any violent attempt, by severe medicines, to effect a passage, would only add to the inflammation, and hasten a fatal result. The best expedient for this purpose is, besides the medicines to be mentioned hereafter, injections of tobacco smoke, which relieve the pain, or lukewarm baths, which destroy the spasms, often remaining after the inflammation is subdued.

TREATMENT. — *Aconite* must be used first, as directed in gastritis. It is the principal remedy, as long as the accompanying fever is intense—the skin very hot.

In severe cases, *Arsenic* and *Veratrum* after it, in a similar manner, as directed in gastritis.

In less severe cases, *Lachesis*, *Belladonna*, and *Mercury*, in alternation, will suffice, if the skin has become cooler, from the application of *Aconite*, but there still remain great soreness of the abdomen and intense thirst.

These medicines must be separately dissolved in water (twelve glob. to half a teacupful), and every two hours, alternately, a teaspoonful given, until three or four teaspoonfuls of each are taken, or until better.

But in the severest cases, when vomiting of fæces appears (*ileus miserere*),

*Opium* must be administered, every hour a dose (four glob.), and if not relieved within six, eight, or twelve hours,

*Plumbum*, in the same manner. It is in this stage where injections of an infusion of tobacco, or injecting tobacco smoke, is of the greatest benefit; also a lukewarm bath, to relax the whole system, and facilitate a decided reaction.

DIAGNOSIS of the *symptomatic* or *subacute* form. — This is always a concomitant symptom in typhus, or other malignant fevers; also in consumption of the bowels. The pain of the abdomen consists of a soreness, aggravated by hard pressure, and after cold drinks and indigestible food; the tongue is red, smooth, sometimes shining; loss of appetite; thirst variable; constant dryness in the mouth and throat; nausea and vomiting are more or less present, the more so, the nearer the inflamed portions of the intestines are to the stomach. When the lower parts are more inflamed, which is indicated by the pain in that region, diarrhea of a slimy mucus, often mixed with blood,

is present as in dysentery. This is particularly the case in the beginning. Consumption of the bowels, such as nursing females are sometimes subject to, see "Diseases of Females."

TREATMENT.—If this disease occurs in typhus or other malignant fevers, see their respective chapters. In the commencement of a typhus, if the pain in the abdomen is excessive, give a few doses of *Aconite*, every two or three hours one (four glob.); after which give *Bryonia* and *Rhus tox.* (in a half a teacupful of water, twelve glob. dissolved), every two hours, alternately, a teaspoonful. See "Typhus Fever."

If it resembles a bloody dysentery, after the use of *Aconite* as above, give *Belladonna* and *Mercury* in alternation (dissolve of each twelve glob. in half a teacupful of water), every hour a teaspoonful, until better; after which *Nitric acid* generally suits, particularly if the patient has already taken a good deal of calomel; or *Colocynth*, if bilious, *green* matter is vomited or discharged by the dysenteric operation. For the application of these medicines, and of *Sulphur*, which is important in this disease, see "Dysentery."

If worms are suspected to be the cause of the inflammation of the bowels, or are complicated with it, see "Worms."

If the disease resembles or runs into consumption of the bowels, see "Diarrhea" and "Diarrhea of lying-in Women."

The *chronic* form of this disease requires the same remedies, only administered not so often, because the symptoms are less urgent.

In all severe cases of this kind, it would be indispensably necessary to apply to a homœopathic physician, if one can be had.

DIET AND REGIMEN the same as in "Gastritis."

INFLAMMATION OF THE LIVER. (*Hepatitis.*)

DIAGNOSIS. — This inflammation differs according to its place in the liver.

*a.* When the surface and the convex, or upper part, of the liver is inflamed, there are in the right hypochondria sometimes stitches, at other times burning pains, often as violent as in pleurisy, shooting to the breast-bone, the right shoulder-blade increased by deep respiration, frequently accompanied with cough and vomiting; in this form the patient *can not lay on the right side.*

*b.* When the concave, or lower part, and the substance of the liver is inflamed, there is not so much pain, but more a feeling of pressing, heaviness, as of a bag lying in that region; the color of the eyes and face becomes yellowish, as if jaundice were present; bitter taste, and vomiting; in this form the patient can not lay on the *left side.*

In both forms general inflammatory fever is present more or less, and the region of the liver sensitive to pressure, and apparently bloated and hot. The first form may be mistaken for pleurisy, as the symptoms are very similar; but, as in such a case the treatment would be the same, no danger can result from such a mistake.

If hepatitis is not well treated, it has the tendency to pass into the chronic state, forming ulcerations, which either appear externally or communicate internally with the intestines or the lungs, causing inflammation of these organs; indurations, adhesion, gangrene are also issues of neglected hepatitis.

CAUSES. — Summer heat and hot climate predispose to this disease; also, the abuse of ardent spirits; beside, it is excited by external lesions, concussion of the brain, suddenly suppressed diarrhea or dysentery, hemorrhoidal congestion, or strong emetics or purgatives, the abuse of mer-

cury, the presence of stones in the gall-bladder, indigestion, violent passion or anger, particularly when immediately followed by drinking ardent spirits.

TREATMENT. — The following are the principal remedies necessary for the *acute* form of this disease: *Aconite*, *Chamomile*, *Bryonia*, *China*, *Belladonna*, *Mercury*, *Nux vomica*, *Pulsatilla*, *Sulphur*, *Lachesis*.

For the species *a.* are more suitable: *Aconite*, *Belladonna*, *Bryonia*, *Mercury*, *Lachesis*, *Sulphur*.

For the species *b.*, *Aconite*, *Chamomile*, *Mercury*, *Pulsatilla*, *China*, *Nux vomica*, *Sulphur*.

For *chronic* affections, such as enlargement and induration of the liver (liver complaint): *Nux vomica*, *Sulphur*.

(Other serious chronic diseases must be treated by a homœopathic physician.)

SYMPTOMATIC DETAIL. — *Aconite*. At the commencement of the treatment, when there is violent inflammatory fever, *shooting* pain in the region of the liver, tossing about, fear of death, and anguish; pain seems to be insupportable.

*Belladonna*. Pressing pains extending to the chest and shoulders; fullness in the pit of the stomach and around the short-ribs; difficult and anxious respiration; anxious tossing and sleeplessness; thirst, dryness of the mouth and throat; headache, with clouded sight, and vertigo, with fainting. Suitable after *Aconite*, and alternately with *Lachesis* or *Mercury*.

*Bryonia*. Pressing pains, with fullness in the hypochondria; *violent oppression of the chest*; rapid and anxious breathing; tongue yellowish and thickly coated, followed by fever and thirst; pains worse by movement; constipation. Suitable after *Aconite*, or alternately with *Mercury*.

*Mercury*. Patient can not lie long on the *right* side; pains are pressing and shooting; bitter taste in the mouth; thirst



and shivering ; skin and eyes yellowish. After it *Lachesis* and *Sulphur* are suitable.

*Lachesis*. If *Belladonna* or *Mercury* are insufficient, or when the patient is in the habit of drinking ardent spirits.

*Sulphur*. In every case of hepatitis, either at the end of the cure, or when the preceding remedies have produced no perceptible amelioration within a few days.

*Chamomile*. In hepatitis after taking cold ; when caused by violent anger, or when there are dull, pressing pains, *not aggravated by movement, respiration, or external pressure ; yellow color of the skin ; pressure in the stomach ; gastric bilious state of the stomach ; tongue coated ; bitter taste in the mouth ; fits of anguish.*

*Pulsatilla*. In hepatitis from indigestion, alternately with *China* ; nausea, bitter taste in the mouth, tongue coated ; oppression of the chest ; tension around the short-ribs ; *fits of anguish in the night, with loose, greenish, and slimy evacuations.*

*China*. After indigestion, or when there is shooting and pressing, swelling and hardness of the hepatic region ; pressing headache ; bitter taste in the mouth and yellowish coat on the tongue.

*Nux vomica*. After a fit of passion, or when there is shooting and beating under the short-ribs of the right side, excessively tender when touched ; nausea, with sour and bitter taste ; *fits of anguish ; shortness of breath, thirst, vertigo, and headache ; suitable for drunkards, and in chronic liver affections, previous to Sulphur.*

ADMINISTRATION. — Dissolve of the remedy selected twelve globules in half a teacupful of water, and give every two hours a teaspoonful, until three or four spoonfuls are taken, or until another remedy is indicated by the change of symptoms ; as the patient gets better, lengthen the intervals ; as soon as greatly relieved, discontinue all medi-

cine for forty-eight hours, when one dose (four glob.) of *Sulphur* may be administered to complete the cure.

In *chronic* hepatitis, give of the necessary remedy only once in twenty-four hours a dose (four glob.) for two or three days; then discontinue six or eight days before the next remedy is given.

DIET AND REGIMEN. — *No meat or meat soups of any kind*, but farinaceous substances—rice and oatmeal gruel, sweetened and made palatable by lemon-juice, lemonade, cold or warm, prunes stewed or boiled in gruels, baked apples, toasted bread, toast-water, cold water alone. Keep the patient quiet and comfortable as regards temperature.

#### JAUNDICE. (*Icterus*.)

DIAGNOSIS. — This disease, well known by the yellow color it imparts to all the white parts of the body, is not to be considered dangerous of itself, but becomes only so by neglect, when other serious disorders, hectic fever, dropsy, etc., may follow. In most cases it is the result of neglected liver diseases, or is caused by taking cold which affected the liver; also, by worms or gall-stones, obstructing the gall-duct, or is induced by excessive mental emotions. In jaundice the bowels become constipated, and if fæces pass, they are hard and whitish; the urine is orange colored, and the skin dry, or, if moist, it is from weakness, and the perspiration imparts a yellow color to the patient's linen.

TREATMENT. — The principal object must be to increase, by degrees, the healthy action of the skin; the patient, therefore, ought to keep warmly covered in bed in a warm room, and take, for the first three or four days, *Mercury*, morning, noon, and night a dose (four glob.); if not better at the expiration of four days, give *Hepar*, in the same manner; but if signs of amelioration appear, give it at longer inter-

vals ; this course generally removes the disease in eight or ten days, during which time the patient must keep in an equal, warm temperature and perspiration. In very obstinate cases *Lachesis*, *Nitric acid*, *Sulphur*, may be given afterward in the same manner.

If jaundice is complicated with fever, see "Inflammation of the Liver."

If it is caused by abuse of *Mercury*, see medicinal diseases, under the head of "*Mercury*."

DIET AND REGIMEN, as in "Hepatitis."

#### INFLAMMATION OF THE SPLEEN. (*Splenitis*.)

DIAGNOSIS.—Sharp, pressing, or shooting pains in the left side, below the short-ribs ; the spleen on pressure is painful, and when pressed upward excites cough and nausea in the stomach, sometimes vomiting of blood ; if fever is present, it is very high, with all the concomitant symptoms of inflammation of internal organs. The region of the spleen is enlarged and congested.

CAUSES.—Similar to those of inflammation of the liver, besides the abuse of *quinine* or *Peruvian bark* ; also, living in a marshy region causes congestion of the spleen, and consequently inflammation, under otherwise trifling circumstances.

TREATMENT.—We have an acute and chronic form. For the *acute*, *Aconite*, *Bryonia*, and *China* are the principal remedies, to be administered as directed in hepatitis.

If vomiting of blood is present, or sometimes even the blood is discharged by the stools, *Arnica*, *Rhus tox.*, and *Arsenic*, followed by *China*, are necessary, one remedy after the other, in the same manner as above.

For the *chronic* form, *enlargement*, *induration* of the spleen (fever cake), *Sulphur*, *Calcarea carb.*, *Ferrum*, *Lycopodium*, *Carbo veg.* are the principal remedies ; administration as directed in the chronic form of hepatitis ; but

it is strongly recommended to procure the aid of a skillful homœopathic physician for its treatment.

DIET AND REGIMEN, as in "Hepatitis."

### WORMS.

Every part of the system can, under favorable circumstances, create and sustain worms, or other imperfectly organized animals; thus, the skin, liver, lungs, kidneys, even the brain and eyes, in some animals, are favorable places for their generation, the species of animals varying according to the organ attacked. *Worms* are most frequently found in the intestinal canal; there are three kinds of worms met with, the *Ascaris*, or pinworms (*ascaris vermicularis*), *Mawworms* (*ascaris lumbricoides*), and the *Tapeworm* (*taenia*), of which there are two varieties:

a. The *broad one* (*taenia lata*), varying from three to twelve feet in length, seldom comes away entire, but in joints; and

b. The *solitary tapeworm* (*taenia solium*), with long and slender joints, sometimes reaching the length of thirty feet.

DIAGNOSIS.—It is frequently not an easy matter to decide positively on the presence of worms in the system, as most all those symptoms which could lead to such a conclusion are also met with in other diseases, save the one symptom, surest of all, that is, the passage of some worms, or undoubted pieces of them. This uncertainty in the diagnosis of the worm disease is no less annoying and embarrassing for an allopathic practitioner; as his prescriptions are dangerous and, in most cases, without good or decided result for the little patient, as with the worms their cause is not always expelled. Homœopathy, on account of its guiding principle, avoids either of the two dangerous issues, and, prescribing for the ostensible symp-

toms as they reflect the inward disease, cures and eradicates this surely, safely, and mildly. Merely expelling the worms, by remedies which have no *specific* bearing upon the worm disease, is of no essential use, and requires often enormous quantities of poisonous and dangerous stuff before one of the worms is sickened, killed, or expelled. Yes, I have seen the foundation of painful chronic diseases laid by the over-medication in presumed or real cases of worms; there are even cases where children have died under the operation of those drugs, which allopathists believe to be obliged to apply in such dangerous doses. The most irrational, nay, a real lottery business, is the use of those manifold nostrums, in the shape of lozenges, syrups, etc., which generally contain one or more specific anti-worm remedies; for instance, worm-seed, or pink-root, which in small and not offensive doses given would produce salutary effects, but in the combination of dangerous purgatives expose the little patients to death or misery; at least, the smallest number escape unhurt, most of them receiving injuries of the abdomen, the bad consequences of which impede not a little the future development of the child.

Worms, as such, are not dangerous. They have no teeth, and consequently can not eat children up alive. It is true, they can and will disturb the quiet and rest of a child, and make it pick at the nose, or cough, or cry out suddenly, or even have fits, etc.; yet all this is only produced by the worms having been made restless, by the presence of sickness in the system, which causes them to touch the walls of the intestines, usually avoided by them—living in and of the mass of mucus and other nutritive substances, with which the system had been indiscriminately, and in too large quantities burdened with. As soon as the system recovers from its own disease, either from cold or irregu-



larity of diet, the worms return to their usual, unobtrusive quiet again, giving no signs of their existence, nor causing alarm. Every person, more or less, has worms, without the least injury to the system. Neither is it absolutely the quantity of them, which is the most dangerous quality of this disease; a few worms can cause as many fits, or other spasmodic symptoms, as whole clusters of them. It depends altogether on the accompanying disease; and this it is which we will have to treat, before we can expect the worms to leave the system (which I have seen frequently, after a dose of the homœopathic preparation of *Sulphur*), or to be absorbed (digested and reduced to mucus, from which they originated.) This latter is the most common result of homœopathic medicines in this disease. In such cases, the children show signs of improvement, as soon as their abdomen becomes reduced in size and hardness; their complexion improves, they become more agile, grow better in height, and sleep quiet.

And this result is always attained by the judicious use of homœopathic medicines in this disease, in whatever form it may appear; for instance, as fits, diarrhea, scrofula., etc. Fits, occasioned by worms, are not in the least dangerous, and only become so by unwisely dealing with them.

The principal symptoms, by which the presence of worms in the system is indicated, are the following: the complexion of the patient is pale and changeable; livid circles around the eyes; accumulation of saliva in the mouth in the morning, and when not having taken nourishment for some time; irregular appetite; nausea; fetid breath; great hunger; picking and boring at the nose, with sneezing; abdomen enlarged, but not hard; frequent distressing pain in the stomach, in the region of the navel, which forces the patient to draw up the limbs against the abdomen, or lay on the stomach when going to sleep; pupils of



the *eye dilated*; bleeding at the nose; starting, as if from fright, during sleep, also grinding of the teeth; disposition to spasms, and unusual emaciation. The surest sign, however, is the discharge of worms, or parts of worms.

Signs of *tapeworm*, in particular, are: a sensation, as of something suddenly rising from the left side into the throat and falling back, or as if a lump in one or the other side was making an undulatory motion; dizziness, particularly in the morning, before eating.

Signs of *ascarides*, or *pin-worms*, especially, are: itching in the anus, more in the evening or at night; difficulty of making urine; tenesmus; apparent hemorrhoidal complaints; discharge of mucus from the rectum, bladder, or vagina (fluor albus.)

Signs of *maw-worm*, or *lumbricus*, particularly, are: all the above general symptoms of worms, but frequently stomach-ache, which is relieved after drinking sweet, cold milk.

### *Diseases caused by the Presence of Worms.*

Infancy is the age, when, on account of the general predominancy of nutrition, laxity and weakness by accumulation of mucus in the intestines preponderate; and consequently, those remote causes for the generation of worms, which may be favored even by circumstantial ones, such as flabby constitution of the child, damp habitation, and other epidemical and endemical influences.

This variety of the remote causes of worms gives rise to an equal variety of diseases, flowing from the same source (the worms), yet different, according to their exciting causes. Thus these diseases vary, from the most trifling illness, to the most severe colics, and dangerous fevers. The worms create, operating on the *nervous system* by sympathy, convulsions, all kinds of spasms, epilepsy, St. Vitus'

dance (*chorea*), somnambulism, periodical paralysis, insanity, fury.

On the *vascular* or *blood system*: fevers, congestions to bowels, hemorrhages, eruptions on the skin, appearing in irregular blotches of a scarlet color.

On the *reproductive system* (the intestines themselves): pains, spasms, increased secretion and accumulation of mucus, diarrhea, dysentery, worm colic.

TREATMENT. — The principal remedies for the diseases produced by worms on the *nervous system*, as above stated, are: *Ignatia*, *Cina* <sup>cc.</sup>, *Nux vomica*, *Hyosciamus*, *Belladonna*, *Spigelia*, *Sulphur*.

On the *vascular* or *blood system*: *Aconite*, *Belladonna*, *Hyosciamus*, *Cina*, *Ferrum*, *Mercury*, *Sulphur*, *Silicea*.

On the *reproductive system*, or the *intestines*: *Aconite*, *Belladonna*, *Cina* and *Cina* <sup>cc.</sup>, *Mercury*, *Sulphur*, *Spigelia*. (For the detail of these medicines, see below.)

Especially against *tape-worms*, the most suitable remedies are: *Mercury* and *Sulphur*, to be given every four or six days, alternately, a dose (six glob.), after which *Calcarea carb.*, *Graphites*, *Stannum*, may be given in the same way, if necessary. If not then relieved, apply to a homœopathic physician.

Against *ascarides* (*pin-worms*): *Aconite* and *Ignatia*, or *China* and *Mercury*, or *Calcarea carb.* and *Ferrum*, or *Cina* and *Sulphur*, alternately, every second or third evening a dose, until better. An injection every evening of an ounce of pure sweet oil (Olive oil), without any other admixture, is an excellent adjunct against these troublesome worms. The *ascarides* lodge never higher up in the abdomen than the rectum; therefore, can be expelled easily, or killed.

Against *maw-worms*, or worms in the intestinal canal, are generally appropriate: *Aconite*, *Belladonna*, *Hyosciamus*, *Mercury*, *Cina*, and *Cina* <sup>cc.</sup>, *Sulphur*, *Silicea*, *Spi-*

*gelia*, *Ferrum*, *Ignatia*, *Nux vomica*; but in particular diseases caused by maw-worms, as follows:

a. For *colic*, with *fever*: *Cina* <sup>cc.</sup>, *Aconite*, *Belladonna*, *Hyosciamus*, *Mercury*.

b. For *colic* with *convulsions*: *Cina* <sup>cc.</sup>, *Belladonna*, *Hyosciamus*.

c. For *colic* with *diarrhea* and voracious appetite: *Spigelia*.

d. For *fever*: *Aconite*, *Spigelia*, *Silicea*.

For *slow fever* in *scrofulous* children: *Silicea*.

SYMPTOMATIC DETAIL. — *Aconite*. In the commencement of all cases, where considerable fever exists; restlessness at night; irritability of temper; also, in cases without fever, when there is, from the presence of ascarides, continual itching and burning at the anus; and then alternately with *Ignatia*.

ADMINISTRATION. — In fever, every two or three hours a dose (infant children above one year of age three glob.), repeated two or three times, until the fever abates, when another remedy will be necessary. In cases of ascarides, alternately with *Ignatia*, every evening a dose (two or three glob.)

*Ignatia*. In pin-worms, or ascarides (see *Aconite*.) In maw-worms, when there are spasmodic twitchings in the muscles of the arms, as if the St. Vitus' dance was commencing. In this latter case, alternately with *Sulphur*, every evening a dose (two or three glob.)

*Sulphur*. At the end of every case of worm affection, after other remedies have either subdued the fever, or mitigated the nervous symptoms. Give every evening one dose (two or three glob.) for three evenings, and then wait from six to eight days, before another remedy is applied, or the same repeated, if necessary.

*Cina*. This is one of the most important remedies, par-

ticularly when there is boring at the nose, fever (after the application of *Aconite*); irritability; restlessness at night; desire for things which are rejected when offered; pale, bloated face, with livid circles around the eyes; craving for food, even after a meal; the child wants bread, potatoes, and other gross victuals; colic; griping; distention and hardness of the abdomen; discharge of *pin* and *large* worms; diarrhea; weakness of the limbs; the child can not stand any more on its feet; spasmodic movements in the limb.

Administration, see *Aconite*.

*Cina* <sup>cc.</sup>. When the children are very nervous, cry out in their sleep, or have *severe* colic pains, give every one or two hours a dose (two or three glob.), until better, or another remedy is necessary. Where children cry out in the night, groan, and are restless, give every evening only one dose (two or three glob.) until better.

*Spigelia*. In severe cases of worm colic, fever, and diarrhea, with craving appetite and chilliness, or where the symptoms appear regularly at the same time of the day.

Administration in acute cases the same as *Aconite* in fever. In chronic cases, every evening a dose (two or three glob.)

*Hyosciamus*. In worm fevers, when the patients are very nervous, restless, cry out, want to run off, and stool or urine passes off involuntarily.

Administration as in *Aconite*.

*Belladonna*. Worm colic after the administration of *Aconite*, when great pains remain in the stomach, which are relieved by lying on the bowels, starting in the sleep, sensibility to light, headache; on awaking, the child does not know where it is, etc. Particularly useful in scrofulous children, with distended abdomen and red cheeks, or where red blotches, like erysipelas, appear on different

parts of the body. In such cases, in alternation with *Silicea*.

Administration in *acute* and *chronic* cases, the same as *Aconite*.

*Mercury*. In *worm diarrhea*, swelling and hardness of the abdomen, and when there is unusual flow of saliva from the mouth.

Administration as in *Spigelia*, after which it is suitable.

*Nux vomica*. Worm symptoms with constipation (the child has frequent, but insufficient calls to stool), and irritability.

Administration as in *Mercury*.

*China*. In worm diarrhea, when it has already lasted some time; distention of the abdomen; debility; great appetite; pains in the stomach after every meal. Suits well in alternation with *Veratrum*, particularly when the stools are light colored.

Administration as in *Mercury*.

*Ferrum*. When there is vomiting and accumulation of watery fluid in the mouth.

*Silicea*. If a worm fever assumes the slow, chronic form, with or without diarrhea, particularly in scrofulous children.

Administration as in *Sulphur*.

DIET AND REGIMEN.—In worm fever, adhere to the diet recommended for common fever. In severe worm colic, give as a palliative, cold, sweet milk, from time to time. In diarrhea, avoid acids, fruits, and all indigestible articles; give, rather, soups of mutton or lamb, with rice. Avoid, in chronic worm diseases, heavy, gross nourishment, such as too much bread and butter, potatoes, heavy puddings, pies, raw or boiled vegetables; but meat-soups of any kind, meat, roasted or broiled, plenty of cold water

and milk, are good. Besides, bathing in cold water, and exercise in the open air. (See "Hydropathy.")

#### CONSTIPATION. COSTIVENESS.

This term must not always be taken in the absolute sense as a diseased state of the bowels; as there are persons who naturally have only every two or three days an evacuation—a habit which frequently promotes rather than disturbs general health, if the latter is not interfered with by an unwise use of medicines, such as taking pills or aperients of a drastic or saline nature. As a general rule, we can consider it the normal state of the bowels, to have one evacuation every day. This ought to become a fixed rule and habit, as it frequently regulates costiveness of the bowels, without the use of any medicine.

Some occupations of life predispose to costiveness, particularly those which allow but little exercise. Persons, under such circumstances, must abandon these occupations, if possible, for a length of time, take a great deal of exercise in the open air, change the diet (see below), and make use of cold water, externally and internally, in large quantities (see "Hydropathy.") Nothing is more injurious than the continued use of medicinal aperients. They render the bowels, in time, more costive and torpid. The best expedient for facilitating an immediate action on the bowels, if wanted, is an *injection*, either of *cold water*, if the person otherwise is in health or the patient prefers it, or lukewarm water, sweet oil, or lard, and a small piece of castile-soap dissolved in it, if the constipation has been very obstinate, or a quick evacuation is wanted, as in children, when they have fits, etc.

In most cases of acute diseases, constipation is only one of the many symptoms constituting the disease—not the



disease itself; because it disappears as soon as the whole complex of symptoms has been taken away.\*

If, however, constipation is the only or principal symptom in a disease, as in miserere, lead colic, etc., we have remedies which combat it more safely and successfully than all the aperients commonly used. Cases of this kind, however, must be treated by experienced homœopathic physicians.

**TREATMENT.** — We will now give first a tabular view of the causes and complications of constipations, with their principal remedies, the detailed symptoms of which follow below. These must be consulted before a selection is made.

*Constipation in general, or a disposition for it: Bryonia, Nux vom., Lachesis, Lycopodium, Sulphur, Sepia.*

*In persons who lead a sedentary life: Bryonia, Nux vom., Sulphur, Opium, Platina, Lycopodium.*

*In drunkards: Lachesis, Nux vom., Opium, Sulphur.*

*In old persons, or those of weakly constitutions, where constipation often alternates with diarrhea: Antimon. crud., Conium, Opium, Phosphorus.*

---

\* One of the most striking cases of this kind, showing in a high degree the truth of the above statement, presented itself to me a few years ago, in a lady who had been persuaded to take castor oil, three days after her confinement, although in former cases of this kind she always had suffered very much after its use, from irritation of the intestines. After the administration of the castor oil, a violent and painful diarrhea appeared, threatening fearful consequences for the life of the patient. This was, however, soon removed, by the exhibition of *China, Mercury, and Pulsatilla*, and her health decidedly improved, from day to day, her appetite and strength returned, and she moved about as if she was well; yet there was no disposition for evacuating, until on the *twenty-second day* after the last evacuation, when it appeared, without any artificial means, in a perfectly healthy condition. During the eight days previous to it, the patient had been allowed a good nourishing diet, and enough of it, too.

I have frequently had cases, where patients had no evacuations for eight, ten, or twelve days; but I never have experienced their suspension for twenty-two days, except in the above case.

In *pregnant women* : *Nux vom.*, *Opium*, *Sepia*.

In *nursing infants* : *Bryonia*, *Nux vom.*, *Opium*, *Sulphur*.

In *lying-in women* : *Bryonia*, *Nux vom.*

In *consumptives* : *Stannum*, *Silicea*, *Sulphur*.

By *traveling in a carriage* : *Platina*, *Opium*.

During *sea-voyages* : *Cocculus*, *Nux vom.*, *Silicea*.

From *poisoning by lead* : See "*Painter's Colic*."

From *abuse of purging medicines* : *Opium*, *Nux vomica*, *Lachesis*.

From *abuse of quinine* : *Pulsatilla*, *Carbo veg.*, *Veratrum*.

SYMPTOMATIC DETAIL. — *Bryonia*. In all recent cases of constipation during *fevers* or *bilious attacks*, where headache, chilliness and rheumatic pains predominate ; particularly, also, in *warm weather* ; the patient is of a nervous, irritable temperament. After it, or in alternation with it, *Nux vomica*.

*Nux vomica*. Is of the most frequent application in *hypochondriacal persons*, or those suffering from *piles* ; also, after derangement of the stomach, when there is *frequent but ineffectual effort* to evacuate, or a feeling as if the *anus were closed or contracted* ; ill humor, fullness of the head, can not bear the clothes tight around the waist. After it, or in alternation with it, *Sulphur*.

*Sulphur*. In all cases of *habitual costiveness*, or where *Nux vomica* was insufficient.

*Opium*. In all cases where there is *great torpidity* of the bowels, and a feeling as if the *anus were closed*, but *without* the frequent and ineffectual efforts to evacuate, as is the case in *Nux vomica* ; the patient only feels full in his bowels, has congestion to the head, with dark redness of the face, slow and full pulse ; this remedy is of frequent application in children, old persons, pregnant women, drunkards, and hypochondriacal persons.

*Sepia.* In cases where there is, beside constipation, *flashes of heat, sick-headache*, a pressing down, particularly in pregnant women, or where *Nux vomica*, *Opium*, and *Sulphur* have been insufficient.

*Platina.* In nervous females and weakly hypochondriacal persons, where the stool only appears in small pieces, followed by tenesmus and tingling in the anus; shuddering, and a sensation of weakness in the abdomen; sometimes, also, a constrictive pain in the abdomen, and ineffectual desire to evacuate. Compare *Nux vomica*.

*Lachesis.* A very useful remedy in obstinate constipations during fevers, particularly where abscesses of internal organs are existing or presumed to exist, as in the liver, mesenteric glands, etc., or in alternation with

*Lycopodium.* Where an obstinate disposition to constipation is present, from constitutional causes, such as scrofula, which had fallen on the bowels after it disappeared from the skin, etc.; if, in such cases, these two remedies remove the costiveness, but the disease attacks the lungs and produces cough, which is worse when a part of the body gets cool by exposure, give *Hepar sulph.* See “*Dyspepsia.*”

*Antimon. crud.* In cases where *constipation* and *diarrhea* alternate, if one ceases the other sets in; the stools are never regular.

*Phosphorus.* If *Antimon.* is insufficient, or when the patient is of a consumptive habit or has the consumption.

*Stannum.* If *Phosphorus* does not suffice, or when there is frequent but ineffectual desire to evacuate, particularly in consumptives.

*Silicea.* Same as *Stannum*, when this is insufficient, or in children whose stomachs are distended and hard, indicating worms or swollen glands; also, in dyspeptic persons, with heart-burn, sour taste in the mouth, etc.

*Cocculus.* See "Sea-sickness" and "Colic."

*Conium.* For *old* and *weakly* persons, who are troubled with frequent but ineffectual efforts to evacuate (after *Nux vomica*); also, in scrofulous children, where the mesenteric glands are swollen (after *Silicea*), particularly accompanied with soreness of the abdomen.

*Pulsatilla.* In similar cases, where *Nux vomica* would suit, but the temperament is mild and phlegmatic; or after derangement of the stomach by fat food, the patient appearing morose and silent; also, after abuse of *quinine*, either with or without the return of the fever and ague, and then in alternation with,

*Carbo veg.* This remedy, particularly, when persons have become very much debilitated, complain of *rheumatism*, *piles*, or have the *consumption*.

*Veratrum.* After *Opium*, when there is great torpor in the bowels, deficiency of expulsive power; or where the abdomen becomes *very tender to the touch*, (and then in alternation with *Arsenic*); see "Bilious gastric Fevers;" also, after abuse of *quinine*, when there is great congestion to the head and hypochondriacal feeling.

ADMINISTRATION. — In cases where constipation exists as the only troublesome symptom, give the medicine selected in quick succession and large doses; for instance, *Nux vomica*, every three or four hours a dose (six to eight glob.) for one or two days, until relief is obtained, or another remedy is chosen; or *Lachesis* and *Lycopodium*, in alternation, three times a day a dose (six glob.); but where constipation is habitual or constitutional, the remedies must be administered at longer intervals; for instance, *Nux vomica*, every evening a dose, (four glob.) for three or four days; then discontinue an equal length of time without taking medicine, after which either repeat it or select another remedy, if necessary, which in most cases will be *Sulphur*.

DIET AND REGIMEN. — These are important agents in the cure of constipation ; first of all is the habit of attending to the calls of nature carefully and regularly, at a certain time, say after breakfast, though there should be no actual result from it ; secondly, the careful avoidance of all nourishment which is confining in its character, such as salted meats, cheese, rice, wheaten flour, except these things are mixed in a sufficient quantity with their opposites ; a loosening quality have : fresh meat, and soups made of it, green vegetables, and fruits, except almonds or nuts ; but the chief promoter of a healthy action on the bowels is the frequent and liberal use of cold water, internally and externally, with bodily exercise in the open air ; see “Hydrophathy.” The cold and ice water can be drank freely during and after a meal, without the least inconvenience, and ought to be preferred to the use of wines, brandies, tea or coffee.

#### DIARRHEA. LOOSENESS OF THE BOWELS.

This complaint varies so much in regard to time of duration, form, and cause, that we are obliged to give here a general view of its varieties. It is either recent, lasting only for a few hours or days, or is chronic, lasting for months or years. The discharge itself may vary as regards consistency or character, either watery fluid, fecal, slimy, mattery, bilious, or bloody (see “Dysentery.”) Sometimes a diarrhea is dangerous ; at other times salutary, as in biliousness or after indigestion (see these articles.) In general, however, the popular idea of the harmless or beneficial nature of diarrheas, which the common practice of the old school of medicine has freely propagated, must be restricted, in so far as any diarrhea is of itself an indication of a diseased state of the intestines, and no disease ought to be created or wished for where we



can do without it. It has been, and is yet, to a great extent, a common expression that a good cleansing of the bowels is necessary, etc.; now, these ideas, when carried out in practice, lead sometimes to awful consequences, while the good which they may occasionally produce might have been obtained by mild and rational means. See "Constipation" and "Biliousness." On the other hand, it is just as dangerous to stop a diarrhea suddenly by violent means, such as brandy, paregoric, or astringents generally, as thereby the disease is not cured, but merely its natural outlet stopped; as if an ulcer could be cured by covering it over to confine and hide the matter, which it has to discharge. It is always fortunate if such a diarrhea reappears.

TREATMENT. — We will now give first a general view of the varieties of diarrhea, as regards causes, form, and appearances, with their principal remedies, whose detailed symptoms may be found below, and must be consulted before a choice is made.

*Diarrhea without pain* : *China, Phosphoric acid, Ferrum, Secale.*

With *gripping or colic pain* : *Mercury, Colocynth, Chamomile, Rheum, Pulsatilla, Bryonia, Rhus tox., Arsenic, Sulphur.* (See "Bilious Colic.")

With *tenesmus*, during or after the discharge, in the anus : *Ipecac., Mercury, Belladonna, Nux vomica, Sulphur.* (See "Dysentery.")

With *vomiting* : *Ipecac., Veratrum, Arsenic, Rhus tox.* (See "Cholera.")

With *debility* : *China, Arsenic, Ipecac., Veratrum, Secale, Phosphorus.* (See "Cholera Morbus" and "Asiatic Cholera.")

Of *mucus and blood* : See "Dysentery."



Of bile : See “Bilious gastric Derangement ;” beside, *Bryonia*, *Rhus tox.*, *Arsenic*, *Mercury*, *Pulsatilla*.

Of watery (rice-water) liquid : see “Cholera Asiatica.”

After eruptive fevers, such as small-pox, scarlet, measles, etc. : *Pulsatilla*, *China*, *Phosphoric acid*, *Arsenic*, *Sulphur*.

After sudden mental emotions : after joy : *Coffea*, *Aconite*, *Pulsatilla* ; after fright : *Chamomile*, *Veratrum* ; fear of cholera or other dangers : *Chamomile* ; after grief : *Phosphoric acid*, *Ignatia* ; after anger : *Colocynth*, *Chamomile*, *Nux vomica*. (See “Mental Diseases.”)

After indigestion : from partaking of milk : *Bryonia*, *Sulphur* ; of acids or fruits : *Lachesis*, *Pulsatilla*, *Arsenic* ; of other indigestible substances, fat meats, etc. : *China*, *Pulsatilla*, coffee without milk. (See “Indigestion.”)

After the abuse of drugs : of *Mercury* or calomel : *Hepar*, *Pulsatilla*, *China*, *Carbo veg.*, *Nitric acid* ; of *magnesia* : *Pulsatilla*, *Rheum* ; of *rhubarb* : *Pulsatilla*, *Mercury*, *Colocynth* ; of tobacco : *Pulsatilla*, *Chamomile*, *Hepar*, *Veratrum* ; of quinine : *Pulsatilla*, *Carbo veg.*

After cold or iced drinks : *Pulsatilla*, *Carbo veg.*, *Arsenic*.

After taking cold, cold on the bowels : *Dulcamara*, *Chamomile*.

In summer : *Bryonia*, *Mercury*, *Arsenic*.

In autumn and spring, during the changeable weather : *Bryonia*, *Rhus tox.*, *Dulcamara*, *Carbo veg.*

In weak and exhausted persons : see “With Debility.”

In consumptive persons : *China*, *Phosphorus*, *Calcarea*, *Ferrum*. (See “Consumption.”)

In chronic diarrhea, of a scrofulous character : *Iodine*, *Phosphorus*.

In *aged persons* : *Secale*, *Antimon. crud.*, *Bryonia*, *Phosphorus*.

In *pregnant and lying-in females* : *Phosphorus*, *Antimon. crud.*, *Dulcamara*, *Pulsatilla*, *Sepia*, *Rheum*. (See "Female Diseases.")

In *children* — by *worms*, *teething*, and *scrofula* : see these articles.

**SYMPTOMATIC DETAIL.** — *Antimon. crud.* Watery diarrhea, from disordered stomach, with foul tongue and nausea.

*Dulcamara*. In *most all diarrheas from taking cold*, or when there are greenish, or yellowish, slimy, or watery evacuations, particularly at *night*, with colic and griping pains in the region of the navel; great thirst, *nausea*, debility, and restlessness.

*Colocynth*. *Bilious* or *watery* diarrhea, with *severe cramp-like pains* (see "Bilious Colic" and "Dysentery"); or after *Chamomile*, when caused by anger or vexation.

*Chamomile*. *Bilious*, *watery*, or *slimy* diarrhea, of a *green (grass-green)* or *yellowish white*, curled appearance, like *scrambled eggs*, with or without *vomiting of bile* (see "Biliousness"); griping, tearing colic; distention and hardness of the abdomen; fullness in the pit of the stomach, and under the short-ribs; bitter taste in the mouth; this medicine is particularly useful in diarrheas from cold, fear, anger, vexation, and in those of infants (see "Diseases of Children"), when they draw up the legs, cry, toss about, want to be carried, etc.

*Rheum*. *Sour smelling* evacuations, with ineffectual effort *before* and *after*, at the same time *contractive colic* in the abdomen, and *shuddering* when evacuating; grayish brown evacuations, *mixed with mucus*; very useful in diarrheas of *infants*, particularly when they cry, toss about (see *Chamomile*), and in diarrheas of *lying-in women*.

*Ipecac*. Diarrhea of *fomented* evacuations, greenish,

yellowish, putrid, *bloody*, or *slimy*; dysenteric, when there are discharges of white flakes, and great tenesmus (see this article), or diarrhea with vomiting (see “Cholera Infantum,” “Cholera Morbus,” and “Asiatic Cholera.”)

*Veratrum*. In all diarrheas resembling the different varieties of *cholera* (see these articles); or where there is great *debility* or *gripping pains*, as if *knives were cutting the intestines*, which happens often in fevers.

*Secale*. In *cholera*, or when *painless* evacuations are attended by *great weakness*; in *old people*, particularly *females*, where drowsiness ensues, in alternation with *Belladonna*. (See “Cholera Asiatica.”)

*Carbo veg*. In *cholera* (see this article), or where there are *thin, light-colored* evacuations, of a *putrid* or *fetid* smell, particularly in *children*; or after taking cold in wet weather; also, after abuse of *quinine*, in alternation with *Pulsatilla*.

*Arsenic*. *Burning* evacuations, with *severe colic pains*, alternately with *Veratrum*; or *fetid, putrid, indigested* diarrhea, alternately with *Carbo veg.*; or *watery, slimy, burning* evacuations, principally after *midnight*, when *Dulcamara* or *China* have afforded no relief; or in *cholera* (see this article.)

*Bryonia*. Diarrhea from the heat of summer (summer complaint of children), or where *bilious rheumatic* (see this article) symptoms accompany the bowel complaint, alternately with *Rhus tox*.

*Rhus tox*. In *cholera infantum* (see this article); or in bilious rheumatic disorders. See *Bryonia*.

*Pulsatilla*. In all diarrheas from indigestion or disordered stomach, where, after some gripping pain, *watery, green, or bilious slimy* stools appear, particularly at *night*; sometimes the color changes, frequently from yellow to white, or the discharge is mixed with blood; bitter taste

in the mouth, and foul tongue ; in diarrhea from abuse of *quinine* and *calomel*.

*Mercury*. In critical diarrhea after fevers or bilious states of the stomach, when it threatens to become too debilitating and irritating ; this critical diarrhea generally commences with griping pain and rumbling of the bowels, after which a copious, bilious, slimy, and frothy evacuation of a very fetid smell follows, with signal relief ; yet this may repeat in a short time after, and finally, if not relieved by *Mercury*, run into a dysentery. *Mercury* suits for almost any diarrhea when accompanied with griping in the bowels before, and tenesmus or burning in the anus after the discharge.

*Nitric acid*. Diarrhea after taking too much calomel or mercury, particularly when the stools are *bloody*, with tenesmus at the anus, sometimes of a fetid smell.

*Nux vomica*. *Frequent but scanty evacuations* of watery and greenish, lightish substance, with colic and tenesmus, and then in alternation with *Mercury*.

*Ignatia*. Diarrhea of a bloody slime, with rumbling in the bowels, or after continued grief.

*Belladonna*. Frequent, small evacuations of mucus, sometimes with fever heat in the head, pressing down (see "Dysentery") ; or in those congestions to the head which follow severe diarrheas, or accompany them in old persons, and then in alternation with *Secale*. (See this remedy.)

*Hepar sulph*. In diarrheas of children, when they smell sour, are lightish, greenish, yellowish, bloody, slimy, and like dysentery ; also, in those of nursing women. (See "Diseases of Females.")

*China*. Diarrhea after *eating* (*lienteria*), or in the *night* ; also, when it is very debilitating, with rumbling in the abdomen. (See "Cholerina.")

*Phosphoric acid.* Diarrhea without pain, after *grief* and distress of mind; or when it passes involuntarily. (See "*Cholerina.*")

*Ferrum.* Painless evacuations after *eating* and *drinking*, or discharges in the *night*, with griping, flatulency, thirst, pain in the stomach, back, and anus. (See "*Ascarides.*")

*Hyoscyamus.* Watery diarrhea; *involuntary discharges*, particularly in the *night*.

*Sulphur.* In most cases where other remedies seem to have failed, particularly when there are *frequent stools* at *night*, with *griping*, of a slimy, watery, frothy substance, and of a sour and fetid smell, or bloody; also, if a diarrhea returns easy on taking the least cold.

*Phosphorus.* In chronic diarrhea without pain, particularly in the diarrhea of consumptives.

*Iodine.* In chronic diarrhea of a whitish color, in scrofulous persons, and which will not yield to any other remedy.

ADMINISTRATION. — In *recent* cases dissolve twelve glob. of the selected medicine in half a teacupful of water and take after every evacuation a teaspoonful until better, or another remedy is necessary; children half the quantity. In *chronic* cases take every evening and morning a teaspoonful of such a mixture, or if taken in the dry state three or four glob. at a dose.

DIET AND REGIMEN. — Acids or acidulated drinks, fresh vegetables, and fresh meat and meat soups are forbidden; allowed are, rice, toasted bread, boiled milk, and gruels; no coffee and spiritous liquors. Cold water can be drank, if it is otherwise not injurious. Keep quiet as much as possible, even to lying down.

#### DYSENTERY. BLOODY FLUX.

DIAGNOSIS. — This complaint is, properly speaking, not a diarrhea; as no fæces are discharged, but only mucus

and blood, accompanied by constant urging to go to stool, violent tenesmus in the anus and rectum, or severe pains in the abdomen, if the disease is located higher up in the intestines. As long as this irritation or inflammation of the mucous membrane of the intestines lasts, no fæces are discharged; but when these make their appearance, even mixed with bloody mucus, the dysentery may be said to be at an end, or at least ameliorated.

Its causes and course vary—sometimes it appears suddenly, sometimes is preceded by diarrhea and other gastric rheumatic symptoms.

We distinguish a *white* dysentery, where only *light-colored* mucus (scrapings of the intestines) are discharged; this is soon followed by the *bloody* dysentery, when the blood comes from the highly-inflamed mucous membrane. In such cases, more or less fever is always present. If the patient is relieved, the fever subsides, and gradually with it the pains, and frequent urgency of the calls, the skin becomes moist, and sleep and rest ensues. Death may come from gangrene and exhaustion, but happens very seldom, under a judicious homœopathic treatment. Yet such severe cases of dysentery ought to be treated by a skillful homœopathic physician. The causes of this disease may be local, by irritating the intestines directly in eating and drinking, or general, by suppression of perspiration (hot days and cool nights promote this cause), or by an epidemic influence, which affects in a similar manner the mucous membrane of the intestines, as the influenza-miasm attacks the nose and bronchia; also, low, marshy regions, where intermittents prevail, incline to the propagation of dysentery, which is sometimes very fatal at such places.

TREATMENT.—Principal remedies are: *Aconite*, *Belladonna*, *Mercury*, *Colocynth*, *Ipecac.*, *Nux vom.*, *Bryonia*, *Rhus*, *Sulphur*, *Nitric acid*, *Carbo veg.*, *Arsenic*.



*Aconite.* In all cases at the commencement, when there is fever, pain in the head, neck, and shoulders, heat and thirst alternate with shivering. This is particularly the case when we have cold nights and hot days.

*Belladonna.* After *Aconite*, or in the beginning, when there is dryness in the mouth and throat, tenderness of the abdomen, and blood passing with the mucus; constant feeling of bearing down.

*Mercury.* This is by far the most important medicine in this disease, and beneficial in all stages, but particularly when there is *violent tenesmus* before and after the evacuations, as if the *intestines would force themselves out*, and yet nothing but *pure blood*, sometimes streaked with *white mucus* or *greenish matter* appears; at other times the evacuation resembles scrambled eggs. Concomitant symptoms are: violent colic, *nausea*, *shivering* and shuddering, cold perspiration on the face, great exhaustion and trembling of the limbs. (It alternates well with *Aconite* or *Belladonna*, if there is fever, and with *Colocynth*, if the discharges are mixed with green, bilious matter.)

*Colocynth.* After *Mercury*, the principal medicine, when the colic pain appears periodically, and is excessive; the discharges are mixed with green matter or lumps. In such cases, alternate with *Mercury*, if this remedy had not been given already.

*Ipecac.* Is also of great benefit in this disease, when it occurs in the fall, or when the mucus or slime appears first, afterward only mixed with blood. After, or in alternation with it, give *Colocynth*, if necessary.

*Nux vomica.* *Small, frequent evacuations* of *bloody slime*, with heat and thirst, mostly in summer; alternately with *Mercury*, when there is great *downward forcing*.

*Bryonia* and *Rhus*. In alternation, under the same conditions as stated in the article on "Diarrhea."

*Sulphur.* In the most critical cases, where no other remedy seems to afford relief; after *Mercury*. It must be allowed to have its effect at least thirty-six hours, before another remedy is given.

*China.* Dysentery in *marshy* countries, or when the discharges are *very offensive*.

*Nitric acid.* One of the best remedies, after *Belladonna* and *Mercury* have been given without effect, particularly when the discharge is either mucus without blood or blood without mucus; in either case, however, the tenesmus which follows, is violent.

*Carbo veg.* and *Arsenic.* In alternation, if the discharges become putrid in smell and involuntary; stupor ensues; red or bluish spots appear on the skin; the patient is very weak, and his breath begins to be cold; the pains in the bowels are *burning*. *China* may follow these two remedies, if the putrid smell of the discharges will not change.

ADMINISTRATION. — Dissolve twelve glob. of the selected medicine in half a teacupful of water, and give every hour or half hour a teaspoonful, or, if the discharges are not so frequent, after every discharge a teaspoonful, until six, eight, or ten teaspoonfuls are given, when the medicine must be omitted for a couple of hours, to await its effect. If then necessary, either repeat the same or select another remedy, if the symptoms have increased. But as soon as an amelioration takes place, discontinue all medicine, even if there is yet frequency of stool and violent pain at times. If very restless at night, give *Belladonna* <sup>cc.</sup> and *Coffea* <sup>cc.</sup>, every hour a dose (four glob.) for four hours.

If the disease becomes more *chronic*, give the medicine at longer intervals, say three times a day, finally only once a day. Cold water injections are of the greatest benefit, if the patient can bear them, and the dysentery is in its inflammatory stage.

DIET AND REGIMEN. — The patient ought to lay in or on the bed constantly, even during convalescence, if possible, well covered, but not too much heated. For food, if wanted, toasted bread, softened in water or black tea; for drink, toast-water, or small quantities of warm lemonade, in the inflammatory period, and when not taking *Aconite* at the time.

Gruels of oatmeal, rice flour, or farina, are allowed. Meats or soups of meat, eggs, and all animal food, are strictly forbidden, even during convalescence. Wine and spirits are real poison, in this disease, and long after it.

*After diseases.* — If *dropsy*, *paralysis*, or *rheumatism* follow the dysentery, when it has either been treated badly, or was suppressed by allopathic remedies, see these articles, or send for a homœopathic physician in time.

#### CHOLERA MORBUS.

This disease, well known under the above name, occurs mostly during the summer, in all parts of the country, north and south, resembling in its appearance the much dreaded Asiatic cholera, from which, however, it entirely differs, in character and result. (See "*Cholera Asiatica.*")

DIAGNOSIS. — Cholera morbus generally attacks suddenly, without any precursory symptoms, with vomiting and purging, accompanied with pains in the stomach and bowels, anxiety and tenesmus; sometimes the vomiting and urgency to go to stool are very violent and incessant, at other times less so. In very violent cases, exhaustion soon follows, pulse becomes small, scarcely perceptible, extremities cold; finally spasms and convulsions appear.

At first, the discharged substances consist of the contents of the stomach and of bile; afterward of a watery, lymphatic liquid; and finally nothing is thrown up; when violent retching and gagging, with urgency to go to stool, remains. The discharges from the bowels consist first of

fæces; afterward of a watery, bilious liquid; each stool preceded and accompanied with violent, burning, cutting colic, especially in the region of the navel.

CAUSES. — Intense heat in the summer; irritable temperament; teething in children (see “Cholera Infantum”); sudden suppression of habitual discharges, such as menstruation or diarrheas, or of cutaneous eruptions and gout; errors in diet, such as eating unripe, watery fruit, or drinking acrid, sour liquids, beer, etc.

TREATMENT. — The principal remedies in this disease are: *Chamomile*, *Ipecac.*, *Arsenic*, *Colocynth*, *Dulcamara*, *China*, *Veratrum*.

In the beginning, give *Ipecac.* (administration as below); if not better within half an hour, give *Veratrum* and *Arsenic*, in alternation, in the same manner; but if this treatment does not succeed within one and a half or two hours, select from among the remedies described below, one which is more homœopathically suitable, and give it as stated under “Administration.”

SYMPTOMATIC DETAIL. — *Chamomile*. In the beginning of the disease, or when the tongue is coated yellowish; colic in the region of the navel; pressing pains from the stomach up to the heart; great anguish; cramps in the calves of the legs; watery diarrhea and sour vomiting; particularly after fits of passion, and with an irritable temperament.

*Ipecac.* When vomiting predominates, alternately with painful diarrhea. It is generally *always* given in the beginning of the disease, in alternation with *Veratrum*.

*Arsenic.* Violent vomiting, and diarrhea of watery, bilious or slimy, greenish, brownish, or blackish substances, with severe pains in the abdomen, great restlessness, sudden prostration, cold extremities, and clammy sweat. In alternation with *Veratrum*.

*Veratrum.* For the same symptoms as those mentioned under *Arsenic*, only that the pains in the bowels, especially around the navel, are *violent, cutting* as from *knives*; also, pains or cramp in the calves of the legs; *prostration* is *very great*. See "*Ipecac.*" and "*Arsenic.*"

*Colocynth.* *Green* vomiting, with *violent colic*; the stools at first greenish and bilious, afterward become more colorless and watery.

*Dulcamara.* In cholera morbus caused by iced drinks or exposure to sudden changes of the temperature; frequent greenish stools, with bilious vomitings; great debility and severe pains in the abdomen.

*China.* Cholera morbus from indigestion; vomiting and diarrhea after eating ever so little (*lienteria*); painful eructation, with oppression on the chest; rumbling in the bowels; flatulency.

ADMINISTRATION. — Dissolve of the selected remedy twelve globules in half a teacupful of water, and give every ten or fifteen minutes a teaspoonful, until four teaspoonfuls are taken, or another remedy is necessary, which must be prepared and given in the same manner.

Externally, apply nothing but a mustard poultice on the stomach, and heated bricks, and warming bottles to the feet.

DIET AND REGIMEN. — For a drink, thin oatmeal gruel or cold water; if wanted. After the diarrhea has ceased, and reaction has taken place, and if the patient is feverish, give gruels with lemon-juice. In general, treat as stated in "*Bilious Fever.*"

Besides, consult the article on "*Asiatic Cholera.*"

#### ASIATIC CHOLERA. (*Cholera Asiatica.*)

This disease appeared hitherto among us as an epidemic. Before it, the star of Allopathy grew pale, and is ever since



on its downward course ; with it Homœopathy battled most successfully, and ever since her banner floats unfurled in every clime.

DIAGNOSIS. — Sudden prostration of strength ; diarrhea of watery substances (rice-water discharges), without smell or color ; similar, profuse vomitings, which follow each other quickly. Although these evacuations may give the patient a little relief, it is only momentary ; his prostration increases constantly until cramps appear in his feet, hands, limbs, and arms, sometimes in the bowels and breast ; insatiable thirst, with a burning sensation under the sternum (in the region of the heart.) Agonizing restlessness precedes the last stage—that of collapse, in which, to the above symptoms are added, cessation of circulation of the blood in the extremities, which assume an icy coldness, while the heart palpitates violently, no pulse is perceptible ; tongue and breath grow cold, agonizing oppression in breathing ; hollow, hoarse, and shrill voice ; blue lips and nails ; livid countenance ; sunken eyes ; shriveled skin on hands and feet. During all this time, the secretion of urine is stopped. Finally, the skin of the patient is covered with a cold, clammy perspiration. Up to this time, the patient was perfectly conscious, although feeble ; but now he sinks into stupor, with bloodshot upturned eyes, and expires.

Although Asiatic cholera presents itself in the form of a diarrhea, yet it is not such ; because no fæcal matter is found in the discharges ; its nature consists in a decomposition of the blood, produced by an unknown agent (miasm or animalcula), and aided by mental and physical debilities, such as fear, care, overpowering sympathy, weakness of body and mind by night-watching, exposure, etc. This decomposition of the blood in its solid and liquid parts (coagulum and lymph) may be the work of days, hours,



or minutes. Generally, however, it commences in the early morning hours, at which time most of the cholera attacks begin. The watery part of the blood (lymph) separated from the solid, is emitted in the thoracic and intestine lacteal duct, and emptied the absorbing vessels of the intestines in the intestinal canal, and from thence into the stomach, whence it is ejected in violent gushes, or runs from the bowels in streams. Thus we see that the first appearances of *cholera asiatica* are those of a hemorrhage of the white part of the blood—a view with which the consequent symptoms in the following stages of the disease can easily be harmonized. The cramps in the muscles of the extremities and bowels are caused by the cessation of the action of arterial blood on the muscles, and cease immediately, as soon as the arterial irritation reaches them again. It would carry us too far, to explain all the following symptoms of the disease; nor would it be of any practical use. This, however, we must state yet, that in that form of cholera Asiatica, called the *sicca* or dry cholera, where no discharge from the stomach and bowels takes place, this disagreement with the above theory is only apparent; because, by the suddenness of the attack, the decomposition of the blood is rendered so general and great at once, that the cramps follow immediately, closing the absorbing vessels in the intestines, before the lymph had time to escape.

PREVENTION. — During a cholera epidemic, a person must not give way to fear and despair, which debilitates the system too much, allowing, thereby, an easier entrance to the enemy. The best remedy against the weakening thoughts is a constant and vigorous activity of body and mind, which draws our attention away from disease and death to our duties and objects of life. Necessary, above all, is regularity in all our actions, in sleeping and waking; modera-

tion, but sufficiency in eating and drinking. The most substantial nourishment is the best, such as beef, mutton, potatoes and bread; the best drink is cold water, or ice water, which, with dry bread in sufficient quantity, is the best calculated to appease that craving and gnawing appetite, and the rumbling of the bowels which, in time of cholera, almost every one feels daily. A flannel bandage, even in summer, covering stomach and bowels, is strongly recommended. To be avoided are: over-exertions of body or mind, grief, anger, fear, as well as great muscular labor, running, walking to exhaustion, and sexual excesses. Besides these precautionary measures, every one ought to have a firm reliance in the will of God, who has placed within our reach a safe and sure remedy against this disease.

The best medicine for prevention is *Sulphur*, which, in the thirtieth dilution, may be taken every week twice, six globules each time, in the evening. This remedy, more than any other, destroys within the system the liability of attack, by diminishing the psoric diathesis, more or less inherent in every one.

*Cure of an attack of Asiatic Cholera.*—Not every diarrhea, in time of cholera, is real Asiatic cholera, yet it predisposes to it; and we must, therefore, *immediately* and energetically attend to it. The first and most imperious duty in such a diarrhea is to lie down; as the horizontal position is indispensable to a speedy cure. With the rice-water discharges or vomiting, commences the first stage of the disease, in which *Camphor* is the specific remedy, having actually the power of killing or destroying the animalcula or malignant agent which has infested the system. The application of the *Camphor* must be *effectual* and *immediate*, in the following manner:

Cover the patient, up to the chin, well and abundantly, particularly his feet, do not allow his arms out of bed;

then give him quietly, without showing anxiety by unnecessary hurry, of the tincture of *Camphor*,\* every two, three, or five minutes one drop, on a little piece of sugar; also, after fifteen or twenty minutes, some brandy and water, if he can retain any thing on the stomach. As soon as the patient begins to get warm, and to perspire, give the doses less frequently, and discontinue them altogether when he perspires freely. A few doses of brandy and water will then do him good and strengthen him. In this perspiration the patient must remain without change for eight or ten hours, when he will generally have a good appetite for a lunch of cold beef or mutton, with salt, dry bread, and cold water. If, however, the slightest indisposition yet remains, he must not leave the bed, as a relapse might take place. During the reaction, following the use of the camphor, frequently a headache ensues from congestion to the head, which in children and middle-aged persons disappears after the exhibition of one or two doses of *Belladonna* (for children three glob.; for adults six glob.); in aged persons, when stupor is combined with it, *Secale corn.* (thirtieth dilution) is yet necessary, three hours after *Belladonna* is given.

If costiveness follows the use of camphor, a cup of coffee without milk, and twenty-four hours afterward a dose of *Sulphur* (six glob.) is indicated. If the diarrhea is not entirely checked, but changed into a dysenteric one, with straining and bloody tinge, *Mercury* will suffice, followed by *Sulphur*, twenty-four or thirty-six hours afterward. If by the *Mercury*, however, the dysenteric stool is changed into the choleric, the use of *Camphor* is necessary again, followed by one or two doses of *Veratrum alb.* (twelfth dilution.)

If, however, the cholera proceeds into the *second stage*,

---

\* The tincture of *Camphor* ought to be made out of one part of gum to five parts of alcohol.

where cramps appear in different parts of the body, *Veratrum* and *Cuprum* are necessary; the former, when the cramps are more in the extremities and bowels; the latter, when in the breast, or when great oppression in breathing is present. It is frequently best to give these two remedies alternately, in the following manner: Dissolve of each remedy twelve or sixteen globules, in four tablespoonfuls of water, and give every fifteen minutes a teaspoonful (each remedy having its separate teaspoon), giving less frequently as the symptoms decrease in violence. In this stage, it is not so much the diminution of diarrhea, or vomiting, which indicates an amelioration, but the greater regularity and lesser rapidity of the pulse, which always will be followed by a discontinuance of the cramps, although these may yet threaten to break out again.

If these remedies are not successful within four or six hours, or if the cramps are more in the calves of the legs, where the slightest motion will excite them, or when stupor threatens, which is particularly the case in old persons, or when the diarrhea still continues, *Secale corn.* must be given, in the intervals of half an hour a teaspoonful, dissolved in water like *Veratrum* and *Cuprum*. If a few doses are given, wait an hour or two, if the case does not grow worse; if, however, there is a great deal of nausea or retching, increased by motion of head or body, *Tabacum* (six glob.) must be given, at any time during the attack.

If in this stage congestions to the lungs and heart appear, indicated by violent stitches in the side, almost preventing respiration, and producing restlessness and great anguish, give *Cuprum* and *Hydrocyanic acid* (*Prussic acid*), in alternation, every ten or fifteen minutes a dose (dissolved in water, a teaspoonful as a dose. N. B. *Prussic acid* must be given in the second dilution.)

The covering of the patient during the second stage

ought still to be warm, but already more according to the feeling of the patient; his drink, cold or warm water, left to his choice. Ice pills may be given to him freely, or injections of ice water if the bowels are cramped; rub slightly the cramped parts with the hands; warming bottles, and all the other known apparatus heretofore applied, are not of much use, although a homœopathist may allow them, not interfering with his medicine.

If the disease should go into the *third stage (collapse)*, the scene and treatment change entirely. No pulse, livid countenance, hoarse voice, and sunken eyes characterize this period. Yet our hope is not gone; particularly in cases which, up to this stage, were treated strictly homœopathically, or where the disease ran quickly through the first stages. This period generally lasts much longer than any of the former, and medicines must be given at longer intervals. The disease frequently stays in this stage for two or three days, almost unaltered—a slight improvement in the pulse is sometimes perceptible only within two days. In such cases, the medicine must not be changed, or repeated often. The principal remedy is *Carbo veg.*, in the thirtieth dilution, prepared as *Veratrum* above, of which for the first six hours, every hour a teaspoonful should be given. If a great deal of burning remains in the stomach, with drinking frequently, but little at a time, *Arsenic*, thirtieth dilution, prepared similarly, may be alternated with *Carbo veg.* But if thus four or six teaspoonfuls of each remedy have been given, their use should be discontinued for ten or twelve hours—particularly if the pulse has shown evidence of returning during that time. The improvement of the pulse is, in this stage the most important amelioration; the coldness of the surface is not of so much importance, as it often remains for a longer time without injury. If, after twelve hours, the patient's condition is not much improved, the same remedies



may be repeated, and in this manner continued for a couple of days, when, in most cases, a healthful reaction takes place.

The covering in this stage must be altogether left to the feelings of the patient, and only so much put over him as decency requires, as the patient generally refuses all covering, and is restless. To try to warm patients in this stage by external means, is not only useless, but cruel; as they actually complain of burning up internally, such means only hasten dissolution. We rather recommend to wrap the patient in a sheet, wrung in cold water, which may, after a few hours, be repeated, if he desires it. But, in such a case, the patient ought to be covered besides with a blanket or two, to follow up the beginning reaction. Cold drinks, particularly ice-water, are preferable in this stage.

In the period of convalescence, which is usually very short, great care ought of course to be taken in diet and exercise—rice and other gruels, afterward small quantities of broth, meat, and vegetables.

If a typhoid fever appears, it must be treated accordingly.

In that torpor or stupor which sometimes succeeds a severe attack of cholera, when the patient can not easily be aroused, is very weak, pulse slow, yet the expression of his countenance is natural, *Spiritus nitri dulcis* is of the greatest benefit, as *laurocerasus* is in those cases of stupor or lethargy, which border on paralysis of the brain, or on exhaustion of the nervous system, where the expression of the countenance is indicative of great suffering, the patient very weak, pulse slow, eyes half-closed, in continual lethargy, only broken by deep sighs and moaning.

We must mention here yet a disease, called *cholerina*, which often precedes and follows cholera. It is a diarrhea of fæcal matter, accompanied with rumbling of the bowels,



generally caused by the constitutionality of a cholera season, under debilitating influences of mind and body, such as *fear* and *grief*. If caused by the latter, *Phosphoric acid* and *China*, in alternation, after every evacuation a dose (six glob.) will suffice; if excited by fear, *Chamomile* (six glob.) a few doses; if, however, attended with bilious rheumatic symptoms, tending toward a typhoid state, such as headache, pains in the limbs, arms, and back, with occasional chilliness, *Bryonia* and *Rhus*, in alternation, two doses of each, every two hours one (six glob. a dose), and then waiting twenty-four hours, will change the symptoms, so that then either *Sulphur* will suit, or, if the diarrhea still continues, *China* and *Phosphoric acid*, in alternation as above stated.

*Sulphur* frequently takes away that disposition to diarrhea, so common in times of cholera. It ought to be taken only twice, on two succeeding evenings, or one evening and the next morning a dose (six glob.); after which no medicine of any kind should be taken for twenty-four or forty-eight hours.

We would recommend to every one, to have with him, in cholera times, a small bottle of *Camphor*, *Veratrum*, and *Cuprum*, in case of need; if not necessary, it certainly increases the feeling of safety.\*

---

\* In the cholera epidemic of 1849 which raged in Cincinnati worse, perhaps, than in any other city of the same size. every homœopathic family was provided with a small case of four vials. containing *Camphor*, *Veratrum*, *Cuprum*, and *Sulphur*, as the principal remedies for the prevention and cure of cholera. Printed directions accompanying these remedies enabled persons to make immediate use of them, when necessary; thus preventing entirely the application of allopathic medicines before the homœopathic physician could arrive. These simple but effective means kept the homœopathic practice free from many obstacles and difficulties, and made its results so glorious, and, even in the history of homœopathy, unrivalled. The average loss which was sustained by the homœopathic prac-

## PILES. HEMORRHOIDS.

DIAGNOSIS. — The piles form on the anus tumors or lumps, which protrude externally (*external piles*), or remain inside the rectum (*internal piles*.) These lumps either bleed (*open piles*) or do not bleed (*blind piles*.) The cause of the piles is a constitutional taint, which breaks out at certain times in this local affection of the rectum, and is then called the piles. It becomes an important disease, when bad, or a mere external treatment suppresses the outward piles, without curing the internal disposition. This latter, consequently, not unfrequently attacks more important organs, such as the lungs, brain, etc., where it generates incurable diseases. Just as detrimental to general health is the painful and abominable practice of the surgeons, of cutting off these lumps or tumors. It is a blessing, however, that in most cases such operations are of no avail.

CAUSES. — The use of strong and heating drinks, coffee, liquors, highly-seasoned food, sedentary habits, costiveness, suppression of long-continued discharges, etc., are among the exciting causes of this disease, and ought, consequently, to be strictly avoided in its treatment.

---

tice in 1849 amounted to only three and a half per cent., viz: of 2 410 patients treated homœopathically 85 were lost. In 1850 the attack of cholera in Cincinnati was not so severe—the cases generated not so much by epidemic influence as by indigestion, and other weakening causes. Such cases, however, were more difficult to manage, as the exciting cause could not be so easily removed, being of a material nature; yet the same homœopathic medicines proved entirely successful—even more so, in the aggregate, than the previous year. The loss of Drs. Pulte and Ehrmann was not more than three per cent.; from other practitioners, the ratio has not been as yet obtained.

The above description and treatment of Asiatic cholera is carefully made after our own experiences during these two epidemics, in which nearly two thousand cholera patients were attended by us.

Besides the medicines, use cold water externally, as a sitz-bath, or washing, and drink of it freely and often.

TREATMENT. — In all cases of piles, begin the treatment with

*Nux vomica*, as the principal remedy, which suits for all varieties of this complaint. After it, or, in chronic cases, in alternation with it, give

*Sulphur*. These two remedies are mostly sufficient to effect a cure; given alternately, every evening a dose (four glob.) until better. If these do not suffice, give of the following remedies, *Ignatia*, *Sepia*, *Belladonna*, *Hepar*, *Colocynth*, one after the other, in intervals of twenty-four or forty-eight hours.

The cure of chronic piles must be directed by a skillful homœopathic physician; as there are too many and too various constitutional considerations to be observed, which could not be given in a work like this.

DIET AND REGIMEN. — The same as stated in the article "Constipation."

#### ITCHING OF THE ANUS.

If it occurs in children, it is caused by the so-called pin-worms (see "Ascarides.") If caused by piles, *Nux vomica* and *Ignatia* are the principal remedies, given as stated under the heading "Piles." Besides these medicines, the washing or sponging in cold water, or even cold water injections are of the greatest benefit. Sometimes sweet oil, used in the same manner as the water, will be of use.

#### PROLAPSUS ANI. FALLING OF THE BODY.

By this term is meant the protrusion of a portion of the lower intestine or rectum. It occurs more frequently in children than adults, caused by severe strainings when at stool. The rectum can easily be brought back again by a

gentle pressure of the thumb and fore-finger, greased or oiled for that occasion, the patient bending forward during the operation, and reclining on his back after it. We have remedies which diminish the tendency to this troublesome disease.

TREATMENT. — *Ignatia* is the principal remedy, and must be given once in twenty-four hours (four glob.) for six or eight days ; then discontinue eight days, and if not better, give

*Mercury*, in the same manner. After that, *Nux vomica* and *Sulphur*, in the same manner, until better. *Sepia*, *Lycopodium*, *Colocynth*, are also suitable to be given, if necessary, in the same manner.

Externally, wash with cold water, as in “Piles.”

DIET AND REGIMEN. — As in “Diarrhea.” If this treatment is of no avail, consult a homœopathic physician.

#### RUPTURE.    HERNIA.

A protrusion or swelling in the inguinal region or groin, generally indicates a rupture, by which the intestines descend into the protruded part. As a disease of such consequence and importance is beyond the sphere of an exhausting discussion in a domestic work, we must limit our remarks to a few dangerous points, to which the disease may run at times, leaving the treatment of the rupture itself to the combined efforts of the surgeon and homœopathic physician, who must be consulted.

Sometimes an *incarceration* of the rupture takes place, when the tumor becomes very painful, inflammation in the surrounding parts ensues, with vomiting, quick and hard pulse, and fever. If the rupture can not be brought back, or the inflammation reduced by other means, mortification and consequent death are unavoidable.

First, try the reduction in the following manner: place

the patient on his back on the bed, grasp the swelling with one hand gently, and rub and press it with the fingers of the other hand, pressing upward toward the body constantly, sometimes even in a rotary manner. Continue these efforts for half an hour; if the swelling is repressed, keep it in its place with the palm of the hand for some time, giving, besides, the following medicines:

*Aconite*. If the fever is *very high*.

*Nux vomica*, followed in two hours by

*Opium*. If the *strangulation* of the rupture continues, and afterward in alternation with *Nux vomica* until better.

Also, *Lachesis*, *Belladonna*, *Rhus*, *Arsenic*, *Veratrum*, can be given in extreme cases, if necessary; one after the other, every two hours one remedy, until better.

ADMINISTRATION. — Dissolve of the necessary remedy twelve globules in half a teacupful of water; give every fifteen or thirty minutes a teaspoonful, for two hours, or until better.

DIET AND REGIMEN. — As in fevers. Externally, the application of cold water or ice is recommended. This will do when the parts are not inflamed too much; in which latter case warm water fomentations are preferable. Complete rest is indispensable, and a truss must not be worn until all soreness has subsided.

#### DROPSY OF THE ABDOMEN. (*Ascites*.)

As this is one of those diseases which require a most skillful medical attendance, in order to secure a successful termination, we can not pretend to give more than what is necessary to a knowledge of its rise and progress. This will enable persons to be aware of danger in time to seek for help, when it is yet possible to render it successfully.

DIAGNOSIS. — A swelling and tension of the abdomen, which follows the position of the patient; a fluctuation in-

side the abdomen is distinctly felt by the palm of one hand while clapping gently against one side, when the other hand presses the opposite side of the abdomen, in the lowest part of which the fluctuation is felt the strongest, while the patient is standing erect. Urine is scanty, brown, like beer (peculiar to ascites); stools are scanty and dry; also, the skin, tongue, and mouth dry; when the water is collected in a sack (*hydrops saccatus*), the swelling is unequal in the beginning; the urine less scarce and brown. As the disease progresses, the feet and other parts swell successively; fever and dry cough appear, which consume the strength more and more, until death ends the misery.

CAUSES. — As such, we find previous acute fevers, scarlet or other eruptions, diseases of important abdominal organs, especially the liver, badly treated, intermittents of too long standing, which dililitate the absorbing vessels, gout, abuse of spiritous liquors, mechanical lesions, such as blows on the abdomen, falls, etc.

This disease sometimes lasts very long, according to the causes, and the constitutionality of the patient.

A pregnancy, not closely watched in its progress, may be mistaken for a dropsy of the abdomen; but, in such a case, a skillful physician must decide.

TREATMENT. — As this disease can be cured, if early attended to in the right manner, we recommend every one to apply immediately to a good physician. Until one can be procured, give,

*Arsenic* and *Sulphur*, in alternation, every four days one dose (six glob.)

The tapping may be allowed in cases where the accumulation of water is very rapid; but no one must trust in it as a curative means; the necessary internal treatment must therefore be continued, until cured entirely.



DIET AND REGIMEN.—As in all chronic diseases, the usual homœopathic diet.

#### EXTERNAL INJURIES OF THE ABDOMEN.

On receiving large wounds in the abdomen, do not despair immediately, as a great many are cured who received the severest cuts and shots in the abdomen, if only properly treated. Clean the wound with lukewarm water, and replace the entrails as soon as possible, handling them not with the hands, but with clean linen cloths. If the patient is faint, from loss of blood, give *China*; if frightened, *Coffea*, and afterward *Arnica*. Sew up the wound carefully with a silken thread waxed, leaving an opening on the lowest corner, but cover it with scraped linen, dipped in a mixture of *Arnica* (twelve drops to a teacupful of water.) If diarrhea ensues, give *Colocynth* alternately with *China*.

If a child has swallowed a button, or any other metal substance, do not give it aperient medicine, in the hope of purging it out of the system more quickly; this is impossible; on the contrary, the substance will come out less quickly, as the liquid state of the fæces prevents them from carrying it along. In the meantime, observe carefully each stool, to be sure the substance has passed. If a needle is swallowed, it may work itself through a different part of the body; in such a case, give *Silicea*, every six days a dose (four glob.)

If live animals are swallowed, such as frogs, insects, etc., let the patient drink plenty of sweet oil.

## CHAPTER XV.

### AFFECTIONS OF THE URINARY AND GENITAL ORGANS.

WE invite the special attention of our readers to the study of the anatomical position and structure, as well as the physiological bearing, of the organs named in the above heading, as it will be important for their true diagnosis. These diseases are various and severe.

#### INFLAMMATION OF THE KIDNEYS. (*Nephritis.*)

DIAGNOSIS. — In the region of the kidneys (on both sides of the spine, above the hip, and below the short-ribs), appears a pungent, pressing pain, shooting downward to the bladder; difficulty of urination; stranguria, or ischuria (the latter only when both kidneys are inflamed, which rarely ever occurs); the urine red and hot; the testicle drawn near to the abdomen on the affected side, painful and swollen; the foot on the same side sometimes spasmodically affected and benumbed; in severer cases, vomiting; colic pains; the pains in the kidney are worse when lying on the affected part, and the back when standing and walking.

If this inflammation is not cured entirely, the kidney hardens and indurates, or suppuration takes place, which leads to the formation of abscesses and their opening, either outside through the skin, or inside in the intestinal

canal. This chronic affection of the kidneys requires medical skill for its successful termination.

The principal causes are : stones or gravel in the kidney ; violent blows and concussions, or any other external injury bearing on this part ; excess in the use of wines and liquors ; lying on the back for a long time ; lifting heavy loads ; suppression of piles, and their congestion transferred to the kidney ; the use of acrid diuretics and the abuse of Spanish fly, sometimes even when only applied as a blister, externally.

TREATMENT. — *Camphor*, when the disease is caused by the application or use of Spanish fly (*cantharis*), or when no urine passes, or only in a slender stream very slowly, with burning in the urethra and bladder.

*Aconite*. If the fever is high, give several doses (four glob. each) of this remedy, every hour one, before the following remedies are resorted to :

*Cantharis* is the principal medicine, when there are shooting, tearing, and incisive pains, with *painful emission of a few drops of urine only*, or *complete ischuria* ; sometimes urine mixed with blood.

*Belladonna*. In most cases, where the pains are worse periodically, shooting from the kidney down into the bladder, with great anguish and colic ; in fleshy women, at the time of change of life (critical age), when the menses stop. (*Hepar sulph.* will suit after *Belladonna*.)

*Nux vomica*. When this disease is caused by suppressed piles, or when there is any other abdominal congestion, with tension, distention, and pressure in the kidney.

*Pulsatilla*. In delicate females, whose menses have stopped, or are very scanty ; also, when the urine appears turbid, leaving a purulent sediment.

*Arnica*, externally and internally, if caused by external

injuries ; and after it the above remedies may be resorted to.

N. B. If nephritis is caused by the pressure of stones in the kidneys, the fever which usually accompanies this disease does not appear so quickly, but, instead of it, the foot and thigh of the affected side becomes numb, and the testicle retracts ; sure signs of stones in the kidneys or the ureter. In such a case apply on the most painful spot, where the stone is lodged, narcotic, warm fomentations of hops mixed with tobacco leaves ; as soon as the stone or gravel reaches, in its descent, the bladder, the pains and inflammation caused by them cease at once. Beside this external treatment, give the patient the above medicines as indicated.

If the kidney indurates, give *Mercury*, every evening a dose (four glob.)

If suppuration takes place, give *Lachesis* and *Hepar* alternately, every evening a dose (four glob), for eight days, or until better ; apply to a homœopathic physician.

ADMINISTRATION. — Dissolve, of the necessary medicine, twelve globules in half a teacupful of water, and give every half hour, hour or two hours, a teaspoonful until better, or until another remedy is needed.

DIET AND REGIMEN as in fevers generally ; but particularly wines, liquors, and beer are prohibited in convalescence.

#### INFLAMMATION OF THE BLADDER. (*Cystitis*.)

DIAGNOSIS. — Burning pains in the region of the bladder, with external swelling ; tension ; heat and pain on touch ; urine red and hot ; difficulty or inability of making water ; constipation, with tenesmus ; fever, with a hard pulse ; in some cases, vomiting and hiccough.

The issue of this inflammation is either resolution, indi-

cated by a thick urine, or suppuration discharged with the urine, or abscesses and fistulas from the bladder outside, or induration, which leads to a thickening of the bladder, impeding, thereby, its action, and leading to various chronic disorders ; or, if fatal, gangrene takes place.

Its causes can be external injuries ; suppressed piles and menstruation ; rheumatic, gouty, or syphilitic transfers ; stones in the bladder ; pressure on the bladder in pregnancy, or during the birth of the child.

TREATMENT is the same as in “ Inflammation of the Kidneys ;” see this article.

DIET AND REGIMEN, also, the same.

#### STRANGURY, DYSURY, ISCHURY, AND ANURY.

These four terms signify only different degrees of the same affection.

In *strangury*, the discharge of urine is *painful* and *difficult*.

In *dysury*, *difficult* and *incomplete*.

In *ischury* and *anury*, it is totally *suppressed*.

Strangury and dysury are troublesome, but not dangerous complaints, while the true ischury, or total retention of urine, ranks among the most dangerous diseases, and kills, either by gangrene or rupture of the bladder, effusing the urine into the abdomen ; in its chronic form, however, the urine can be absorbed and carried into the circulation, producing, thereby, eruptions on the skin, and making the perspiration, saliva, etc., smell urinous.

The causes of these affections are various. Mechanical injuries or impediments, such as gravel, stones, callosities and strictures, spasmodical, or caused by syphilitic disorders ; tumors in these parts ; falling of the womb, or its dislocation ; congestion of every kind ; paralysis of the bladder ; the use of ardent spirits, Spanish fly, violent diuretics, etc.

TREATMENT.—For *strangury* and *dysury*, if caused by the abuse of *ardent spirits*, *Nux vom.*, *Pulsatilla*, and *Sulphur*; if by the use of *Spanish fly*, *Camphor*, *Pulsatilla*, and *Aconite*; if by *suppressed piles*, *Nux vom.*, *Pulsatilla*, *Sulphur*; if by *taking cold*, wet feet, etc., *Dulcamara*, *Nux vom.*, *Mercury*, *Pulsatilla*, *Belladonna*; if after a *fright*, *Aconite*; if after a *fall, blow*, etc., *Arnica*; if in *pregnant females*, *Cocculus*, *Pulsatilla*, *Nux vom.*; if in *children*, *Pulsatilla*, *Aconite*, *Belladonna*, *Mercury*.

For *ischury* and *anury*, if it is *spasmodic*, *Nux vom.*, *Pulsatilla*, *Opium*, *Hyoscyamus*; *inflammatory*, *Aconite*, *Cantharides*, *Belladonna*, *Nux vom.*, *Pulsatilla* (see “*Cystitis*”), *paralytic*, *Hyoscyamus*, *Dulcamara*, *Arsenic*, *Lachesis*.

ADMINISTRATION the same as in “*Inflammation of the Kidneys (Nephritis.)*”

DIET AND REGIMEN the same as in *nephritis*; besides this, in *strangury* the drinking of large quantities of cold water, or the moderate use of gum-arabic, dissolved in water, is strongly recommended. In cases of disease of the bladder, consult, as soon as possible, a homœopathic physician.

INCONTINENCE OF URINE. (*Enuresis, incontinentia urinæ.*)

This troublesome disease exists in three varieties:

1. It is a *complete* or *paralytic* incontinence of urine (*enuresis completa s. paralytica*), when the patient is constantly troubled with an involuntary dribbling of urine, owing to a variety of causes, all having the tendency to paralyze, more or less, the contractive power of the muscles around the neck of the bladder. Instances of this kind are: *apoplexy*; *consumption*, and *paralysis of the spine*; too great distention of the bladder by long retention of the urine; surgical operations on the bladder; *paralysis*, or weakness of the bladder, particularly in old age; difficult and hard parturitions. This form of incontinence of urine requires



the most careful attention of a homœopathic physician. We only mention some of the principal remedies for this form of *enuresis*, with which a treatment may be commenced: *Hyoscyamus*, *Belladonna*, *Causticum*, *Conium*, *Arsenic*, *Lachesis*.

2. Or it is an *incomplete* or *spasmodic* incontinence of urine (*enuresis incompleta s. spastica*), where the urgency to pass urine is so sudden and great that the patient is immediately forced to yield. This form occurs very frequently, caused by continual irritation in the bladder, or some contiguous part, either by stones, gravel, worms (*ascarides*), (see "Worms"); menstrual or hemorrhoidal congestion; gastric derangement, with rheumatic or gouty complication; scirrhus tumors in the bladder, rectum, prostrate glands; ulcers, fistulas; also, from mechanical pressure of the pregnant or dislocated uterus (see "Diseases of Females;") or, lastly, by the bad habit of urinating too frequently, diminishing, thereby, the size of the bladder.

The principal remedies for this complaint are: *Bellad.*, *Causticum*, *Cina*, *Hyoscyamus*, *Ignatia*, *Pulsatilla*, *Rhus*, *Lachesis*, *Mercury*. (Compare each remedy in "Materia Medica.")

ADMINISTRATION of these medicines as in "Strangury."

3. The urine escapes *involuntary* only *in the night* (*wetting the bed*), (*enuresis nocturna*.) This occurs mostly in children, and depends on bad habits, uncorrected in early years, or on local irritations, such as worms (see this article), or other derangements, enumerated under the form No. 2; (see their treatment.) If it is a bad habit, try to correct it, by diminishing the quantity of drink allowed in the evening, by laying the child on its side in the sleep, by awakening it several times during the night in order to pass urine, or even by chastisement in the morning, the effects of which will be remembered, even in the sleep.

Besides, use the following remedies: *Silicea*, every third evening a dose (three glob.), for two weeks; if not better,

*Sepia* in the same manner; if not better,

*Sulphur*, *Arsenic*, *Carbo veg.* in the same manner, one after the other.

### DIABETES.

DIAGNOSIS.—Any secretion of urine, the excessive increase of which has a weakening, morbid effect on the whole system, may be called a diabetes; although the quality of the urine be not altered, its appearance shows that it contains more water, it being colorless (*nervous urine*); cases of this kind either correct themselves, or are in connection with hysteric or hypochondriacal diseases, the cure of which will, also, take away this form of diabetes. But where the urine is not, or not much, increased in quantity, however changed in quality, having a *sweetish* taste, and containing *saccharine* (sugar) matter (*diabetes melitus*), as high as an ounce in a pound, it is necessary to institute an early and earnest treatment for this disease, which, in its commencement, can be cured easily, and only becomes fatal by neglect. But, as such a complaint is too important in its progress and issue to be treated domestically, we here only indicate the remedies, with which a rational treatment may be commenced.

*Remedies.*—*Phosphoric acid*, *Carbo veg.*, *Conium*, *Mercury*, *Sulphur*. The first of these medicines is almost a specific, as also the two last ones, which may be given in alternation; give, every evening, a dose (four glob.), for one week, and then wait a week for its effect.

DIET.—Only bread, roasted meats, and soups of meat, are allowed; no vegetables whatever; the patient must keep in an even temperature, dry and warm; must wear flannels on the skin, if possible.

PILES OF THE BLADDER. (*Hæmorrhoides Vesicæ.*)

As the pile-disease consists in a congestion, which can tend to any part of the system, and produce piles, where the nature of the attacked organ admits of it, it is not strange to speak of piles of the bladder, because this organ allows the same disorganization in this respect, as the rectum. They are also either *blind*, or *open*, *running* piles.

DIAGNOSIS.—If they are *blind*, or not *running*, they create great difficulty in making water, strangury, ischury, sometimes spasms, and inflammations of the bladder (see these articles), with consequent induration and suppuration.

If they are *open*, *running* piles, the blood coagulates in the bladder, obstructs the passage of urine, and may contribute to the formation of stones.

TREATMENT.—Consult a homœopathic physician, if possible. If not, use the following remedies: *Nux vom.*, *Pulsatilla*, *Sulphur*, besides all those recommended in the articles on *strangury* and *ischury*, in the same manner as there stated.

## POLYPUS OF THE BLADDER.

The neck of the bladder is the place most favorable to the generation of this kind of fleshy excrescences, where they occasion frequent obstructions in urinating, producing similar symptoms to those exhibited in piles and stones of the bladder. Their existence, however, is rendered beyond doubt, as soon as fleshy particles of a round form, with a stringy substance attached to them, pass from the bladder; females are subject to them more than males.

TREATMENT.—If possible, consult a homœopathic physician for this disease, as the cure depends too much on constitutionality to make a prescription of remedies for a domestic work of this kind possible. To commence the

treatment, however, give *Calcarea carb.* every third or fourth evening a dose (four glob.), for four or six weeks.

N. B. I once cured polypus of the bladder with *Chamomile*, in a complicated case, where the concomitant symptoms strongly advised its use. No other remedies were used in this case; after the exhibition of *Chamomile*, the polypus passed off altogether.

*Staphysag.* is recommended after *Calcarea carb.*, in the same manner.

#### GRAVEL AND STONE IN THE BLADDER. (*Calculus.*)

DIAGNOSIS.—It is sometimes very difficult to be certain of the existence of a stone in the bladder, as it produces all the symptoms recorded under *strangury*, *ischury*, and *piles of the bladder*. (See these diseases.)

The only sure guide in establishing a true diagnosis, is the examination by a competent physician or surgeon. In gravel, the diagnosis is easier, as some of it will pass, from time to time, clearly showing the nature of the disease.

TREATMENT.—The treatment for a *stone in the bladder* can not be given here, as such a disease must not be left to domestic practice.

For *gravel*, the best remedy is *Sarsaparilla*, either in a homœopathic preparation (four glob.), or, if not at hand, of the tincture of *Sarsaparilla*, every evening one drop on sugar, until better. Besides this remedy, those stated under “*Strangury*” are recommended.

#### HEMORRHAGE WITH THE URINE.

(*Hæmaturia. Mictus cruentus.*)

DIAGNOSIS.—The blood comes with the urine, either mixed with it, like dark beer (then it proceeds from the kidneys, *hæmaturia renalis*), or it is separated from the urine,

, coagulated (then it comes from the bladder, *hæmaturia vesicalis*), or the blood comes alone, without urinating (then it proceeds from the urethra, *stymatosis*.) In the first case, pains are felt in the region of the kidneys; in the second, the region of the bladder is painful; and in the third case, the pains are in the urethra.

CAUSES.—This disease occurs always in consequence of another already there located, to which we refer the reader. These are: inflammation of the kidneys and bladder, and their causes, mechanical and external, stones and gravel, strangury, etc.

TREATMENT.—The principal remedies are: *China*, *Ipecac.*, *Arnica*, *Pulsatilla*, *Lycopodium*, to be given as stated under “Strangury,”

#### AFFECTIONS OF THE PENIS.

If the front of the penis—the gland and the prepuce—is inflamed, red and swollen, give, when caused by friction, *Aconite* and *Arnica*, in alternation, every two or three hours a dose (four glob., infants half a dose), until better; keep quiet on the bed or a couch; abstain from all meat diet, and bathe, if necessary, with cold water alone, or water and milk; if caused by want of cleanliness, give *Aconite* (four glob.) alone, and bathe. If it was caused by the touch of poisonous plants, give, in the same manner, *Belladonna* and *Rhus*, after *Aconite* has been given. If a burning discharge appears from the urethra, give *Mercury*, and if not better in twenty-four or thirty-six hours, give *Capsicum*, and in a few days after, *Hepar*; if the parts become hardened, callous, and bluish, give *Lachesis* and *Arsenic*, in alternation.

DIET AND REGIMEN.—Nothing but milk and water, dry toast, and gruels.

N. B. For syphilitic disorders, chancres, and gonorrhea, see these articles.

#### AFFECTIONS OF THE TESTICLES.

*Inflammation*, with painful swelling of the testicles, from *External injuries* : *Aconite* and *Arnica*, as above under the "Affections of the Penis;" *Arnica*, also, externally.

*Gonorrhea suppressed* : *Pulsatilla*, externally and internally, as above.

*Metastasis of the mumps* : see this disease ; *Pulsatilla* is the principal remedy.

*Abuse of Calomel* : *Pulsatilla* and *Carbo veg.*, in alternation, as above.

If necessary, give *Sulphur* after the above remedies have ameliorated, but not altogether cured.

DIET AND REGIMEN as in "Affections of the Penis."

*Hydrocele*, or dropsy of the scrotum : if in infants shortly after the birth, washing with a mixture of six drops of *Arnica* tincture, in a teacupful of water, three or four times a day ; if in older children, the same, externally, and *Pulsatilla* every evening a dose (three glob.) internally ; if in adults, *Pulsatilla*, *Silicea*, *Sulphur*, each remedy for eight or twelve days, every fourth day a dose (four glob.) during that period, until better, and if not, apply to a homœopathic physician.

Diet and regimen as usual in chronic diseases, when treated homœopathically.

#### SEMINAL WEAKNESS.

(*Pollution. Impotence. Onania.*)

These diseases are very weakening, and frequently fatal to happiness, health, and even life. As their introduction



here would carry us beyond the sphere of a domestic treatise, we advise our readers not to neglect, for a single day, the above complaints, which explain their nature in their names, but to have the advice of a skilful homœopathic physician at once.

#### ERUPTIONS ON THE GENITAL ORGANS.

Eruptions on the glans: *Bryonia*, *Rhus*; on the hairy part, *Lachesis*; on the penis, *Graphites*, *Phosphoric acid*; on the prepuce, *Sepia*, *Silicea*; between the thighs, *Petroleum*.

*Prurigo on the scrotum* (tetter) requires, mostly, *Sulphur Dulcamara*, *Nitric acid*, *Petroleum*, *Thuja*.

ADMINISTRATION.—Every evening a dose (four glob.), and thus every medicine used for four days, before another one is commenced; if it is not better, wash with a solution (twelve glob. in half a teacupful of water) of the same remedy during its internal use.

DIET AND REGIMEN as in chronic diseases.

*Erysipelas of the scrotum* requires *Belladonna*, *Rhus*, and *Arsenic*, in the same manner as stated under “Erysipelas.” (See “Diseases of the Skin.”)

Diet and regimen as in “Erysipelas.”

#### SYPHILIS.—GONORRHEA.

We can only mention here the commencement of the treatment for these complaints, as inveterate or chronic disorders of this kind are so complicated and difficult of treatment, as to require the utmost attention of a skillful homœopathic physician.

##### a. *Syphilis. Chancre.*

If infection has taken place, and a slight soreness manifests itself on the glands, with a pimple which soon becomes a

small ulcer, marked by its hard surface and elevated edges, take, internally, *Mercury*, three times a day (six glob.), and wash, externally, with a solution of *Mercury* (twelve glob. in half a teacupful of water), until the advice of a physician can be obtained.

*b. Gonorrhea. Clap. Gleet.*

If, after infection, a soreness of the whole urethra ensues, with inflammation of the glands in particular, take *Aconite*, every two or three hours a dose (four glob.), for six or eight times, and after it, if a running appears, *Mercury*, every three hours a dose (four glob.), until medical aid can be procured. Outwardly, use cold water bathing; to subdue pain and inflammation, keep *perfectly quiet*, and abstain from all meat diet.

DIET AND REGIMEN are alike for both complaints: no meat whatever; but milk, water, and bread; the most perfect rest.

## CHAPTER XVI.

### DISEASES OF FEMALES.

THERE are diseases peculiar to the female sex, whose distinctive character is founded in the different physical organization of the female from the male. It shall be our endeavor to be as minute in treating of them as the object of this work will allow, although a great many of them must, and will, always demand the especial attendance of a physician, on account of their importance to life and difficulty of treatment; yet a right knowledge of their nature will diminish fear, and create confidence and trust—a great lever in surmounting medical difficulties.

#### 1. MENSTRUATION.

With the appearance of the menses (*courses*) the age of puberty commences for the female, and various changes take place in the mental and physical development; such as change of voice, expansion of the chest, enlargement of the breast, a marked reserve in manners, and an increased attention for outward decorum and appearance. This time generally comes, in northern climates, between the thirteenth and fifteenth year; in southern, between the eleventh and thirteenth. The same difference of time is observed in the disappearance of the menstrual discharge, which takes place, in the northern climates, generally between the forty-third and forty-fifth year; in the southern, between

the forty-first and forty-third year. The menses appear quite regular in a healthy female, every twenty-eight days (a slight deviation of from two to three days earlier or later does not constitute of itself disease), and last, at an average, five days, although this is no absolute rule.

We can not recommend too strongly a strict adherence to the general principles of Hygiene, as the departure from these lays, in most cases, the foundation to those derangements in the menstrual functions which prove so fatal to the life and happiness of a female. And here we may remark that, in the course of a long practice, we have found the wearing of thin-soled shoes to be the most fruitful source of the decay of female beauty, and the decline of female health. The damage of tight lacing (although considerable) is nothing in comparison with the fatal habit of appearing in all kinds of weather in thin-soled shoes; the consequences of the latter are beyond description, fearful, and destructive. Most all diseases which follow a stoppage of the menses, consumption, fluor albus, etc., can be traced to this source. Let reform set in!

#### *Obstructions of the First Menses.*

When the menses do not appear, although the time of puberty has come, the young girl will show signs of deranged health which demand our immediate attention. She has, perhaps, frequent bleeding at the nose, congestions to the head, flushed face, constriction of the chest, palpitation of the heart, etc. If these symptoms manifest themselves, give first,

*Pulsatilla.* If, besides other symptoms, she looks rather pale, is slender and feeble, melancholy and sad; feels better in the open air.

*Bryonia.* If she looks flushed in the face, her nose bleeds frequently, inclined to constipation.

*Veratrum*. If she is chilly and inclined to diarrhea.

*Sulphur*. If either of the above remedies have not brought on the menses.

ADMINISTRATION.—Of the selected medicine, give every evening a dose (four glob.) for four nights, then discontinue four days, and give *Sulphur* in the same manner. If not better in four or six weeks consult a physician.

DIET AND REGIMEN.—Let the diet be simple, but nutritious ; exercise as much as possible—apply warm foot-baths before going to bed.

### *Chlorosis.*

DIAGNOSIS.—Pale, white color of the skin, cheeks, and lips ; chilliness, lassitude, weak, slow pulse, yet easily excited on the least exertion ; difficulty of breathing, and palpitation of the heart when exercising ; swelling of the feet ; no appetite, but sometimes an immoderate desire for unusual things, earthy substances, as chalk, etc. The blood is thin, watery ; if not cured, general dropsy and consumption, or nervous diseases can set in. It is mostly a disease appearing in young girls, when the first menses do not appear, or after several hemorrhages, or other diseases, carrying with them great loss of life-sustaining fluid.

TREATMENT.—If it occurs in young girls, the treatment mentioned for obstruction of the first menses in the preceding article will suit ; yet, after *Sulphur*, we would recommend *Calcareæ carb.* ; if it occurs, however, after severe sickness, or hemorrhages, give *China* and *Carbo. veg.*—every other evening a dose (six glob.) alternately, for at least five or six weeks. In both cases, if the above medicines do not relieve, apply to a skillful physician without delay.

DIET AND REGIMEN. Let the diet be very nutritious, exercise plenty, particularly in open air—if mineral springs

are chosen for a summer resort, give the preference to the chalybeate, containing iron.

*Suppression of the Menses.*

Or their *temporary cessation*, when once well established. This occurs either suddenly by taking cold (from wet feet mostly), overheating, violent mental emotions, faults of diet, etc., and often produces violent congestions to the chest, head, or stomach, with cramps, convulsions, inflammations, etc.; there is no disease which could not appear in consequence of it; or the menses have gradually disappeared, without creating any immediate symptoms in the female economy, giving rise to suspicion of pregnancy.

TREATMENT.—Besides the remedies recommended in “Obstructions of the First Menses,” which will be of the greatest benefit here, and of which *Pulsatilla* is the principal, we recommend

*Aconite*. In alternation with *Bryonia*, every hour or two hours a dose (four glob.), in a sudden check of the menses, producing congestions to the head and chest, or *Cocculus*, in alternation with *Veratrum*, in the same manner, if severe cramps in the abdomen are present.

*Opium*. If all the blood seems to have rushed to the head, producing *heaviness there*, with a dark redness of the face, and drowsiness, even convulsive jerks of the muscles.

*Capsicum*. In spasms of the chest, and in alternation with *Veratrum*, if, also, the *abdomen* suffers; or with *Opium*, if the *head* is congested.

If the menses have been checked by mental agitation or depression, take those remedies recommended for fright, fear, etc., in “Affections of the Mind.” See this article.

ADMINISTRATION the same as in *Aconite* above.



*Painful Menstruation.*

The causes of these tormenting distresses of females are generally laid in the early part of womanhood, and owing chiefly to improper treatment of other diseases, suppression of eruptions, or habitual discharges, rheumatic disorders, colds, etc. The pains either appear before or during the flow of the menses, sometimes resembling real labor pains, with bearing down and forcing; at other times as a constant aching in the loins, hips, and limbs. They generally diminish in violence as soon as the regular flow has commenced, but not always.

TREATMENT.—If possible, lay down, cover well, and take *Coffea*, *Pulsatilla*, and *Veratrum*, in alternation, every half hour a dose (four glob.) *until better*. If this does not suffice, take *Nux vom.*, if the *forcing* pains predominate; *Cocculus*, if *colic pains* in the abdomen appear, with shortness of breath; *Chamomile*, if with discharge of dark, colored blood there are pains like labor-pains, together with colic pains and tenderness of the abdomen.

If a profuse perspiration sets in while in bed, do not leave it soon after the pains cease, nor cool off too quickly, else the pains return. Avoid the use of heating substances, either externally or internally, save a warm brick on the feet or stomach.

ADMINISTRATION.—As above, under *Coffea*, etc.

DIET.—No coffee for the first two days.

*Menstruation too Early.*

If the menses appear too early, say every two or three weeks, the disorders causing it are too complicated frequently to be prescribed for in a domestic treatise; apply to a physician; yet the cure may be commenced with the following remedies:

*Ipecac.* Almost a specific in all passive hemorrhages, where feebleness, dullness, nausea, and coagulated discharges prevail.

*Ignatia.* Where the menses return every two weeks, spirits are depressed ; in alternation with *Ipecac.* if its symptoms are also present.

*Belladonna.* Heat and pains in the head, with cold feet and dryness of the throat, bearing down in the abdomen.

*Calcarea carb.* is almost a specific in this disease, where with the menses a diarrhea or frequent discharges from the bowels appear, with pains in the bowels ; suitable after *Belladonna.*

*Sulphur.* If the menstruation is *too early* and *too profuse.*

*Natrum muriaticum.* If *Calcarea carb.* is insufficient.

ADMINISTRATION.—Just before, or in the beginning of the menses, one or two doses of the selected medicines every three hours one dose (six glob.) ; after the menses are over, take one more dose, and then wait till the next appearance of the menses, and observe whether they are more regular as regards time.

#### *Menstruation too Late and too Scanty.*

*Pulsatilla* is the principal remedy, when the above difficulty exists, and also when the menses appear irregular, sometimes too late and too profuse (as this is the case particularly at the critical period—change of life), sometimes too early and too scanty. In the former, *Lachesis* alternates well with *Pulsatilla.* In most all cases,

*Sulphur* is necessary, to complete the cure.

Compare, also, “Suppression of the Menses,” “Obstruction of the First Menses.”

ADMINISTRATION, DIET, AND REGIMEN, the same as stated there.

*Menstruation too Copious. Flooding.*

In cases of this kind, causes mental and physical may operate, to enumerate all of which here, would be impossible. We content ourselves in directing the reader to the various sources, where he can find their remedies.

If *mental* causes, such as fright, fear, etc., exist, give the remedies stated under “Affections of the Mind,” for the several exciting causes.

If *external injuries*, see this chapter. In cases of this kind, *Arnica* is the first remedy, after which others may be selected.

In general, however, *Ipecac.* is the principal remedy (see “Menstruation too Early”), followed by *China*, if there is great weakness, buzzing in the ears, faintness when raising the head off the pillow; *Belladonna*, if there is downward pressure; *Pulsatilla* and *Lachesis*, if it occurs during *change of life*; also, in such a case, or in aged women, *Ipecac* and *Secale*.

*Chamomile*. If dark, clotted blood is discharged.

*Coffea* and *Camphor*, in alternation, when there is, besides the above symptoms, *exceedingly painful colic*.

*Platina*. After *Belladonna* or *Chamomile*, when the discharge is too profuse, or of too long duration — of *black* and *thick blood*, with great nervousness, sleeplessness, and constipation.

*Secale*. Particularly with *great weakness* and *coldness of extremities*.

If real *flooding* ensues, resisting the above medicines, the application of *cold water*, or *pounded ice*, over the lower part of the abdomen, externally, is necessary to coagulate the blood in the vagina and uterus, which stops the hemorrhage. There can be no fear of getting cold in doing this, if it is done well; always keep the patient lightly but well

covered. In such a case, the patient must lay with her head low. This treatment will be effectual in the severest cases of flooding.

ADMINISTRATION. — Dissolve six glob. of a remedy in six teaspoonfuls of water, and give every fifteen, twenty, or forty minutes a teaspoonful, sometimes in alternation with another, and lengthen the intervals as the patient gets better, when no medicine is needed any more.

DIET AND REGIMEN. — Every drink must be cold and not stimulating, except when faintness appears, with deadly paleness of the face, no pulse, and cloudiness of sight, give wine and brandy in frequent, small quantities. *Camphor* and *China*, in such cases, are of the greatest service. The patient ought to lay perfectly quiet.

*Menstruation of too long Duration.*

Compare “Menstruation too Copious,” and give of the selected medicine every evening and morning a dose (six glob.) until better. (See, also, next article, on “Change of Life, or Critical Period.”)

DIET in such cases must be highly nutritious. It is best to consult a physician early.

*Critical Period. Cessation of the Menses. (Change of Life.)*

This period, commencing about at the age of forty-five years, forms one of the most important in the life of a female. If not guided through this critical time by the counsel and aid of a skillful physician, she gathers the seeds of endless miseries, or even early death. This period may be a blessing to her, as well as a source of great distress; as after it her health either becomes more confirmed, or disorganizations in internal organs are formed, which soon carry her off. Without enlarging further on the subject, we recommend the early and constant advice

of a skillful physician during this time, which generally lasts from one and a half to two years.

First, an irregularity of the courses is experienced ; they stop for three months, then reappear with great violence, then stop for four or six months, during which time the woman shows more or less symptoms of congestions ; piles appear ; the limbs swell ; pruritus (violent itching of the private parts) troubles them ; cramps and colics in the abdomen ; asthmas and palpitations of the heart ; sick-headache ; hysterics ; apoplexy, etc. Sometimes they remain perfectly healthy. These maladies are so various, and continually changing and complicated, that they require the constant, watchful care of a family physician. Do not neglect them.

TREATMENT.—We can here only give general rules : eat and drink moderately ; sleep in airy, well-ventilated rooms ; avoid violent emotions or exercise, but contrive to be busy mentally and bodily ; shun exposure to inclement weather, wet feet, etc. (See “Hygiene.”)

*Pulsatilla* and *Lachesis* are the principal remedial agents in this period, of which take every six days one dose (six glob.) alternately, unless other remedies are necessary. If diseases otherwise occur, see their respective chapters.

### PROLAPSUS UTERI. (*Falling of the Womb.*)

A great deal of abuse has taken place of late with this disease, which has almost become a fashionable one, as neuralgia, etc., through the neglect and ignorance of practitioners, who call every feeling of pressure which they can not cure a “falling or dislocation of the womb,” supported by the feelings of the patients, who really believe they have such a complaint, while in the majority of cases, they have only the peculiar sensation of forcing or bearing down, which results often from congestion to the womb,

caused by other inherent chronic ailments. These ought to be ascertained and cured, and the poor sufferer not tormented with endless and senseless supporters and bandages, explorations by speculums, attempts to replace the womb, etc., by practitioners who frequently afterward assert that they have replaced the womb, what really they could not have done, because it was not dislocated. If after all the patient is not cured, and the same feeling returns, it is said, "A relapse has taken place; we must replace it again;" and so the poor sufferer, confiding in her medical attendant, never receives relief. There are, undoubtedly, cases of falling of the womb, which require manual replacement; but these occur mostly after mechanical injuries, falls, blows, missteps, etc., or after severe labor in childbirth, and an examination gives unequivocal signs of dislocation. To such cases, the above remarks have no reference. I have seen frequent cases of the former kind, however, which were pronounced and treated by eminent physicians as prolapsus uteri, where no dislocation whatever existed; in women, too, who never had children, and who had not suffered external injuries. These patients were cured without external helps of pessaries and supporters, and became entirely healthy by a rational internal treatment, after their several chronic maladies, congestions to the womb, neuralgias, and indurations of the womb, etc., had been removed. Yet they frequently would insist upon it, that they had a falling or dislocation of the womb; judging from their own feelings, and having been indulged or strengthened in them by their former physicians.

**TREATMENT.** — If a falling of the womb, or its dislocation is certain, from the preceding external injuries above stated, and the consequent constant feeling of downward pressure, less when lying, worse when moving or walking, compelling her to lean forward; the patient ought to lie



down, and remain quiet, in a horizontal position, sometimes for weeks. During that time, let her take,

*Belladonna* and *Sepia*, alternately, every other morning a dose (six glob.) until better, or until one week has elapsed, when she discontinues the medicine for one week, yet remains lying; after which, she may repeat the same, or take,

*Nuxvomica*. Afterward *Calcarea carb.* in the same manner.

Sometimes a dose of *Platina*, *Cocculus*, and *Ignatia* may be given, if the patient exhibits great nervousness, with constipation.

DIET.—The usual homœopathic diet in chronic diseases.

#### LEUCORRHEA. FLUOR ALBUS. (*Whites.*)

This troublesome and weakening complaint consists of a discharge of mucus, variously colored, and of different consistency, from the private parts. It occurs, generally, between the age of puberty and the critical period, and is seldom seen later than this, except when discharges of this kind are excited, in consequence of the disorganization of the womb. If it manifests itself in children, or even infants, it is either on account of want of cleanliness of these parts, or local irritations, such as pin-worms (*ascarides*), etc.

Weakly females, of a nervous, relaxed, or excited temperament, are more prone to it; and the more our present state of society becomes over-civilized, with its legion of pleasures, inactivity of body, idle and late hours, bad literature, and immediate use of tea, coffee, and other spices (we mention here, only as an instance, the increased use of vanilla), the more easily will this disease be engendered.

It would carry us too far, to go into the practical detail of a disease which requires the most skillful attention of a

medical attendant; and we earnestly recommend an application to him, in an early stage of the disease.

Besides this, we recommend cold water, in all its various applications, as the best means to restore the tone and strength of all the weakened parts of the system.

The principal medicines to be taken domestically, are :

*Pulsatilla*. Discharge thick, like cream, sometimes creating an itching around the affected parts.

*Cocculus*. Discharge of a reddish hue before and after menstruation, with colic and flatulency.

*Sepia*. After *Pulsatilla*; parts become excoriated; discharge yellowish, greenish, fetid.

*Calcareo carb.* *Whitish, corrosive discharge* in children; in adults it is milky before menstruation or after lifting; particularly in fat, corpulent females.

*Sulphur*. If none of the above remedies should prove sufficient.

ADMINISTRATION. — Four doses of a remedy selected, every evening one dose (six glob.); then abstain from taking medicine for four days, and if not better at the end of that time, select another remedy, and take it in the same manner.

DIET AND REGIMEN. — Diet must be nourishing, but not flatulent (see “Dietetic Rules.”) Avoid the causes which excite this complaint; particularly colds and excitements of any kind.

#### PREGNANCY.

As this state of the female is an entirely physiological one, it must not be considered abnormal of itself. It only becomes so, by deviating from the rules of health. As regards diet, a pregnant female may live as she has found it for herself most beneficial heretofore. She must ab-

stain, however, from excesses in diet, which previously would not have injured her, for the sake of the welfare of a being, bound to her by the most endearing ties; the moral and physical health of this expected offspring depends upon its mother's actions.

Be careful in all things—careful in not making a misstep, in not dressing too tight, in not sitting up too late at night; in walking too great a distance, and in eating and drinking, be careful and moderate. But enjoy life, as far as is possible within these conditions; and the reward will be the happiness of others.

Exercise in the open air is good and necessary, but it must not be too violent; for instance, on horseback, etc.; walking in the open air is the best. Do not walk out immediately after dinner; but early in the morning or at night. Do not take purgative medicines.

We will now consider the most common ailments during pregnancy.

#### DISEASES DURING PREGNANCY.

Pregnant females are subject to diseases which, when not in that condition, would not befall them. These we will treat of in particular; while, for other ailments, we refer the reader to their respective headings, giving here only the most suitable medicines:

For *moral affections* (see “Affections of the Mind”):  
*Aconite, Pulsat., Bellad., Platina, Lachesis, Stramon., Verat.*

*Headache: Belladonna, Bryonia, Nux vomica, Platina, Veratrum, Cocculus.*

*Toothache: Chamomile, Belladonna, Pulsatilla, Mercury, Sulphur, Sepia.*

*Convulsions and spasms: Belladonna, Hyosciamus, Ignatia, Chamomile, Platina.*

*Dyspepsia: Nux vomica, Pulsatilla, Sepia, Phosphorus.*

*Constipation* : *Bryonia*, *Nux vomica*, *Opium*, *Lycopodium*, *Sepia*.

*Dysury and strangury (scanty and painful urination)* : *Nux vomica*, *Pulsatilla*, *Cocculus*, *Sulphur*.

*Diarrhea* : *Antimon. crud.*, *Phosphorus*, *Pulsatilla*, *Mercury*, *Sepia*.

*Piles (hemorrhoids)* : *Nux vomica*, *Ignatia*, *Aconite*, *Sulphur*.

*Incontinence of urine* : *Pulsatilla*, *Belladonna*, *Stramonium*, *Cina*, *Silicea*.

*Sleeplessness* : *Coffea*, *Belladonna*, *Nux vomica*, *Hyoscinus*, *Opium*.

*Fainting (hysterics)* : *Ignatia*, *Pulsatilla*, *Aconite*, *Belladonna*.

*Melancholy (low spirits)* : *Aconite*, *Pulsatilla*, *Ignatia*, *Belladonna*.

*Cramps in the abdomen* : *Belladonna*, *Nux vomica*, *Pulsatilla*.

*Cramps in the back* : *Ignatia*, *Rhus tox.*

*Cramps in the hips* : *Colocynth*, *Rhus*.

*Cramps in the legs (calves)* : *Veratrum*, *Secale*.

*Cramps in the feet* : *Calcarea carb.*

*Pain in right side (see article "Liver")* : *Chamomile*, *Aconite*, *Pulsatilla*.

*Heart-burn (sour stomach)*, *Water-brash* : *Nux vomica*, *Phosphoric acid*, *Pulsatilla*, *Sulphur*.

In all the above diseases, compare their respective chapters.

### *Nausea and Vomiting.*

This distressing complaint is present during the pregnancy of some females ; in others it is wanting. It begins usually about six weeks after conception, and lasts up to the fourth, fifth, even seventh month, more or less violently. It is of

a constitutional origin, and can not be taken away easily, as the exciting cause still exists. The same may be said about the spitting of frothy saliva, with which some women are troubled.

Exercise in the open air, cheerful society, strengthening food, and refreshing drinks, such as ale, lemonade, or salt fish, sardines, smoked herring, are frequently means to alleviate the sufferings. The following medicines, also, often have a good effect :

*Ipecac.* If bile is thrown up, coated tongue, loss of appetite.

*Nux vomica.* If with sickness at the stomach there is headache, constipation, pressure in the stomach.

*Arsenic.* Vomiting after eating and drinking; great weakness.

*Pulsatilla.* Sour vomiting; white, coated tongue.

*Petroleum.* (See "Sea-sickness.")

### *Pruritus. (Itching.)*

This distressing disease of the private parts often takes away rest and comfort from the pregnant females, and must be removed as soon as possible. Try first, frequent washings with water, and if not better, wash with a solution of borax and water, three or four times a day.

Internally, take the remedies recommended under this head, in "Diseases of the Skin."

### *Varicose Veins.*

A complaint which consists in an extension of the veins on the lower extremities, owing to the pressure of the pregnant uterus on the large veins of the abdomen, impeding the speedy return of the venous blood upward, creating, thereby, stagnation. If they are not painful and large, frequent washing in cold water is sufficient, as they speed-

ily disappear after the birth of the child. But when they are very large and painful, the patient should lie down for a few days, and apply besides, if necessary, a bandage or laced stocking, to compress the extended veins.

The following remedies can be taken with benefit, every three days one dose (six glob.), changing the medicines every week, until better :

*Arnica, Nux vomica, Pulsatilla, Lycopodium.*

*Spots on the Face during Pregnancy.*

Yellowish or brownish spots on the face, which often appear in pregnant females, require the use of *Sepia*, every eight days one dose (six glob.), until better.

*Miscarriage. (Abortion.)*

Miscarriage can take place at any time between the first and seventh month. If it occurs after the seventh, it is no longer so called, but *premature birth* ; as at this period the child can be saved, and the object of pregnancy be gained—to give birth to a living child. But if the child is born before the seventh month, it can not live ; thence the name “*miscarriage*.” The more advanced the pregnancy is, the less is the danger which might result from a miscarriage. The oftener a woman has miscarried, the more her constitution inclines to new misfortunes of the same kind. Miscarriages are more liable to occur again at the same time of pregnancy, at which the former or last one happened ; if once past that period, the danger to miscarry diminishes. Miscarriage can become a habit ; it sometimes appears epidemically, in so far, at least, as in certain seasons the uterine congestion generally increases (menses appear more frequently and profusely, hemorrhages take



place spontaneously, etc.), and in the same ratio, the possibility and occurrence of abortion multiplies.\*

If a woman approaches the time where formerly she had miscarried, she must be careful not to provoke a return by walking great distances, lifting, running down stairs, riding over a rough road, etc. These practices might excite at any time a miscarriage in females who never were predisposed to it; certainly much more in those already predisposed. A weakening, luxurious mode of living, great mental excitement, are causes of miscarriage, and must be strictly avoided.

The premonitory symptoms of a miscarriage are: chilliness, followed by fever and pressing down sensation, which afterward increases to labor-pains; cutting, drawing, mostly in the loins and abdomen. A discharge of bright-red blood either immediately issues from the vagina, or coagulated, dark blood appears afterward in spells; frequent repetitions of these efforts of nature are usually necessary to expel the foetus; varying in duration from two hours to two days.

TREATMENT.—As soon as the above symptoms appear,

---

\* I observed this phenomenon, on the approach of the Asiatic cholera in Cincinnati in December, 1848, when there was a sensible increase of miscarriages and hemorrhages of every kind, increasing during the spring of 1849, and being at its height in July and August, when the cholera was raging the most fearfully. Women miscarried at this time who never before had had any predisposition to it, although mothers of many children; premature births occurred without any apparent cause, bodily or mentally. After that time, the tide of the disease turned, and with it the abortions decreased in number; also, other kinds of hemorrhages. Undoubtedly, the small quantity of positive electricity in the atmosphere caused a general debility in the nervous system, and consequent muscular and vascular laxity, which even involved, in its general effects, the uterus in all its functions. Hence the epidemical appearance of abortions. Such facts must only be known and recognized as such, and it becomes an easy task to prevent misfortune.

even if in a slighter degree, the patient must lie down and keep perfectly quiet, without moving; if the attack is brought on by mechanical injury—a fall, blow, misstep, or walking, lifting, etc., take

*Arnica*. Twelve globules dissolved in half a teacupful of water, every fifteen or twenty minutes a teaspoonful, until better, or until after one hour another remedy becomes necessary.

*Secale* will be the next remedy to be given, in the same manner as *Arnica*, particularly in females who have miscarried more than once; in older ones, or in those who have a weak and exhausted constitution.

*China*, in alternation with *Secale*, becomes necessary, when the loss of blood is considerable, and weakness and exhaustion evidently increases; buzzing in the ears; cloudiness of sight; fainting when raising from the pillow.

*Ipecac.* becomes a necessary remedy, in alternation with *Secale*, if with *flooding* there is nausea, fainting, cramps.

*Platina* and *Belladonna* in alternation, either at the beginning or, after *Ipecac.* has failed to relieve.

Administration the same as stated in *Arnica*.

DIET AND REGIMEN.—As in “Menstruation too Copious,” “Flooding,” which article may also be consulted, if the above remedies are insufficient for the *flooding*. After the foetus has passed away, treat the patient in the same manner as would be necessary after a regular birth. The patient must keep the bed for the same length of time as in regular childbed. This fact must be well remembered.

Do not neglect to procure a physician, if possible, immediately.

N. B. *Premature birth*, or a birth occurring after the seventh, and during the eight month, must be treated as a regular birth; but frequently needs medical aid, which ought to be procured.

## PARTURITION. REGULAR BIRTH.

This is a physiological and healthy process, setting in at the end of the ninth month of pregnancy, by contractions of the womb, called labor-pain, which, at first light, increases in violence and strength, until the child is expelled. If labor occurs after the ninth month (sometimes it delays for two weeks), it becomes harder on account of the child becoming larger, and its head more closed, consequently less compressible.

*Labor-pains.*

The following medicines will be of benefit to correct the irregularity and painfulness of labor-pains :

For *ineffectual* or *spasmodic* pains : *Coff. Cham.*, *Nux.*, *Sec.*

For the *absence of labor-pains* : *Pulsatilla*, *Secale*.

For *excessively painful labor-pains* : *Coffea*, *Aconite*.

For *sudden cessation of labor-pain*, with congestion to the head, redness of the face, drowsiness : *Opium*.

ADMINISTRATION.—Every half hour a dose (four glob.) until better, or another remedy is chosen.

ADHERENCE OF THE AFTER-BIRTH. (*Placenta.*)

When, after delivery, no more pains appear, to expel the after-birth within half an hour, give *Pulsatilla* and *Secale*, alternately, ever fifteen or twenty minutes a dose (three glob.), until new contractions ensue, after which, the placenta will soon appear. If the patient's head is congested, full, red, give of *Belladonna* four globules in preference to the above remedies.

## TREATMENT AFTER DELIVERY.

Perfect rest and sufficient covering are the first necessities after delivery. No change of clothes or position ought to take

place within the first eight or ten hours after delivery ; during this time the patient generally enjoys a good sleep, into which she may be allowed to fall one hour after delivery ; nothing restores the lost strength better than a good, quiet sleep.

If *flooding* threatens, rub the region over the womb by regular frictions with the hands, until the womb contracts again, and after-pains appear, which diminish the danger of flooding ; or else give her *Belladonna*, if there is a great deal of bearing down ; *Chamomile*, if there is coldness of the limbs and pains around the abdomen.

*China* and *Ipecac.*, in alternation, in the worst cases, when the above remedies do not succeed ; or,

*Pulsatilla*, if the discharge of clotted blood appears at intervals, ceases, and reappears ; also, *Crocus*, *Platina*, and *Sabina*. As the last remedy, apply cold water in wet bandages, which must be frequently renewed, or pounded ice on the abdomen, which allays the flooding soon. (See “Menstruation too Profuse, Flooding.”)

#### AFTER-PAINS.

These are necessary evils, as their beneficial effect in contracting the womb more perfectly is too evident. Yet, they sometimes become harrassing and almost unbearable ; in such cases, use the following remedies :

*Arnica*, as the first medicine, to soothe the irritability of the womb after such heavy exertions ; next, or in alternation with it,

*Pulsatilla*, almost a specific for these pains ; or, if not better,

*Chamomile* and *Nux vom.*, in alternation.

*Secale*, in weakly persons, or those who have already had many children.

*Coffea*, if the pains seem to be so severe that the patient almost despairs.

ADMINISTRATION.—Every half hour or hour a dose (four glob.), until better.

#### DURATION OF CONFINEMENT.

If thus far this process is a perfectly healthy one (and neglect or mismanagement only can make it a source of suffering), its remaining part—that of confinement in bed after delivery—passes off, generally, in the same healthy manner. During this time, usually fourteen days, positively *no visitors* should be admitted *on any consideration*. In the best case, where nothing is wanting in regard to appetite, strength, or milk, and other natural discharges, the patient must not be allowed, before the fifth day, to sit up, and then, during the next five or six days, only as long as is necessary to have the bed made and aired. The reason is, to give the womb ample time to reduce and replace itself in the proper location; and that can *safely* be done only while lying. Attending to this advice, *literally*, may save years of misery. The diet in this time may be more substantial, if wished for, and if the mother and child feel well otherwise.

#### BREAST. NIPPLES. SECRETION OF MILK.

It is of the utmost importance for happiness and health, that the greatest care should be bestowed upon the breasts and their secretions, the milk, the fountain of life for a being so dear to a mother. Months before the birth of the child, the breasts, which sometimes already enlarge at that time, must be prepared for their duty afterward, by washing them every day in cold water, particularly the nipples, so as to harden them for the service to come.

*Milk fever* generally, if it is not too severe, passes off without injury, and requires no other action on our part than to watch the patient closely, day and night, attending

scrupulously and kindly to her wants, because mental excitement at this period often becomes fatal.

If, however, chilliness is followed by a very high fever, flushed face, etc., give *Aconite* in alternation with *Coffea*, if great restlessness is present; every hour a teaspoonful of a solution made of twelve globules in half a teacupful of water. *Bryonia*, *Belladonna* and *Rhus* are afterward necessary, one after the other, every six or eight hours a dose (four glob.), if the pain and swelling in the breast still continues. The child may continue to draw the breast, or it may be drawn by the nurse or a breast pump.

DIET very light; gruel, toast and tea.

### *Sore Nipples.*

The soreness and tenderness, even bleeding of the nipples, require, at first, applications, of

*Arnica tincture*, six drops of it to half a teacupful of water; wash with it every time *after* the child has nursed. If no better in eight days, and the nipples *inflame, swell*, and threaten to *ulcerate*, give

*Chamomile*, particularly when the pains seem to be insupportable, like toothache—every four or six hours a dose (four glob.), until better. If the nipples ulcerate or suppurate, dissolve of

*Silicea* twelve globules in half a teacupful of water, and wash three or four times a day, or after every nursing; if not better in four or six days, take the following medicines, one after the other, in the same manner as *Silicea*, until one relieves, when all medicines must be discontinued until worse again: *Mercury*, *Sulphur*, *Graphites*, *Lycopodium*, *Calcarea carb.*

DIET as usual.



*Ague in the Breast. Gathered Breast.*

Give, immediately, *Chamomile* and *Bryonia* in alternation, every hour a dose, four glob. for four hours, after which discontinue for four hours and let the fever pass off by perspiration, without giving any more medicine ; if the fever will not disappear, however, or returns, give

*Aconite* and *Belladonna*, particularly if the breasts are swollen, hard, and very tender ; externally, apply hot brandy cloth. If lumps remain in the breast, rub with sweet oil, or lay over the breast a plaster of beeswax and sweet oil.

If a *gathering* of the breast can not be avoided, abstain from putting on the warm poultices as long as possible, as they only implicate a still larger part of the breast within the suppurating sphere ; give, during this time,

*Phosphorus* and *Hepar*, in alternation, morning and evening one dose (four glob.), until better or until four doses of each are taken, after which discontinue the medicine, awaiting its effects for at least three or four days ; if no signs of improvement are visible, give

*Mercury* and *Lachesis* in the same manner ; and then again *Phosphorus* and *Hepar*, until the abscess has opened, or the swelling is diminished.

After the abscess has opened and discharged the matter, give

*Silicea*, internally, every evening and morning a dose (four glob.) ; externally in a wash on the breast, three times a day, made of twelve globules of *Silicea* in half a teacupful of water, well shaken. This remedy may be followed in two weeks by

*Sulphur*, internally, same as *Silicea*, and in alternation with it, until the breasts are healed.

DIET, nourishing, but not stimulating.

*Deficiency of Milk.*

Besides using drinks which have a tendency to increase the milk, such as milk-punch (milk with brandy), beer and milk, tea or coffee with milk, give

*Pulsatilla*, *Causticum*, *Calcareæ carb.*, each one for a week, every third night a dose (four glob.)

*Suppression of the Milk.*

If the secretion of the milk is suppressed suddenly, either by mental or physical causes, great injury is to be feared, if the proper measures are not adopted immediately ; one of them is to take

*Pulsatilla*, twelve globules dissolved in half a teacupful of water, every two or three hours a teaspoonful. If congestion to the head, lungs, or abdomen takes place after the milk has disappeared, give *Belladonna* and *Bryonia*, in alternation, in the same manner as *Pulsatilla*.

If *mental emotions* were the cause, give

*Aconite*, *Coffea*, *Chamomile*. (See in "Diseases of the Mind" for the consequences of fear, fright, etc.)

If *taking cold* was the cause, give *Chamomile*, *Bryonia*, and *Rhus*, particularly when the head and limbs ache, in the same manner.

If *diarrhea* sets in, give *Pulsatilla*, *Bryonia*, *Rhus*, *Belladonna*.

*Milk bad, too thin, or repugnant to the Child.*

Give the mother *Cina*, *Mercury*, *Silicea*, every third day a dose (four glob.), each remedy for ten days ; *Silicea*, particularly if the child vomits immediately after sucking.

*Milk, excessive flow of it. Weaning.*

Internally, give *Pulsatilla*, *Bryonia*, *Belladonna*, *Calca-*

*rea*, in succession, each remedy for thirty-six hours, every twelve hours a dose (six glob.) Externally, wash with spirits of Camphor, and wear constantly cotton bathing. This is the best external application which can be used at the time of weaning the child.

*State of the Bowels.*

On account of the great changes going on at this time in the female organism, the secretion from the womb and breasts, whereby a great quantity of liquids is disposed of, the secretion in the bowels by the stool is retarded, and, consequently, no evacuation designed by nature, for at least five days after delivery; if any stool is forced, artificially, before that time, it is that much liquid taken away from places where nature needed it the most. If, after the lapse of five days, no motion on the bowels has appeared, give of

*Bryonia*, four doses every three hours one (six glob.), and await its effects twenty-four hours, after which, give *Nux vom.* in the same manner, if necessary.

*Sulphur* after that, if necessary, in the same manner. An injection, either of luke-warm water and two tablespoonfuls of sweet oil, or the same with a little castile-soap dissolved in it, may be given on the sixth day, if necessary, and as often afterward as needed.

If *diarrhea* sets in, the principal medicines are: *Rheum*, *Phospor. acid*, *China*, *Pulsatilla*. For closer examination of the symptoms and their exciting cause, see article "Diarrhea."

But there is one kind of diarrhea peculiar to nursing women, which is frequently connected with the *nursing sore mouth* (see this below.) The discharges from the bowels are whitish, curled, smelling sour and musty, being copious and frequent, but not very painful. This diarrhea is the beginning of consumption of the bowels, and must be

attended to immediately. I have found the following remedies almost specific in such cases :

*Nux vomica*, and *Hepar*, alternately, every three hours a dose (six glob.), until better, or until six doses of each are taken ; at the same time the patient must lie on the bed or couch.

DIET.—Black tea, dry toast, rice, afterward beef-steak and roasted mutton. (See, besides, “Consumption of the Bowels.”)

### *Sore Mouth of Nursing Women.*

In some females this complaint is constitutional ; it is caused by the peculiar irritation which nursing has upon their digestive organs ; if let alone, it sometimes becomes so bad as to force us to discontinue nursing, or to send the patient to mountainous places (the higher the better), where the digestive organs become stronger and more able to resist this weakening influence, occasioned by the constant loss of fluid (milk.) The following remedies, however, will be used with benefit :

*Mercury*, *China*, in alternation, every evening and morning a dose (four glob.), for a week.

*Nitric acid*, every other evening a dose (four glob.), until better, or for a week ; and, afterward,

*Borax*, *Nux vom.*, *Sulphur* in the same manner. (Besides, see article on “Stomacace.”)

DIET AND REGIMEN. Diet nutritious, but not flatulent, as frequently a diarrhea accompanies this complaint, which requires particular attention. See foregoing article.

Exercise in the cool fresh air (but not fatiguing exertion by walking), riding out in a carriage, etc., are beneficial ; also, cold bathing, or sponging.

## DISCHARGE FROM THE WOMB DURING CONFINEMENT.

*(Lochia.)*

This bloody secretion from the womb is called the "Lochia," and lasts, generally, from nine to fourteen days, becoming gradually paler from the beginning until finally a lightish mucus discharge is left, which diminishes in quantity, until it also disappears.

This is the natural process ; the deviations from it are abnormal, and require attention. If they are *suppressed* entirely, and fever, pain, and congestion follow, particularly in the abdomen and head, give *Bryonia*, *Pulsatilla*, *Hyoscyamus*, *Platina*, *Secale*, *Veratrum*.

(Another means to make them reappear again is, to apply on the womb warm fomentations of hops in a bag.)

If they are *too profuse*, or of *too long duration*, *Ipecac.*, *China*, *Secale*, *Calcarea carb*, *Pulsatilla*, *Bryonia*, *Platina*. If the white lochia becomes bloody again, *Rhus*.

ADMINISTRATION the same as in "Menstruation too Profuse, Flooding."

DIET AND REGIMEN, also, the same.

## PAIN. INFLAMMATION IN THE ABDOMEN.

*Metritis and Puerperal Fever.*

If the abdomen becomes very sensitive to the touch or by movement, shooting pains, fever, headache, give immediately *Bryonia* and *Belladonna*, twelve globules of each dissolved separately in teacups half full of water, every two hours a teaspoonful, alternately ; send for the physician *immediately* ; if the lochia had stopped, apply, externally, a bag filled with hops dipped in hot water and wrung out well. If perspiration ensues, encourage it until the pain has disappeared.

## GENERAL COMPLAINTS DURING CONFINEMENT.

For *sleeplessness* : *Coffea*.

For *colic* : *Chamomile*, *Bryonia*, *Pulsatilla*, *Sepia*, *Veratrum*.

For *convulsions* (spasms) *Ignatia*, *Hyoscyamus*, *Platina*.

For *retention of urine* : *Nux vom.*, *Belladonna*, *Pulsatilla*.

For *debility* : *China*, *Phosphor. acid*, *Veratrum*, *Calcarea carb.*

For *falling off of the hair* : *Lycopodium*, *Calcarea*, *Natrum mur.*, *Sulphur*.

For *white swelling* (milk leg) : *Belladonna*, *Rhus*, *Lachesis*, *Arsenic*, *Sulphur*.

N. B. For further information and the administration of medicines in the above diseases, see their respective chapters.



## CHAPTER XVII.

### TREATMENT OF CHILDREN.

#### APPARENT DEATH OF A NEW-BORN INFANT. (*Asphyxia*.)

When a new-born infant, either from exhaustion or other causes, does not breathe or very imperfectly, order a warm bath, and while that is being prepared, rub the spinal processes, from the neck down to between the shoulders, up and down, gently but firmly for some time, and blow the breath gently into its mouth from time to time, if it does not breathe.

In the bath, pursue the same manipulations, and give, a few drops of a solution of *Tartar emetic*, one grain in half a teacupful of water; also, give an injection of it every fifteen or twenty minutes. If the child looks blue in the face (strangled), give *Opium*, in the same manner prepared, on the tongue, and by injection.

After the child revives, but looks pale and exhausted, give *China*, one glob. every two hours, for six hours.

DIET OF A NEW-BORN INFANT.—After washing and dressing, the infant wants sleep, into which it falls as soon as it is warm enough. Put it in the mother's bed—near her, if possible. After having slept for six or eight hours, lay it on the breast, even if no milk were there, and feed it afterward with molasses and water (under no circum-

stances bread and water), or milk and water, half and half, and sweetened with loaf sugar.

Wash its mouth after feeding every time with cold water; it prevents the *thrush*.

#### COLIC.—CRYING OF INFANTS.

Examine well whether pins, sticking the little sufferer are the cause.

If no cause is apparent but the universal one (colic), give *Chamomile* and *Belladonna*, in alternation, every half hour or hour a dose (one glob.) on the tongue.

If the child bends its body double while crying, and retracts its thighs, give

*Chamomile*, when the face is red; or *Belladonna*, when the face is pale. If the child has greenish stools at the same time, give *Chamomile*. If it has loose evacuations, of a *sour* smell, give *Rheum*. If these will not suffice,

*Ipecac.* and *Jalappa*. All in the same manner. For great restlessness, sleeplessness, and feverish heat, with crying, give *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, in the same manner.

#### ELONGATION OF THE HEAD.

This deformity of the infant's head is only temporary; wash it with *Arnica tincture* and water (six drops to a tea-cupful); in a few days it will disappear.

#### SNUFFLES — OBSTRUCTION OF THE NOSE. (*Coryza*.)

Give *Nux vomica*, one glob. in the evening, and *Sambucus* the next evening, if not better; in that manner alternate, until relieved.

If the nose runs water, give *Chamomile*.

#### SWELLING OF THE BREASTS IN INFANTS.

Do not press or handle rudely these delicate swellings;

as they contain nothing, the least of all milk, to be squeezed out. It is an inflammation, and must be treated like any other inflammation of glands. First, we should try to reduce the swelling; which can be accomplished, in most cases, by covering it with a lint, dipped in sweet oil. If this will not suffice, wash it several times in warm brandy and water, and give the child internally *Chamomile* and *Belladonna*, alternately, every evening and morning a dose (one glob.) If it still grows larger (I have seen them of the height of one inch and more), keep on it a bread and milk poultice, and after the gathering opens, treat it like any other abscess; inwardly give *Mercury* and *Hepar*, every evening one glob., alternately.

#### RESTLESSNESS. SLEEPLESSNESS.

If from any apparent cause, the child can not sleep, give it *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, alternately, every hour one glob.

#### INFLAMMATION OF THE EYES.

This is a frequent complaint among infants. Do not expose the eyes to a light too strong, neither let it take cold in drafts of air. Drop a few drops of the mother's milk into the eyes several times a day. Besides, give *Aconite* and *Belladonna*, in alternation, three times a day one glob., for several days.

If not better, give *Euphrasia* and *Rhus*, in the same manner; and, at last, *Sulphur*, every other evening one glob., for a week.

#### THRUSH, OR SORE MOUTH OF INFANTS. (*Aphthæ*.)

Wash immediately with *Borax*, dissolved in water and sugar (do not rub with the dry *Borax*); cleanse well all spots with a cloth dipped in the solution, and then wash the

mouth well with another clean cloth, dipped in clean, cold water. If we use *Borax* in the dry state, too much of it remains in the mouth, creating there the same disease which we tried to cure ; because it is the true homœopathic remedy ; in the above described manner, however, cases of thrush have been cured, which were previously treated for a length of time with pulverized *Borax*, without the least beneficial effect. At the same time, give the child *Mercury* and *Sulphur*, every three or four days, alternately, one glob.

#### HEAT. RED GUM. HEAT SPOTS.

In most cases, they disappear quickly, without medicine, if the child is not kept too warm, and bathed regularly.

If the child is restless with them, give *Chamomile*, *Aconite*, *Belladonna*, *Rhus*, *Arsenic*, every evening one glob., and every day another remedy, until relieved.

#### EXCORIATIONS. RAWNESS OF THE SKIN.

This happens mostly in very fleshy infants.

*Chamomile*. Every evening one glob., for three or four days ; then discontinue the medicine for four days.

*Mercury*. In the same manner.

*Sulphur*, *Lycopodium*. In the same manner.

Wash often, and dry well ; after which, use externally fine wheaten starch.

#### JAUNDICE.

Often occurs when children have taken cold, or have been purged too frequently with castor oil.

*Chamomile* and *Mercury*, in alternation, every evening one glob., until better.

*Nux vomica*. In the same manner, if the child is restless and costive.

*Hepar.* If the above remedies do not suffice, every other evening one glob.

#### ERYSIPELAS.

(See “Erysipelas,” under “Diseases of the Skin.”)

This disease claims the attention of a homœopathic physician. Yet the following remedies may be given, until one can be procured.

*Belladonna* and *Aconite*, alternately, every three or four hours one glob. ; after the fever has subsided

*Belladonna* and *Rhus*, in the same manner.

The child must be kept dry during this disease, and lint may be applied to the parts affected.

#### CONVULSIONS OR SPASMS.

Order immediately a warm bath, and while that is preparing, select from among the following remedies that which is suitable.

*Belladonna.* Almost in all spasms the first remedy. Give, in preference, the *Belladonna*<sup>cc.</sup>, in the following manner: put directly two glob. of it on the tongue of the child; then dissolve twelve glob. in four teaspoonfuls of water, mix well, and give every fifteen minutes a few drops on the tongue.

After the bath (which may last from five to ten minutes), wrap the child, without drying off, in heated flannel, and lay it with its head higher on a pillow made of quilts, and, if possible, covered with oil-cloth.

If the spasms will not cease, keep the child's feet and legs in warm water, at the same time pouring a stream of cold water on the crown of its head, until consciousness returns. This process must be *repeated frequently*, although the first attempts have proved inefficient; it has been tried with the greatest success.

If by this time the child has not yet recovered from the fit, order an injection of lukewarm water, and a tablespoonful of sweet oil, together with a little soapsuds (castile soap) to be given, and select another remedy, which administer in the same manner as *Belladonna* <sup>cc.</sup>.

*Chamomile*. Convulsive jerkings, moaning, eyes half open, *redness of one cheek*.

*Cina* <sup>cc.</sup>. If worms are to be suspected.

*Coffea* <sup>cc.</sup>. In weakly children, subject to frequent fits.

*Ignatia*. When the fits return in regular intervals.

*Ipecac*. When the breathing is short between the fits, nausea, and frequent spasmodic stretchings.

*Opium*. In fits after *fright*, when the breathing is labored, the face dark, flushed, almost blue.

If after the fit is over the child sleeps, but with its eyes half open, give *Hellebor.* and *Belladonna*, as directed under "Hydrocephalus," page 388.

#### TEETHING. (*Dentition.*)

With seven months, the two middle teeth of the lower jaw appear; shortly after, the two corresponding ones in the upper jaw. From this time, the little organism is constantly putting forth teeth, until, at the end of two or two and a half years, the first dentition, consisting of twenty teeth, is completed.

The process of teething excites in the little ones a variety of diseases, which, if not well treated, too often prove fatal. The mother can do a great deal, in directing the mode of her living, in eating, drinking, and acting, so as not to make her milk another source of suffering for the sickly sufferer. (See "Hygiene.")

The gums swell and are painful; yet the child wants to bite and press something hard on the teeth, to relieve the intolerable itching and irritation underneath the gums. Give



it an ivory ring. If fever appears, with restlessness, retching, etc., give *Coffea* <sup>cc.</sup> and *Belladonna* <sup>cc.</sup>, alternately, every half hour or hour half a teaspoonful (twelve glob. having been dissolved in half a teacupful of water.)

*Chamomile* and *Belladonna*. If convulsive jerking or twitchings occur in the sleep.

*Cina*. If with teething it has a dry, spasmodic cough, or signs of worms—rubbing at the nose.

*Aconite*. If the fever is high.

If a diarrhea of a yellowish color occurs, it is not objectionable at first; but if it becomes of a whitish, slimy color, and curdled, give *Coffea*, *Ipecac*, *Calcarea carb.*

If *convulsions* ensue, treat as stated under that article—page 385.

If the teeth are tardy in breaking through, do not lance them (it can only be of use where they are much swollen and *heated*), but give *Calcarea carb.*, every evening two glob., for a week; then discontinue a week, and give *Sulphur*, in the same manner.

If *constipation* occurs during teething, give *Bryonia* and *Nux vomica*, alternately, every evening a dose (two glob.) for six days.

#### CONSTIPATION.

It is a bad practice to give children physic, castor oil, etc., rather than to take the trouble to prepare an injection for them, which is easily done; as it consists only of warm water, with a little piece of castile soap dissolved in it, and sweet oil. This is much better to apply than to give children something, which, if it for the present relieves them, renders them afterward more constipated than ever. First give them the following medicines:

*Bryonia*, *Nux vomica*, and *Opium*, in succession, each remedy for one day, and three doses (two glob.) of

each in a day. If not better, give *Sulphur*, in the same manner; and then, if necessary, give the above injections.

Besides, allow as much cold water as they will take, and rub their stomachs every evening well with sweet oil.

#### DIARRHEA OF INFANTS.

If it has a *sour smell*: *Rheum*.

If with *colic* and a red face: *Chamomile*.

If with *colic* and a pale face: *Belladonna*.

If not relieved after these, give *Sulphur*.

If the diarrhea always appears in the heat of summer: *Ipecac*, *Nux vomica*, *Bryonia*, *Carbo veg*.

If from *cool weather*: *Dulcamara*, *Antimon. crud*.

If with great emaciation and weakness: *Arsenic*, *Carbo veg*.

ADMINISTRATION. — After every operation, one glob. of the selected medicine. For further information, see “Diarrhea.”

#### FEVERS OF INFANTS.

Always give *Aconite*, two glob. first; in two or three hours, *Chamomile* and *Coffea*, in alternation, every one or two hours two glob. (or in water, as usual); and lastly, *Belladonna*, in the same manner.

By that time, the fever must have shown its character, or the child will be well.

DIET. — Give a thirsty child as much cold water as it wants.

#### DROPSY OF THE BRAIN. (*Hydrocephalus*.)

Children with large heads and of a scrofulous taint are more subject to this disease than others, as their brains are more liable to congestions. The most trivial derangement of the bowels, sudden changes in the atmosphere,

teething, eruptive and other fevers; in fact, all diseases which may befall children, can, under certain circumstances, take their final and fatal issue through the dropsy of the brain, if timely aid does not prevent such a result.

DIAGNOSIS.—Whenever a child becomes drowsy, its head hot, feet cold, with or without nausea and retching, and *sleeps with eyes half open*, be careful and give it immediately the following remedies:

TREATMENT.—*Bellad.* and *Hellebor.* (of each twelve glob. in half a teacupful of water), every hour or two hours a teaspoonful, until four teaspoonfuls of each are given, or until the child becomes more lively. If the same drowsiness reappears, repeat the same medicine; if the third time, it appears, give *Opium* and *Sulphur*, in the same manner, and afterward *Belladonna* and *Hellebor.*, again as above.

DIET must be light. Keep the feet warm, but the head cold, with cloths dipped in ice water.

It is evident, that in diseases of this kind, medical aid should be obtained as soon as possible.

#### ASTHMA OF CHILDREN. (*Choking Fits. Asthma Millari.*)

If little children seem almost to suffocate, fall into a spasm, and have a bluish face, give *Ipecac.*, one glob. every ten minutes; and when it is characterized more by hard and tight distention of the stomach and around the short-ribs, with *shortness of breath, choking*, anxiety, agitation and tossing, cries and retraction of the thighs, give *Chamomile*, one glob. every twenty or thirty minutes, until better.

But if an *asthmatic* attack occurs suddenly and violently in the sleep, with dry, dull cough (resembling croup), the face and extremities become purple, spasms in the hands and feet (*asthma Millari*), give *Sambucus*, every five or ten minutes one glob. until better. Send, however, for a physician directly.

If *Sambucus* should not relieve within two hours, and no physician could be had, give *Ipecac.*, *Pulsatilla*, *Arsenic*, *Sulphur*, *Cuprum*, *Spongia*, in succession, in the same manner as *Sambucus* above.

DIET, of the lightest kind. No meats, but gruels.

#### REMITTENT FEVER OF INFANTS.

DIAGNOSIS. — Languor, irritability, nausea, want of appetite, thirst, slight heat of the skin, and restlessness at night. This is the beginning of the disease above named. Soon, the symptoms increase ; fever ; constipation, or diarrhea of a mucous, fetid substance, sometimes mixed with blood ; heat in the body and head ; extremities cool ; tongue coated, dry, and red on the margins ; drowsiness, listlessness in the day, restlessness at night ; hacking cough.

TREATMENT. — *Ipecac.* As the first remedy, every three or four hours one glob., for one day ; afterward, *Bryonia* and *Rhus*, alternately, every three hours one glob. for two days ; then discontinue the medicine for one day ; after which, if the patient is better, *Sulphur* may be given once or twice, every three hours one glob. ; or, if the head suffers the most, give *Belladonna*, every three or four hours one glob. for one day ; or *Chamomile*, if the fever continues, with one flushed cheek, the other is pale ; irritability ; child does not know what it wants ; in the same manner as *Belladonna* ; or give

*Mercury.* If a diarrhea with tenesmus is predominant, and the tongue coated whitish, or

*Nux vomica.* When constipation is present, with frequent but ineffectual desire.

*Lycopodium.* When the tongue is dry, yet the patient is not thirsty ; very petulant ; does not want to see any one, or talk to any one.

If head symptoms appear, see “Dropsy of the Brain.”

DIET AND REGIMEN. — Gruels, toasted bread, soaked in

milk and water ; no meat or broth ; no eggs ; keep them comfortably warm, always covered, particularly arms and limbs. If perspiration appears, do not check it by exposure, as it frequently breaks the disease, and shortens its course.

In convalescence be careful in the diet, as relapses frequently take place from errors in the diet. See what is said about it in "Typhus."

#### VACCINATION.

It is in accordance with the homœopathic principle—*similia similibus*, or a remedy which produces a similar (but not the same) disease in the healthy—that vaccination can save from an attack of small pox, the latter being a similar (but not the same) disease to the former. Through vaccination the triumphs of homœopathy have been shown to the world by innumerable blessings, in arresting such a loathesome disease as small-pox.

Vaccination, if salutary and truly protective, must not be negligently applied. The virus, or matter, ought to be taken from the cow itself, or from a healthy child, whose parents are healthy too, and in whose family skin diseases or scrofula are not hereditary.

Persons can be vaccinated at any time, from the first hour of their existence to any age after that.\* The best age for vaccination is from six months to one year ; the best time in the year is May or June, when the least sickness generally prevails.

If a child has been vaccinated with bad matter, or scrofulous symptoms develope themselves, give *Sulphur*, every evening one glob. for eight days.

---

\* I was obliged, once, to vaccinate a child only half an hour old, whose mother, at the time, was seriously ill with the varioloid ; the vaccination took well, without rendering the child more sick than common, saving it successfully from an attack of small-pox.

## CHOLERA INFANTUM.

DIAGNOSIS.—Violent, copious vomiting, first of food, afterward of sour liquid; diarrhea mostly at the same time; restlessness, child tosses from one side of the bed to the other; nausea; retching on the least movement, or after drinking water, which is soon thrown up again; head hot; extremities cool; thirst great, drinks greedily; very weak; eyes sunken, and half open; eyelids heavy; pulse feeble, or none at all.

This disease occurs, usually, in the summer, after errors in diet and other weakening influences on the stomach.

TREATMENT.—In nowise different from *cholera morbus* in adults.

*Ipecac.* and *Veratrum* (twelve glob. dissolved in half a teacupful of water), give every fifteen or thirty minutes a teaspoonful, until the severest symptoms have diminished, when the intervals ought to be lengthened. If not better after two hours, give

*Arsenic* (prepared in the same manner) every half hour a teaspoonful, until better.

*Rhus*, in alternation with *Arsenic*, if the child tosses about in the bed, in the same manner.

If the cholera disappears, but drowsiness ensues, child sleeps with eyes half open; head is hot; feet cold; treat as stated under "Hydrocephalus," page 388.

DIET AND REGIMEN.—Cold water; gruel; in convalescence great care must be taken in regard to diet.

## SUMMER-COMPLAINT.

This disease is well known in the large cities, where it creates a fearful mortality among children, which are yet under allopathic treatment. Teething and the warm weather are its prominent causes, which are sufficient to



produce all the subsequent changes, which make up the so-called summer-complaint. In general, careful attention to diet, bathing in cold or salt water, cool, refreshing air, on hills, or mountains, or in the high country, will do much to prevent or cure this disease. But where it has begun, give the following medicines :

*Ipecac.*, for nausea, vomiting of food and bile ; fermented stools, with white flocks, tinged with blood ; no appetite ; great thirst.

*Bryonia*. Diarrhea in *hot* weather ; great thirst ; vomiting after eating ; stools smell putrid, are white or brownish. (In alternation with *Ipecac.*)

*Carbo veg.* after *Bryonia*, if the discharges are very thin and offensive, with pain at the time.

*Dulcamara*. Diarrhea in *cool* weather.

*Antimon. crud.* Nausea ; vomiting ; tongue *coated* ; stools offensive ; urinating frequently.

*Tartar. emet.* Nausea : gagging, tongue *clean* ; stools watery, offensive ; child is very weak. (In alternation with *Bryonia*.)

*Mercury*. Stools, with straining (like dysentery), and colic pain, discharges greenish, bloody, slimy. (In alternation with *Nux vom.*)

*Veratrum*, if the disease assumes the character of cholera infantum. (See this article.)

ADMINISTRATION.—Every six or eight hours, four glob. *Phosphor. acid* and *China*, in alternation, after every evacuation two glob., if great weakness predominates and the stools are frequent, yellowish, watery. (See “Cholerina.”)

If the child becomes still weaker, and, finally, a thrush, white sores appear in his mouth, give *Mercury*, every two hours four glob., until better, and in a few days afterward, *Sulphur*, two glob. evening and morning.

DIET AND REGIMEN.—Arrow-root, farina, fried sweet-pota-

toes, sweet pickled pork, fried smoked herring; if possible, retire with the child to a place in the country high and airy, where the child can be carried out mornings and evenings under the shade of the trees — here it will soon recover.

## PART SECOND.

---

I. ANATOMY AND PHYSIOLOGY.

II. HYGIENE AND HYDROPATHY.

III. MATERIA MEDICA.



## I. ANATOMY AND PHYSIOLOGY.

ANATOMY, or the art of dissecting as applied to the human body, describes the mechanism and structure of the parts of the human system, while

PHYSIOLOGY treats of the laws by which the organism is governed and the various functions in man are performed.

From the above is seen how closely these two sciences are related to each other, and that it is almost impossible to treat of them separately, without great disadvantage. They will appear, in this short treatise, interwoven with each other, as the necessity of a clear exposition of their details may require.

The study of anatomy and physiology must be, to every one, interesting, who wants to know a little more of the wonderful creations and provisions of an allwise Providence, than an outside view of nature around him can give. But it is not alone the thirst for increased knowledge which is satisfied by studying the human system and its laws; the latter is often highly beneficial to our physical welfare, and is particularly necessary for those who, from necessity or philanthropy, undertake to minister to the sufferings of their fellow-beings: and to this category not merely the professional physician belongs, into whose hands life and health are trusted, but all men, if possible, should enrich

their minds with the treasures of a science whose teachings lighten, to so great an extent, the burden and responsibility of the healing art; particularly as no one knows whether he might not be called upon, under forcing circumstances, to practice medicine to the extent of his knowledge. Viewed in this light, it becomes the duty of every one who prescribes at all in diseases, to make himself acquainted with the human system and the laws of health, as knowledge only enables him to decide competently where health ceases and disease commences. For a successful and satisfactory use of the prescriptions laid down in a domestic physician, some knowledge of the anatomy and physiology of the human system is indispensable, and we recommend, strongly, a repeated and careful perusal of the following pages.\*

#### STRUCTURE OF THE HUMAN ORGANISM.

The human body consists of fluids and solids, changing constantly from one into the other; their proportion to each other varies in different individuals and at different periods of life. In youth, the fluids are more abundant than in advanced age. The liquids contain, as it were, the whole body in its elementary particles, which, by organical attraction, are formed into the different solid parts of the system. These are composed of different degrees of solidity, as their intention for use requires it. The simplest form of organized animal substance is a *membrane*, or *tissue*, composed of fibres interwoven like a net-work; all organs are formed by tissues, which are different, and adapted to their uses.

---

\* If a more extended acquaintance is desired with these necessary branches of practical medicine, it can satisfactorily be found in the work of Calvin Cutter, M. D., on anatomy, physiology, and hygiene, which we followed in its admirable and popular arrangement.



The *mucous* membrane lines all the cavities which communicate with the air, as the mouth, nostrils, intestinal canal, lungs, etc., and is filled with small glands, which secrete a viscid fluid, called *mucus*, to protect the inner surfaces of the cavities from the contact with the air.

The *serous* membrane lines all the cavities which are closed, like a sack, as the chest, joints, heart, and lungs, in which these organs lie free, yet closely invested. A *serous* fluid is secreted between the sides of these sacks to keep them moist and movable.

The *cellular* tissue extends throughout the whole system, and fills up space wherever it exists, between muscles or within them, even within the nerves themselves, yet never adheres to them, to impede their action, which is accomplished by the secretion of a fluid.

The *adipose* tissue forms distinct bags or cells, filled with *fat*, and is principally located in the skin, abdominal muscles, and around the heart and kidneys; there are none to be found in the brain, ear, and nose.

The *dermoid* membrane forms the outside covering of the body, called the *skin*, and is similar in its structure, (although harder), to the mucous membrane, of which it forms the external continuations. It secretes, constantly, a fluid called perspiration, if it appears in larger quantities; it exists mostly, however, in an imperceptible gas, which, as it were constitutes the atmosphere of the body.

The *fibrous* tissue forms a thin, dense, strong membrane, and is found where a strong elastic action or protection is needed, as in the lining of the internal surface of the skull, around the bones and at the end of the muscles; here it constitutes the so-called *ligaments* and *tendons*.

The *cartilagenous* tissue forms the ends of the bones, where they concur in forming a joint, and is, on that ac-

count, firm, smooth and elastic. It is the hardest part of the body except the bone.

The *osseous* tissue, which composes the *bones*, varies in different periods of life as regards solidity and density.

The *muscular membrane* consists of many filaments, which united, form fibres, each of which is inclosed in a fine layer of cellular membrane. *Muscles* are composed of bundles of these fibres.

The *nervous* tissue is composed of soft, pulpy matter, inclosed in a sheath, each of these, containing one nerve, is formed of two distinct substances, one pulpy and grey colored, the other fibrous and white. This arrangement is found in the smallest nerve as well as in the largest.

#### ELEMENTS OF THE HUMAN ORGANISM.

These are: first, *inorganic* (*chemical*); or, second, *organic* (elementary products of the system itself.)

1. The *inorganic* or *chemical* elements. These are:

a. *Metallic* substances, as Potash, Soda, Lime, Magnesia, Alum, Silix, Manganese, Copper, and Iron.

b. *Non-metallic* substances, as Oxygen, Hydrogen, Carbon, Nitrogen, Phosphor, Sulphur, Prussic Acid, etc.

2. The *organic elements* are: Albumen, Fibrin, Gelatin, Mucus, and Osmazome.

#### ANATOMY OF THE BONES.

The bones, giving strength and solidity to the system and its movements, are so united among themselves, and adapted to each other, that they admit of the most numerous and various actions. The elevations or protuberances of the bones are called processes, which generally form the points of attachment for the muscles and ligaments. They are composed of earthy and animal matter; the former giving strength, the latter vitality. At the earliest stage

of formation, the bones are cartilaginous, soft, and tender, and become hard and ossified as soon as deposits of phosphate and carbonate of lime commence, which are called *points of ossification*.

The *periosteum* or fibrous membrane, investing the bones, except where they are tipped with cartilage, has minute nerves, and gives vitality and nutrition to the bone.

There are two hundred and eight bones in the human body, beside the teeth. They are divided into four parts :

First. The bones of the head.

Second. Of the trunk.

Third. Of the upper extremities.

Fourth. Of the lower extremities.

1. *The bones of the head.* They consist of those of the *skull, ear, and face*.

The bones of the *skull*, of which there are eight, consist of two plates or tablets of bony matter, united by a porous portion of bone ; the external tablet is fibrous and tough, the internal, dense, hard, and glossy. Thus, the skull is admirably adapted to resist the penetration of sharp instruments, or diminish any vibration occasioned by falls and blows.

The skull is convex externally, and at the base much thicker than at the top or sides ; its bones are, as it were, sewed together, united by sutures, whose rugged edges interlock with each other, producing a union, called, by carpenters, *dovetailing*. By this wonderful arrangement, the vibrations from external injuries are interrupted, and fractures prevented from extending as far as they otherwise would, in one continued bone.

In early infancy the bones are not united, leaving interstices of considerable extent, which fill up slowly with bony matter ; thus allowing, in the early time of infancy, sufficient room for the expansion of the brain, which, in

this time of life, is particularly liable to destructive congestions. No part of the human system shows the distinctions in form and features, between the different races of mankind, more than the skull ; and here, too, are located those guides of phrenology, the bumps of the brain, which show, by their position or relative eminence, the propensities of the individual.

In each *ear* are four very small bones, which aid in hearing.

In the *face* we account fourteen bones, some of them serving for the attachment of those powerful muscles for masticating food, others to retain in place the soft parts of the face. This forms the most interesting part of the human system, the face, with its rays of intelligence and joy, or its clouds of distress or sorrow ; besides the wonderful play of passional expression in the features of the face, which is thus made the dial-plate of the inner man.

2. The *trunk* has fifty-four bones : viz., twenty-four ribs, twenty-four bones in the spinal column (back-bone), four in the pelvis (forming the hips), the breast-bone (sternum), and the bone at the base of the tongue (*os hyoides*). They are so arranged as to form, with the soft parts attached to them, two cavities, called the *thorax* (chest) and *abdomen*.

The *thorax* is formed by the breast bone in front, the ribs at the sides, and the twelve dorsals of the spinal column in the back ; this cavity contains the lungs, heart, and large bloodvessels ; its natural form is a cone, with its point above, yet fashion often inverts this order boldly, but never without paying the due penalty in diminution of life and happiness.

The *breast-bone* (sternum) consists of eight pieces in a child, which after uniting, form but three parts in adults ; the lower part of the sternum reaching over into the pit of

the stomach, remains cartilaginous and flexible until old age, when it is often converted into bone.

The *ribs* are connected with the spinal column, and increase in length as far as the seventh, where they become shorter. The seven upper ribs are fastened to the sternum by cartilages, to facilitate their motions in breathing; they are called the *true ribs*. The next three are united to each other by cartilages not touching the sternum — therefore called *false ribs*; while the latter two, or lowest ribs, are not connected either with the sternum or the other ribs, therefore, called *floating ribs*.

The *spinal* column contains twenty-four pieces of bone, called *vertebræ*. Each of the *vertebræ*'s has seven projections, called *processes*, four of which, called *articulating* processes, join similar ones of the adjacent *vertebræ*'s to form the column; two of the remaining processes are called *transverse*, and the other the *spinous* process; these last three receive the muscles of the back, to allow a firm and elastic motion to the spinal column in all directions.

A canal or tube runs through all the *vertebræ*'s in one continuous descent from the round aperture in the skull, to contain the *spinal cord* (*medulla spinalis*), the immediate continuation of the small brain.

Between the *vertebræ* is a cartilaginous and highly elastic substance, which facilitates the movements of the spine, and diminishes any shock in walking, running, or leaping, which would otherwise hurt the spine or brain.

The *pelvis*, or the bones forming the *hip*, consists of four parts, two of which are called *innominata* or *nameless* bones; these, in particular, form the hip, having externally, at each side, a deep socket like a cup, for the reception of the round head of the thigh bone; internally these bones, in conjunction with the sacrum, form a cavity, which contains a great portion of the abdominal organs, as the bladder, rectum, sexual organs, etc. The *sacrum* belongs

to the pelvis, forming its posterior part; it is placed between the hip-bones, to which it is bound by ligaments; it is, in reality, the continuation of the vertebra, and forms part of the spinal column, which is terminated, finally, by a small appendix called *coccyx*, which, in infants, consists of seven pieces, unites, however, in youth, forming one bone, which, in females, is more movable than in males, for a wise purpose.

3. The bones of the *upper* and *lower limbs* are enlarged at each extremity, called projections, to which the tendons, or muscles, and ligaments are attached. The bones of the extremities are mostly in the form of cylindrical and hollow shafts.

The *upper extremities* contain sixty-four bones, as the shoulder-blade (*scapula*), collar-bone (*clavicula*), upper arm bone (*humerus*), two bones of the forearm (*ulna* and *radius*), the wrist (*carpus*), the *metacarpus* (palm of the hand), and the fingers and thumb (*phalanges*.)

The *collar-bone* (*clavicula*) is attached at one end to the breast-bone, at the other to the shoulder-bone: it is shaped like the italic *f*, and its use is to prevent the arms from sliding toward the breast; a fracture of this bone requires a more complicated bandage to keep the broken parts together, than any other within the range of surgery.

The shoulder-blade (*scapula*) lies on the upper and back part of the chest, forming the shoulder, where the upper arm bone inserts itself in a deep (*glenoid*) cavity, surrounded by a strong ligament (*capsular*); the shoulder blade itself is a thin, flat, and triangular bone, covering the back, and remains in its position, or moves in different directions, by numerous muscles which adhere to its projections.

The *upper arm* (*humerus*) forms a connection at the elbow with the *ulna* of the forearm, and at the shoulder its round head sinks into the *glenoid* cavity of the



scapula, in which it is firmly confined by the capsular ligament, thus forming the most movable joint in the whole system.

The *lower or fore arm* is composed of two bones, called *ulna* and *radius*; the former articulates with the humerus at the elbow, forming a perfect hinge-joint; this bone is located on the inner side of the fore arm, while the latter, the *radius*, lies on the outside of the fore arm (on the side where the thumb is placed), and articulates with the bones of the wrist, and forms the wrist-joint; the *ulna* and *radius*, at their extremities, articulate with each other, so that the hand can rotate, permitting its complicated and varied movements.

The *wrist (carpus)* consists of eight bones, ranged in two rows, and firmly bound together, thus allowing only a small amount of movement.

The *palm of the hand (metacarpus)* contains five bones, four of which are joined with the first range of the finger-bones, and the other with the first bone of the thumb.

The *fingers (phalanges)* have three ranges of bones, while the *thumb* has but two.

N. B. The mechanism of the hand and its wonderful adaptation to all the various purposes of life, is one of the many facts for man's superiority over the rest of the animal creation, and must excite in us the deepest interest and astonishment at the Divine wisdom and power.

The *lower extremities* comprise sixty bones, analagous in construction and form to the upper extremities; the *thigh-bone (femur)*; the cap of the *knee (patella)*; the *shin-bone (tibia)*; the *small bone of the leg* on the outside of the *tibia (fibula)*; the *instep (tarsus)*; the middle of the foot (*metatarsus*); and the *toes (phalanges)*.

The *thigh-bone* is the longest and *strongest* bone in the system, supporting the weight of the whole body. Its

upper part, a large round head, fills a corresponding cavity in the pelvis and forms the so-called *hip-joint*, thus admirably fitted, in its mechanism, for its various offices requiring strength and elasticity.

The *cap of the knee* (*patella*) is a small bone in front of the knee, connected with the thigh-bone by a strong ligament; it acts like a pulley, in the extension of the limb.

The *shin-bone* (*tibia*) is the largest bone of the lower part of the leg, of a triangular shape, and considerably enlarged at each extremity.

The *small-bone* of the leg (*fibula*) is of a similar shape with the former, but smaller, and bound firmly to it at each extremity. It lies on the outside of the leg.

The *instep* (*tarsus*) is formed of seven irregularly-shaped bones, so firmly bound together by ligaments as to allow of but little movement.

The *palm of the foot* (*metatarsus*) consists of five bones, corresponding to the five toes, with which they articulate on the front extremity, while on the hinder one, they articulate with one range of the bones of the instep.

By this arrangement, an arch is formed, convex above and concave below, which gives elasticity to the step, preventing the jarring of the whole frame by any weight thrown upon the feet, in their various uses.

The *toes* (*phalanges*) are composed of fourteen bones; each of the small toes has three, while the great toe has but two ranges of bones.

The *joints*, which have such important functions to perform, are composed of the extremities of two or more bones, lined and surrounded with *cartilages*, *synovial membranes*, and *ligaments*.

The *cartilage* (gristle) is a smooth, solid, elastic substance, of a pearly whiteness. It appears as a thin layer, on the articular surfaces of the bones—thicker in the cen-

ter upon convex surfaces, while the opposite arrangement exists upon concave surfaces.

The *synovial membrane* forms in a thin layer over the cartilages a closed sack, in the interior of which a constant secretion of a viscous fluid takes place, to facilitate the movements of the joints.

The *ligaments* consist of numerous straight fibres, arranged into short bands of various breadths, or so interwoven as to form a broad layer, which surrounds the joint, forming a capsular ligament. They are white, glistening, and not elastic, and found mostly exterior to the synovial membrane.

#### PHYSIOLOGY OF THE BONES.

The bones, by their solidity and form, retain every part of the system in its proper shape. They not only afford points of attachment for the muscles and ligaments, to hold the body together and bring it into motion, but they protect, in strong, bony cavities, all the important organs from external injuries, such as the brain, the eyes, etc.

The bones grow and decay, like any other tissue, by new particles being deposited from out the blood, and old, useless matter removed, by the constant action of the absorbing vessels. This has been tested thoroughly. Some of the inferior animals were fed with food containing madder. After a few days, several of these animals were killed, and their bones exhibited an unusual reddish appearance. The rest of the animals were for a few weeks fed on ordinary food, without containing madder, and when killed their bones had the natural color again.

The extremities of the bones are mostly the best suited to form joints; for which purpose their texture is more porous, consequently more elastic, and the external surface of the parts joining immediately each other in a joint is

covered with a cushion of cartilage, to diminish jarring and grinding, and to add to elasticity more beauty and grace. The contemplation of each minute particle, its uses and application in the system, its appropriate location, etc., the thousand wonderful machines which can be observed, acting usefully and quietly in our systems, reveal more and more the wisdom and omnipotence of the great Architect. Wonder and adoration fills the heart of the beholder of such creations as the human body, in its detail and in its whole, as a complete and noble machine, to run the errands of the holier part of man—that of the soul.

The bones serve in the animal frame as a proper basis; the ligaments hold the bones in their proper places firmly, as if clasped by steel, yet with room to move. They surround the joints like a hood, or lay from one bone to the other, in the form of side layers. By these ligaments, the lower jaw is bound to the temporal bones, and the head to the neck. They extend the whole length of the spinal column, in strong bands, on the outer surface, as well as within the spinal canal, and from one spinous process to another.

The joints are different as regards movement and complexity of structure. Some permit motions in all directions, as the shoulder; some in two directions, as the elbow, allowing flexion and extension; others in three and more directions; others, again, have no movement at all, as the bones of the skull, in adults.

#### THE MUSCLES.

A muscle consists of bundles of fibres of different size, each one inclosed in a cellular membranous sheath. Every bundle is composed of small fibres, and each fibre of numerous filaments, each of which is enclosed in a delicate sheath. At the end of the muscle, the fibrous substance

ceases, and the cellular structure predominates, modifying itself into what is called a *tendon* (cord) by which the muscle is adhered to the bone. This union is so firm, that the bone will sooner break than permit the tendon to separate from its attachment.

The form of the muscles varies very much, according to their uses. It is either longitudinal, having a tendon at each extremity, or it diverges in the form of a fan, called a radiate muscle. Others are penniform, converging, like the plumes of a pen, to one side of a tendon; or bi-penniform, where they thus converge on both sides of a tendon.

The color of a muscle is red; each fibre is supplied with arteries, veins, lymphatics, and nerves, both of sensation and motion.

Where a great many muscles congregate round an organ, they are inclosed in a *fascia*—a tendinous membrane, which assists the muscles in their action, by keeping up a tonic pressure on their surface. Besides, it protects the parts underneath; as for instance, in the palms of the hands or feet. The separate muscles are also inclosed in fascias, and arranged in layers, as their use requires it. The interstices between the different muscles are filled with fat, which gives to the body roundness and beauty.

We may arrange the muscles into four parts—those,

1. Of the *head* and *neck*.
2. Of the *trunk*.
3. Of the *upper extremities*.
4. Of the *lower extremities*.

As it is impossible to give an interesting detail of the muscles of the body, their insertions and names, without illustrating them by plates, we content ourselves with the following general remarks.

There are more than five hundred muscles in the human body, forming around the skeleton two layers, a superficial

and deep-seated one. Some muscles are voluntary, under the control of the will, such as those on the fingers, limbs, etc.; others are involuntary, as the heart. On the back, the muscles are arranged in six layers, one above the other, for the complicated movements which the back, neck, the upper extremities, and the abdomen require.

The *diaphragm* is a muscle which needs particular explanation. It is located between the chest and abdomen, separating them completely; being penetrated only by the pipe leading to the stomach, and by the great blood-vessels, leading from and to the heart. It may be compared to an inverted basin, its bottom being turned upward into the chest, its edges corresponding to the outline of the edges of the lower ribs and breast-bone. Thus, the cavity of the abdomen is enlarged at the expense of that of the chest. Respiration, however, changes the natural position of the diaphragm constantly, facilitating this process by its own action. While inhaling, the diaphragm descends into the abdomen, enlarging the chest; while exhaling, the reverse takes place: thus a constant and healthful action is given to both the respiratory and digestive organs. In this place, we should mention the great benefit which results from inhaling deep, to the full extent of the lungs, which is greatly facilitated by exercising the abdominal muscles, causing them to swell out when inhaling, and drawing them back when in the act of exhaling. A little exercise in this way will soon show the good results in the ease, and full extent with which the lungs can be filled.

The action of the muscles, in performing their various functions, can be explained only by the supposition of an inherent contractility, as a peculiar characteristic of the muscular fibre, which can shorten its substance when a sufficient stimulus acts on it. This stimulus is the nervous fluid, directed to the muscle by the will, if voluntary ac-



tions shall be performed, or it is supplied by the necessary demands of the organism, when the involuntary actions are wanted, such as the beating of the heart, etc. The nature of this nervous influence must be of a galvanic or electrical kind ; as these agents, when acting on the muscles, produce similar effects.

The rapidity with which the muscles move is truly astonishing, as any one can observe, in rapid speaking, or playing upon instruments. It is not always the size of the muscle which constitutes its strength ; but, also, the size of the nerves which lead to it, and the size and activity of the brain, which must supply the nervous fluid necessary as a stimulus to action. Yet a great deal depends on training and exercise, as these enlarge both the muscle and nerve. The blacksmith, wielding the hammer daily and forcibly, will have a stronger arm than the student, who merely exercises his muscles with a pen. From this, it is evident that the muscles ought to be educated or trained for the vigorous and healthful performance of their duty. Strict and systematic rules are given for this purpose, in an art called the gymnastic, which, to understand and practice diligently, we recommend strongly. It is indispensable for the young, and not without advantage for the middle-aged, to spend part of a day in systematic, healthful exercise of their muscles. The benefits are too great and palpable to bring them forward here, singly. Washing in cold water, also, invigorates muscular action.

#### THE TEETH.

The teeth differ from other bones in composition, nutrition, and growth ; while bones, when fractured, generally unite, the teeth never do, when broken. They are divided into two parts, *crown* and *root* ; the former, protruding from the jaw-bone, is covered by the highly polished enamel,

the root is placed in the sockets of the jaw, consisting of bony matter, through which several small vessels and nerves pass, giving nutrition and vitality. The teeth, when beginning to grow, are formed within dental capsules, inclosed within the substance of the bone, appearing as a fleshy, bud or granule, from the surface of which exudes the ivory on the bony part of the tooth. In growing, it rises in the socket, which is formed simultaneously around it, passing, finally, through the fleshy part of the gum. The first teeth, called milk-teeth, are twenty in number; they are temporary, and disappear from the seventh year upward; the last, or wisdom-teeth, do not appear until a person is twenty years of age.

The four front teeth are called *incisors*; the next one on each side is the *eye-tooth* (cuspid); the next two are *bicus-pids* (small grinders); and the next two are *grinders* (molars); and the last one is called the *wisdom-tooth*. The incisors, cuspids, and bicuspid have each but one root. The molars of the upper jaw have three roots, while those of the lower jaw have but two.

#### DIGESTIVE ORGANS.

These comprise the mouth and teeth, salivary glands, pharynx, œsophagus (stomach-pipe), stomach, intestines, lacteals (milk, or chyle-vessels), thoracic duct, liver and pancreas.

The *mouth* is, strictly speaking, no organ of digestion, but it contains the instruments of mastication and the organs of taste. The preparation of the food for digestion commences already in the mouth, where it is masticated and mixed with saliva from the *salivary glands*, of which there are six, three on each side of the jaw. Their names and positions are :

The *parotid gland* lies in front of the external ear, and

behind the angle of the jaw ; its duct opens into the mouth, opposite the second molar tooth of the upper jaw ; this gland is swollen and inflamed when children have the *mumps*.

The *submaxillary gland* lies within the lower jaw, anterior to its angle. Its duct opens into the mouth by the side of the string or bridle of the tongue (*frænum linguae*), on each side of which lies

The *sublingual gland*, of an elongated and flattened form ; it empties its saliva through seven or eight small ducts into the mouth by the side of the string of the tongue ; a disease called the “frog” consists in the swelling of this gland.

The *pharynx* or *throat* is the continuation of the mouth, and forms that cavity immediately below the palate where four passages unite, two coming from the nose and mouth, the other two going to the stomach and lungs.

The *stomach-pipe* (*œsophagus*) is a large membranous tube, descending behind the wind-pipe, the heart and lungs, through the diaphragm into the stomach ; it has two membranes, an internal, or mucous, and an external, or muscular coat ; the latter consists of two sets of fibres, one circular, the other longitudinal.

The *stomach* lies on the left side of the abdomen, immediately below the diaphragm, where the stomach-pipe enters its upper part, called the *cardia* ; from this point it enlarges into a sack, which lies with its larger extent on the left side, while its smaller end contracts gradually toward the pit of the stomach, where it empties into the intestines ; this opening in the stomach is called the *pylorus*. The stomach has, beside the mucous and muscular coats, an outer or serous coat, which is very tough and strong, of a smooth and highly polished appearance, and confines the stomach in its proper location. The stomach has a great number of glands, to secrete the stomach juice.

The *intestines*, or alimentary canal, are divided into two parts, the *small* and *large*. The former, commencing at the stomach, measures about twenty feet, while the large intestine, ending at the rectum, is about five feet in length. The latter is divided into three parts, the *cæcum* (the beginning of the large intestine), which lies near the right upper hip-bone; the *colon*, which, from this point on the right side, ascends to the region below the liver, then transversely crosses the upper part of the abdomen, from below the liver, to the lower part of the stomach on the left side, where it bends again, descending to the left hip-bone, and entering the cavity of the pelvis, being called here the *rectum*.

The *small* intestine, commencing at the stomach and ending at the *cæcum*, is also divided into three parts; the *duodenum*, called so from being generally twelve fingers long, commences at the stomach, ascends obliquely to the under surface of the liver, then descends perpendicularly (where it receives the ducts from the liver and pancreas), and passes transversely across the lower portion of the spinal column behind the colon, terminating in the *jejunum*, its continuation; and this, in its turn, is continued by the *ileum*, which empties itself into the *cæcum* at the right hip, after the small intestines have made numerous windings. While the stomach and intestines receive the food for digestion, and assimilate it, various other secretions from the liver, pancreas, etc., enter the alimentary canal to foster this process, and still other vessels are ready to absorb the assimilated juice and carry it into the circulation of the blood. Of the latter are the *lacteals*\* or minute vessels, which commence

---

\* These are the class of absorbing vessels, which perform, during an *Asiatic cholera* attack, such fearful and often fatal offices; their action becoming inverted, permits the lacteal fluid, or the real nourishing juice, to escape in the intestines, drawing after it all what is contained in the

on the mucous membrane of the small intestines; from these they pass between the membranes of the *mesentery* to small glands, out of a collection of which larger vessels run to another range of glands, and so on, through several ranges of these glandular bodies, until they all empty themselves into one large duct, called the *thoracic* duct, which ascends from the abdomen upward, lying in front of the spinal column, and passes through the diaphragm to the lower part of the neck; here it makes a sudden turn downward and forward, emptying itself into a large vein, which passes into the heart; its diameter is equal to a goose-quill.

The *liver* is a gland, which lies under the short-ribs of the right side below the diaphragm, and is composed of several lobes; its upper surface is convex, its lower concave; it is the largest organ in the system, weighing about four pounds; its use is to secrete the bile, a fluid so necessary for chylication. The *gall-bladder*, containing the surplus of bile not immediately necessary for the system, being, as it were, a reservoir, lies on the under surface of the liver, and empties into the gall duct, which carries the bilious fluid to its destination in the duodenum.

The *pancreas* is a long, flattened gland, similar to the salivary glands on the neck; it is about six inches long, weighs three or four ounces, and lies transversely across the posterior wall of the abdomen, behind the stomach; its product, a saliva, or *pancreatic juice* (necessary for digestion), is carried by a duct into the intestines in the duodenum, just opposite where the gall duct opens into the intestines.

The *spleen* (milt), of an oblong, flattened form, lies in

---

thoracic duct, which, after the decomposition of the blood itself has taken place, receives the serum or white blood, and discharges it through the same absorbent vessels into the intestines.

the left side below and touching the diaphragm, stomach, and pancreas. It is a reservoir capable of containing a great quantity of blood, ready for the purposes of the liver, to secrete the bile. It has itself no outlet or duct, showing that the office it has to perform is more of an external nature, as like that, for instance, what the tender has to perform to the engine; for this purpose, large vessels communicate with the liver freely. Further we can not explain the uses of the spleen.

The *omentum* (caul) descending in four layers of serous membrane from the stomach and transverse colon, contain fatty matter, deposited around the vessels, which ramify through its structure. Its functions are: to protect the intestines from cold, and to facilitate their movements.

The whole digestive apparatus is supplied with arteries, veins, lymphatics, and nerves; the latter from the *ganglion* system. (See this.)

One of the most wonderful processes in the animal economy is that of the assimilation of food.

During its mastication (chewing) a considerable quantity of saliva (spittle) is mixed with it, the object of which is, to soften and moisten the food, preparing it for easy deglutition. When it reaches the stomach, it is subjected to the powerful action of the so-called stomach-juice (gastric juice), which is secreted by the glands immediately located in the substance of the stomach. Besides, it is constantly in motion by the action of the muscles of the stomach, which brings the food into contact with the mucous membrane, and thus becomes completely saturated with gastric juice, or dissolved into a pulpy, homogeneous mass of a creamy consistence, called *chyme*, which passes, as fast as it is made, through the pylorus into the duodenum. In the duodenum, the bile and pancreatic fluid mix with the chyme, which, by their action, is converted into a whitish



fluid called *chyle*, and separated from the rejected parts ; at this point, the formation of the fœcal matter commences.

From the above is seen that the bile has no agency in the change through which the food passes in the stomach. The common belief, that bile is in the stomach, is erroneous. If bile is ejected in vomiting, it merely shows not only that the action of the stomach is inverted, but also that of the duodenum. Emetics will, generally, bring bile from the most healthy stomach, by inverting the action of the stomach and duodenum ; the expression, being bilious, having a bilious attack, etc., is generally wrong, the bile having nothing to do with these disorders, for which a better expression would be : the stomach is out of order ; if persons, generally, would know this fact, a great deal of wrong and disastrous treatment would be avoided, as they would not force the stomach to eject bile continually, where there is none located.

#### THE URINARY SYSTEM.

This system, whose duty it is to secrete and carry out of the system the urine, is composed of the *kidneys*, the *ureters*, and the *bladder*, with the *urethra* attached to it.

The *kidneys*, between four and five inches long, and two and a half broad, lay in the lumbar region, behind the peritoneum, on each side of the vertebræ. Their texture is dense and fragile, presenting, in its interior, two structures, an external (cortical), and an internal (medullary) substance. The former contains the bloodvessels, which carry the blood from which the urine shall be separated ; the latter consists of tubes, called the *pyramids of Malpighi*, in which this process is carried on. The secreted urine is then received in the pelvis of the kidney, from which it runs into the *ureter*, the excretory duct of the kidney, a membranous tube of the size of a goose-quill, about

eighteen inches in length ; this runs down along the posterior wall of the abdomen, behind the peritoneum and crossed by the spermatic vessels, to the bladder in front, in which it empties from behind and on the side.

The *bladder* is of an oblong, ovoid shape, situated behind the os pubis and in front of the rectum. It is the reservoir for the urine, which is carried there by the two ureters from the kidneys. It is retained in its position by eleven ligaments, supporting it on all sides. The bladder is composed of three coats, an external, or serous, a muscular, and a mucous, or internal coat.

The *urethra* is the membranous canal issuing from the neck of the bladder. It is curved in its course, composed of two layers, a mucous and an elastic fibrous coat. Through it passes the urine out of the body.

#### THE RESPIRATORY AND CIRCULATORY ORGANS.

These organs are closely connected with each other. All the blood must pass through the lungs, to receive there new life and energy, by being exposed to the oxygen of the air. For this and other reasons, we will treat of these organs here in connection.

After the nutrient portion of the food (the *chyle*—see page 417) is discharged by the thoracic duct into the left subclavian vein, at the lower part of the neck, it is carried to the right cavity of the heart, where it mixes with a large quantity of venous blood. This mixture of fluid, as such, would not be suitable to restore the lost powers of the body, unless subjected to a process, by which the chyle is converted into blood, and the venous blood freed of its carbonic acid and water.

The *respiratory* organs consist of the *windpipe* (*trachea*), the *bronchia* (continuations of the *trachea*), and the *air-cells* (the extreme points of the bronchia.) The *lungs* are

composed of innumerable ramifications of the bronchial tubes, ending in air-cells, lymphatic vessels, and the pulmonary arteries and veins; their connections by cellular tissue, forms the *parenchyme* of the lungs, or its substance. The lungs are divided into two large parts—the left and right, each one of which is inclosed in a serous sack, called the *pleura*. Between the right and left lung, more to the left side, lies the heart, separated from either by a serous membrane around it.

The *trachea* proceeds from the larynx, descending to the pit of the neck, where it divides into two parts, assuming the name of

*Bronchia*; which descends, in numerous ramifications, into the lungs, and forms altogether a surface of twenty thousand square inches, or about thirty times the surface of the human body.

The *air-cells* are small bronchial tubes, and form the ends of these ramifications, retaining the air, when once inflated, except pressed out by force.

The trachea, bronchia, and air-cells are lined with the mucous membrane, and supplied with arteries, veins, and nerves.

The object of respiration is to free the system of carbon and hydrogen, which accumulate in the system, and would make an end to its existence, if not removed. For its removal, an allwise Providence used the inhaling of air; which contains oxygen, in sufficient quantity to form a combination with the carbon and hydrogen, which then is exhaled, in the form of carbonic acid and water. In this process, another wonderful provision was contained, which gives life and motion to the whole system. It is the generation of heat, which is the principal cause of the circulation of the blood. Before the birth of a child, the generation of heat in its body, and the exchange of rejected

particles takes place, through the medium of the placenta, adhering to the mother. But after the birth, that heat is generated in the lungs, and respiration and circulation of the blood are then united closely—the former being sole cause and motor of the latter. For the heat which is generated in the lungs (by a kind of combustion caused by a union of the oxygen of the inhaled air with the carbon and hydrogen of the blood) infuses itself into the blood at the finest air-cells, sufficiently to drive it through the heart and the arteries, until it reaches the capillary vessels where the venous system commences. Here, again, on the confines of the body, the higher temperature of the lungs, draws to these the cooler venous blood. In both cases, the heart, by its regular vibrations, acts as a balance-wheel; the left heart in giving renewed vigor and regularity to the rushing of the arterial blood into the whole system; the right heart acting in the same manner on the venous blood as it rushes into the lungs. The idea heretofore entertained by physiologists, as if the heart alone by the impulse of its contractions was able to propel the blood through the arteries, and have also an effect on the venous blood in returning to the heart, is beginning to be regarded as very doubtful and altogether improbable; in its stead, the above explanation of the motion of the blood by the heat generated in the lungs seems to be much more rational, and, after our opinion, the only true one which can be presented.\*

The *circulatory* organs are the *heart*, *arteries*, *veins*, and *capillaries*.

The *heart* placed obliquely, in the left cavity of the chest, has the form of an inverted cone, the base of which lies upward and backward in the direction of the right shoulder, and its apex points forward to the left side, about three inches, to a space between the fifth and sixth ribs;

---

\* See the work of Mrs. Willard on this interesting subject.

below, it rests on the tendinous part of the diaphragm. The sack, by which the heart is surrounded, is called *pericardium*, the interior surface of which secretes a watery fluid that lubricates the exterior of the heart. The dropsy of the heart has its seat in this place.

The heart itself is a muscle, whose fibres run in different directions, but mostly in a spiral direction. It is divided into two distinct compartments—the right and left *ventricle*. Before the birth, these ventricles are united by an opening, permitting the venous and arterial blood to mix at this place. After the birth, this opening closes and obliterates entirely, preventing the venous blood of the right ventricle from passing directly into the left. If this opening, by a fault of nature, remains unclosed, a disease ensues, called the *blue disease*, because the child looks bluish, on account of the venous blood of the right heart entering the circulation of the arterial blood of the left ventricle. All vessels leading from the heart are called *arteries*; those leading to the heart, *veins*. The right heart sends a *pulmonary artery*, filled with venous blood, to the lungs, and the left heart receives *pulmonary veins*, filled with arterial blood, from the lungs; in this case only, such an exception exists. Otherwise, all arteries in the body carry red or arterial blood, and all veins dark or venous blood.

The venous blood, together with the chyle, is propelled through the pulmonary arteries, by the right heart, into the lungs. Here it receives the oxygen of the air, is converted into red blood, and carried by the pulmonary veins into the left heart, which sends it through the great artery, called *aorta*, in innumerable divisions, into all the parts of the body. In this way, two systems of circulation are formed—one called the *lesser* or *pulmonic circulation*, where the blood rushes from the right heart, through the lungs, and returns, changed into red blood, to the left; the other,

called the *greater*, or *systemic circulation*, where the blood rushes through all parts, and to the very confines of the body; and returns, changed into dark blood, to the right heart. This double circulation be illustrated easily by the figure 8, the centre of which is the place where the heart is located.

The *arteries*, which convey the blood from the heart to every part of the system, are composed of three coats. The external or cellular coat, is firm and strong; the middle or fibrous, is elastic and fragile; the internal is a thin, serous membrane, which gives the artery inside a smooth polish. Communications between arteries are numerous.

The *veins*, which return the blood to the heart, are thinner, and more delicate in structure than the arteries. They are composed, like these, of the same three coats, but each one is more delicate and fine.

The *capillaries* form an extremely fine net-work of vessels, between the ends of the arterial and the beginning of the venous system. Through these, all the blood must pass; and in this point, the capillary system resembles the fine ramifications in the parenchyme of the lungs, where the blood passes through infinitely small vessels.

The *pulse* is caused by the pressure of the blood against the coats of the arteries. In infants, it beats in a minute more than a hundred times; in adults, about seventy times; in old persons, less than that—sometimes only sixty times.

The quantity of blood in a healthy individual varies very much; fat persons have generally less. On an average, a healthy adult has from twenty-five to thirty pounds.

The *lymphatics* are the vessels by which the process of *absorption* is carried on, to remove particles of useless or injurious matter from the system. They are extremely minute at their origin, and exist in great numbers in the skin, mucous membranes, and lungs. As they increase in



size, they diminish in numbers. At certain points, they pass through soft bodies, called *lymphatic glands*, which are mostly located in the groins, arm-pits, on the neck, in the chest, and the abdomen. Sometimes these glands swell producing lumps known as *kernels*, etc.

The *lacteals*, are vessels similar to the former, but designed to absorb the chyle from the small intestines, where they are located in great numbers. They transfer their contents to the thoracic duct, and this conveys the chyle to the subclavian vein, at the lower part of the neck.

### THE NERVOUS SYSTEM.

This system, by which life and sensation is imparted to the whole body, consists of two distinct parts—the *brain*, with its spinal appendix, together with all the nerves, leading from these great nervous centers to all parts of the body, and the *Ganglion* system of nerves, called, also, the sympathetic nerve, located on each side of the spinal column, but having its principal plexuses around the abdominal organs. The constant influence which these two great systems have on each other, constitutes and sustains, very likely, the life of the organism, in all its functions. We may consider them as acting, in this respect, as the poles of a galvanic battery. To explain their workings in detail, or the moving cause of their actions, must remain impossible; as no mortal eye can scrutinize the creative measures of the Almighty.

We give here a short anatomical synopsis of these most interesting parts of the body; requesting our readers, at the same time, to peruse larger works on this subject, if their time and inclination will permit.

The *brain* is located within the skull-bones (cranium), and is divided into two hemispheres—the right and left. This is the case in the upper part of the brain, below the

two hemispheres are connected by a dense layer of transverse fibres, called the *corpus callosum*. In the interior of the brain, there are several cavities, two of which are of considerable size, called the lateral ventricles. These are the spots where the water collects, in that disease, called "Dropsy of the Brain."

The substance of the brain is of a pulpy character, quite soft in infancy and childhood; it gradually becomes more consistent in advanced years.

It is divided into three parts—the large brain (*cerebrum*), the small brain (*cerebellum*), and the medulla oblongata, or that part of the spinal cord which lies within the skull.

These three parts of the brain within the skull are invested and protected by the three membranes of the brain: the *dura mater*, a firm and fibrous membrane, next to the skull; the *arachnoid*, a serous membrane, which envelops the brain and spine, and is like the serous membrane of the heart and lungs—a closed sack; this is the middle membrane of the three; the *pia mater*, a vascular membrane, composed of innumerable vessels, held together by cellular tissue; this is the nutrient membrane of the brain.

From the brain issue the *cranial nerves*, in twelve pairs, to supply the organs of sense and motion with life and vitality. They are arranged, as regards local origin as follows: 1. The olfactory; 2. The optic; 3. The *motores oculorum*; 4. The patheticus; 5. The tri-facial; 6. The abducentes; 7. The portio dura; 8. The facial; 9. The glosso-pharyngeal; 10. The pneumogastric; 11. The spinal accessory; 12. The hypo-glossal.

The *spinal cord*, its membranes, and the roots of the *spinal nerves*, are contained within the spinal column. The cord is the continuation of the brain, or its lowest part, the medulla oblongata, and is divided in two lateral cords

by an anterior and posterior fissure. Each of the lateral cords is divided by furrows into three distinct sets of fibres, called the anterior, lateral and posterior columns; the posterior are the columns of sensation; the anterior, those of motion, and the lateral are divided into their functions between motion and sensation.

From these divisions issue the *spinal nerves*, thirty-one pairs in number, each one from two roots, an anterior and posterior, combining the functions of motion and sensation. After the two roots have united, which constitute the spinal nerve, it passes through the opening between the vertebræ on the sides of the spinal column, and distributes itself to its respective organs in innumerable ramifications.

The spinal nerves are divided from the head downward, into eight cervical pairs, twelve dorsal, five lumbar, and six sacral pairs.

The *ganglion system*, or *sympathetic nerve*, consists of a series of knots or ganglia, lying on each side of the spinal column, united with each other by nervous threads. It connects itself with the brain and spinal nerves. Each ganglion or knot may be considered as a distinct center, from which branches issue in four directions, some for communication with other ganglia, some for distribution. The latter accompany the arteries to the different organs, and form a net-work around them, called plexuses, and named after the artery which they accompany; for instance, the mesenteric plexus, hepatic plexus, etc. In this way all the internal organs of the head and trunk receive branches from the sympathetic nerve, which, therefore, is considered a nerve of organic life.

#### THE SKIN.

The skin covers the exterior of the body, and appears, to the naked eye, as composed of one membrane only. By

closer examination, however, it is found to consist of two distinct layers of membrane, the upper or scarf-skin (epidermis), and the real skin (cutis vera), which lies underneath the former.

The upper-skin (epidermis) is thin, semi-transparent, like a fine shaving of horn; it has no nerves or bloodvessels, and is without sensation. The hair and the nails are of the same nature.

The real skin (cutis vera or corion) consists of minute fibres, interwoven with each other to a firm, strong and flexible web. This manner of composition permits innumerable interstices or pores, which are finer or closer on the superficial part of the skin than on its lower surface. The upper surface of the skin contains blood-vessels and nerves, looks, therefore, more red or pinkish, and is sensitive; while the lower strata of the skin connects with the fibrous web, in which the sub-cutaneous fat of the body is deposited.

It is in the skin, principally, that the finest ramifications of the arteries, and the commencement of the finest veins, with the intermediate microscopic capillary vessels are found. One of the greatest changes in the economy of the system takes place in the skin — the change of the arterial into venous blood; just the opposite from what occurs in the lungs, where the blood changes from venous to arterial. In both instances the blood is carried through infinitely fine and small vessels, where it loses its former identity, even all color, resuming a different one, as it reappears and is collected in different vessels.

Besides, the skin is supplied with nerves in all directions, wherever there are bloodvessels; also lymphatics in great numbers. The oil-glands are contained in some parts of the true skin, in great abundance, as on the face and nose, the head, etc.; they communicate with the surface of the

skin through small tubes, which penetrate the skin, some in spiral windings, others in a straight line. These glands deserve our particular notice, as their usefulness is so great, that neglect in cultivating their healthy action is felt very severely. They appear in great distinction on the eye-lids, where their disease causes the so-called styes; they produce in the ear-passages the so-called earwax: they impart oil to the hair by infusing it into the sheath of each hair.

In some persons these oil-glands can not discharge their contents on the surface of the skin, on account of some diseased state, when it becomes dry and dense, appearing in round, dark spots, particularly on the forehead and nose; this disease is popularly called a "worm or maggot."

The *perspiratory glands* are oval shaped or globular balls, laying in the deeper meshes of the real skin, from whence they communicate with the surface through minute cylindrical tubes, called the "pores." The quantity of these pores is truly astonishing; they average about twenty-eight hundred in the square inch of surface; and if an ordinary sized man has twenty-five hundred square inches of surface, the whole number of pores in the skin amounts to seven million. How important must be, therefore, the culture of the skin, as an organ for secretion and exhalation.

## II. HYGIENE AND HYDROPATHY.

### HYGIENE.

*Hygiene* comprises the knowledge of instituting such a mode of living as is best calculated to preserve health. The clearer this knowledge is, and the closer we follow the rules of health, the surer the prevention of diseases will be. It is certain, beyond doubt, that less misery, disease, and death would mar the happiness of this fair world, if its inhabitants would live more in accordance with the laws of nature, and less after their own often mistaken notions and passions.

Reform, in this respect, has already commenced; and, with a more rational method of healing, and the application of a more efficient and salutary tonic agent (the water), invalid and enfeebled mankind can look forward to a happier, because a healthier future.

In the following pages, we will notice briefly the most important subjects affecting the physical well-being of an individual. These are: *food and drink, sleep, air and exercise, clothing, occupation.*

In all our wishes and necessities of physical and mental things, we must never forget two important principles: first, whether they are really beneficial or useful, and if so, that we use them in moderation.



## 1. FOOD AND DRINK.

The waste of bodily strength must be supplied by suitable food and drink. This is the greatest necessity of nature; and if errors are committed in this respect, our injured system must suffer the bad consequences. It is, therefore, of the utmost importance to pay particular regard to the selection of our daily food and drink, in order to escape disease, and often premature death.

Suitable food must have two paramount qualities: it must be *nutritious* and *digestible*. Not all nutritious food is digestible, and *vice versa*; for instance, fat meat is more nutritious than lean meat, yet it is very indigestible, and requires the strongest stomach, and unimpaired action of the bile, to be digested at all. A great many experiments have been made, to test the relative digestibility of food, and from these, and the experiments on the sick bed, valuable knowledge in regard to the choice of food has been collected. According to these investigations, boiled rice is the most digestible, and roasted pork, fat and lean, the most indigestible substance.

In order to give the reader a more comprehensive view of the relative digestibility of food, we insert Dr. Beaumont's results, as related in Dr. Tarbell's "Sources of Health."

			Hours	Min
Rice.....	Boiled.....	Digested in.....	1	
Pig's Feet, soused.....	" .....		1	
Tripe, soused.....	" .....		1	
Eggs, whipped.....	Raw .....		1	30
Trout, Salmon, fresh.....	Boiled.....		1	30
" " " .....	Fried .....		1	30
Apples, sweet.....	Raw .....		1	30
Venison steak.....	Broiled.....		1	35
Brains .....	Boiled.....		1	45

		<i>H.</i>	<i>M.</i>
Sago .....	Boiled..... Digested in.....	1	45
Tapioca.....	“ .....	2	
Barley .....	“ .....	2	
Milk.....	“ .....	2	
Liver, beef's, fresh.....	Broiled.....	2	
Eggs, fresh.....	Raw .....	2	
Codfish, cured, dry.....	Boiled.....	2	
Apples, sour.....	Raw .....	2	
Cabbage, with vinegar.....	“ .....	2	
Milk.....	“ .....	2	15
Eggs, fresh.....	Roasted .....	2	15
Turkey, wild.....	“ .....	2	18
Turkey, domestic.....	Boiled.....	2	25
“ “ .....	Roasted.....	2	30
Gelatine.....	Boiled.....	2	30
Goose.....	Roasted.....	2	30
Pig, sucking.....	“ .....	2	30
Lamb, fresh.....	Broiled.....	2	30
Beans, pod.....	Boiled.....	2	30
Cake, sponge.....	Baked.....	2	30
Parsnips.....	Boiled.....	2	30
Potatoes, Irish.....	Roasted.....	2	30
“ “ .....	Baked.....	2	30
Cabbage, head.....	Raw.....	2	30
Spinal Marrow.....	Boiled.....	2	40
Chicken.....	Fricassee.....	2	45
Custard.....	Baked.....	2	45
Beef.....	Boiled.....	2	45
Oysters.....	Raw.....	2	55
Eggs.....	Soft Boiled.....	3	
Beef, fresh, lean, rare.....	Roasted.....	3	
Beef-steak.....	Broiled.....	3	
Pork, salted.....	Raw.....	3	
“ “ .....	Stewed.....	3	
Mutton, fresh.....	Broiled.....	3	
“ “ .....	Boiled.....	3	
Chicken Soup.....	“ .....	3	
Dumpling, apple.....	“ .....	3	
Oysters.....	Roasted.....	3	15
Pork-steak .....	Broiled.....	3	15

		<i>H.</i>	<i>M.</i>
Pork, salted.....	Broiled.....	3	15
Mutton.....	Roasted.....	3	15
Bread, corn.....	Baked.....	3	15
Carrot.....	Boiled.....	3	15
Sausage.....	Broiled.....	3	20
Flounder.....	Fried.....	3	30
Oysters.....	Stewed.....	3	30
Beef.....	Boiled.....	3	30
Butter.....	Melted.....	3	30
Cheese, old.....	Raw.....	3	30
Bread, wheaten, fresh.....	Baked.....	3	30
Turnips.....	Boiled.....	3	30
Potatoes, Irish.....	“.....	3	30
Eggs.....	Hard boiled.....	3	30
“.....	Fried.....	3	30
Green Corn and Beans.....	Boiled.....	3	45
Beets.....	“.....	3	45
Salmon, salted.....	“.....	4	
Beef.....	Fried.....	4	
Veal.....	Broiled.....	4	
Fowls, domestic.....	Boiled.....	4	
“ “.....	Roasted.....	4	
Ducks, domestic.....	“.....	4	
Heart, animal.....	Fried.....	4	
Beef, old, salted.....	Boiled.....	4	15
Pork, salted.....	Fried.....	4	15
“ “.....	Boiled.....	4	30
Veal.....	Fried.....	4	30
Ducks, wild.....	Roasted.....	4	30
Suet mutton.....	Boiled.....	4	30
Cabbage, with vinegar.....	“.....	4	30
Suet, beef.....	“.....	5	3
Pork, fat and lean.....	Roasted.....	5	15

The above table is very explicit as regards the digestibility of certain substances, and we can ask its particulars with confidence, whenever we wish to decide this point in the choice of food ; but, in relation to its nutritious qualities, whether highly nutritious or less nutritious food is

necessary, we must follow the researches of Prof. Liebig, who first explained satisfactorily the connection of digestion and respiration. He showed that the lungs receive a greater quantity of oxygen in cold than in warm climates and seasons. A corresponding amount of carbon, which is contained in the food, is necessary, to mix with the blood, and unite in the lungs with this oxygen, to produce such a degree of animal heat, as is required by the external circumstances of the individual. The colder the climate, therefore, the more carbon (or food containing carbon) is necessary, to produce animal heat; the warmer the season or climate, the less of it is necessary. For this reason, is fish-oil the staff of life for the Greenlander; because it contains the greatest amount of carbon or nourishment, for producing a great quantity of animal heat, which he needs in those regions; while persons under the equator live principally upon watery, vegetable productions, affording hardly any nutriment; having no need of a great deal of animal heat; as nature supplies this by the hot climate itself. Animal food, in general, is more nutritious than vegetable; but the structure of our digestive organs, from the teeth to the lowest part of the intestines, shows conclusively that man was destined to partake of both kinds, animal as well as vegetable. The latter must preponderate, however, on account of the reasons above given, in warm seasons and climates, while the animal food takes precedence in cold regions and seasons: in the highest latitudes, even to the exclusion of all vegetable diet. In temperate regions, a judicious combination of both kinds of food is necessary to sustain life harmoniously.

The digestibility of food varies according to individual strength and habit. Pork may be completely indigestible to the invalid and student, although it is highly nutritious,

while it is perfectly digestible for the wood-cutter and hunter in wild regions.

*Animal food* contains, principally, *febrine*, *albumen*, *gelatine*, *oil* and *caseine*, and *osmazome*.

*Febrine* is digestible and nutritious.

*Albumen* is nutritious, but only digestible in its natural or not coagulated state, as the white of eggs, which becomes indigestible when hardened too much by heat.

*Gelatine*, the substance which forms the so-called jelly, is nutritious and readily digested ; but its digestibility is frequently impeded by improper ingredients mixed with it, particularly in the form of acids.

*Oil* is nutritious, but indigestible ; it requires the action of bile in the intestines to assimilate it properly.

*Caseine* is less nutritious and digestible than any of the former substances. It is contained mostly in the cheesy part of the milk.

*Osmazome* is that substance which imparts flavor to the meat and the soups. Its presence shows the nutritious and digestible quality of the meat.

According to the predominance of one of the above named substances, animal food has been divided into three classes :

1. *Fibrine* class ; comprising mutton, beef, pork, ducks, geese, and venison.

2. *Gelatinous* class ; to which belong veal, lamb, young poultry, and most kinds of fish.

3. *Albuminous* class ; containing oysters, eggs, brain and liver, the sweet-bread in calves.

The digestibility of meat varies according to the mixture of oil, or osmazome ; for instance, ducks and geese, although belonging to the fibrine or best digestible class, are more indigestible than beef ; because they are of an oleaginous nature.

*Mutton* and *beef* are the most nutritious and digestible of all meats. The former is rather better for invalids—dyspeptics in particular. *Venison*, also, is very nutritious and digestible.

*Pork* is indigestible, although very nutritious; because it is united to so much oily or fat substance. Besides, its continual use causes diseases of the skin and lymphatic system.

*Veal* and *lamb* belong to the gelatinous class; very nutritious, but not so easily digested, because they are not sufficiently matured or ripened, as it were.

*Poultry* differs, also, according to its age; the young containing more gelatine, and requiring, in consequence, not so much roasting or baking; the older ones containing more fibrine, and requiring, therefore, more cooking and roasting.

Among the fowls, there are none more suitable for invalids than the wild fowls, such as partridges, pheasants, etc.

*Eggs* and *oysters* are very nutritious, and very digestible, in their natural state; but to harden their albumen, by cooking or stewing too much, renders them very objectionable for the sick. Clams are not so digestible.

*Fish*, in general, differ as regards their digestibility, in the same proportion as they contain more oily substance or not. Marine fish generally contain more oil than fresh water fish. Fish without scales contain more oil, and are, therefore more indigestible, than other fish.

*Vegetable food* contains, principally, *starch*, (*fecula*) *gluten*, *mucilage*, *oil*, and *sugar*.

*Starch* is a very nutritious substance, and its predominant quantity in the grains, potatoes and other roots, renders them desirable aliments; it abounds in arrow-root, sago, tapioca, farina.

*Gluten* is that substance in wheat, which enables it to



form a tenacious paste by the mixture of flour with water ; it is similar to the animal fibrine.

*Mucilage*, or gum, is a very nutritious substance, as, for instance, gum-arabic, less digestible, however, than starch or sugar.

*Sugar* is one of the most nutritive and digestible substances in nature ; it forms, with the starch, the great bulk of vegetable consumption.

*Oil*, in the vegetable, is as indigestible but nutritious a substance as that of the animal, and must be used in not too large a quantity.

*Rice* is the most digestible and nutritious vegetable wholesome, for healthy and debilitated persons.

The *potato* ranks next, if roasted or baked, while it is less digestible when boiled ; the sweet-potato is more nutritious than the common potato, but less digestible.

*Arrow-root*, *sago*, *tapioca*, *farina*, are both digestible and nutritious.

*Beets* contain a good deal of sugar, and are, on that account, very nutritious ; but, as the fibrous part of the beet is entirely indigestible, invalids must abstain from its use.

Onion, cabbage, asparagus, and tomato contain but very little nutriment, and must not be indulged in by persons having, or being disposed to the diseases of the abdomen.

Fruits in general are refreshing and wholesome, but not very nutritious. This latter quality makes them very desirable in diseases, where a nutritious diet otherwise would be detrimental. In the convalescence from fevers, apples, baked or roasted, dried prunes stewed, etc., form a refreshing and grateful diet for the recovering patient ; in the green state, fruit produces flatulency, and must be avoided by patients, although a moderate use of ripe fruit, for a

well person, is very good, except in seasons where bowel-complaints predominate.

In order to enable the reader to distinguish between the vegetables as regards their nutritious quality, we subjoin here a list of some of those most in use, as related in Dr. Tarbell's Sources of Health :

100 lbs.	Lentils	- -	contain	- -	94	parts of nourishment.
" "	French Beans	- - - -			92	" "
" "	Rice	- - - -			90	" "
" "	Kidney Beans	- - - -			89	" "
" "	Wheat	- - - -			85	" "
" "	Barley	- - - -			83	" "
" "	Good Bread	- - - -			80	" "
" "	Rye	- - - -			80	" "
" "	Meat (average)	- - - -			35	" "
" "	Potatoes	- - - -			25	" "
" "	Carrots	- - - -			14	" "
" "	Beets	- - - -			14	" "
" "	Turnips	- - - -			8	" "
" "	Cabbage	- - - -			7	" "
" "	Greens	- - - -			6	" "

*Water* is the only liquid which nature has furnished in abundance for drink ; without it, all living beings and organized forms would perish. How important it is, therefore, to procure an article, so indispensable and necessary, in as pure a state as possible. The best water for drink is, without doubt, rain water, as it is free from admixtures of earthy substances ; yet, even rain water frequently becomes impure by falling through an impure atmosphere, particularly that hanging over large towns, or by running on roofs of houses, through unclean spouts before it reaches the cistern.

Equal to rain water in purity and taste is the water procured by the melting of snow and ice, if the latter has been formed in clean ponds or rivers.

But where rain water can not be obtained in sufficient quantity or purity, it is generally taken from the rivers and lakes, which furnish a better water than wells. The latter contains more or less lime, salts, or other earthy impurities, which render it neither palatable nor healthy.

Water, coming from marshy regions, is entirely unfit for drinking; at least, not until, by boiling, the inherent organic matter is destroyed.

A healthy person ought to habituate himself to a free use of cold water. Besides using it externally in baths, he should drink morning, noon, and evening, with his meals, a sufficient quantity to dilute the food. But the water he drinks must be cold, as it is this latter quality which offers to the stomach a natural and strengthening stimulus for digestion; in summer, and in southern latitudes, ice is necessary to render the water fit for hygienic purposes. If a person is once used to drink plenty of cold or ice water at or after his meals, he does seldom suffer from indigestion, and needs not the use of stimulating drinks; and every one can easily habituate himself to this practice. It is a wrong idea, which seems yet to prevail to a great extent, that it is unwholesome to drink water at the meals; but experience has the advantage, and one can soon test the matter for himself. The trial must only be continued for some time, in order to allow the system to accommodate itself to the new habit. Health and good spirits will be the consequence of such a change in a person's living.

It is also very proper to drink a tumbler of fresh water when retiring in the evening, or on rising in the morning. In doing this, the want of other drinks or beverages of a stimulating nature, such as coffee, or tea, is not felt so much, and can be dispensed with easier by degrees.

*Coffee*, especially, is one of those enticing palatable beverages which has laid the majority of the civilized world

under contribution ; it is, to a certain extent, nutritious, and made more so by the addition of cream and sugar ; but the greatest part of its contents is medicinal, and, consequently, not fit for daily consumption, without creating artificial alterations in a person's healthy condition. Hahnemann, the illustrious founder of Homœopathy, was the first who drew the attention of the coffee-drinking community to the destructive tendency their favored article had on the constitution. He distinguished well between a moderate use of this drink and an immoderate consumption in large quantities and strong decoctions. We recommend our readers to read his admirable article on this subject. It exhausts all what can be said in regard to it, and is capable to cure the most inveterate coffee-drinker. Homœopathy forbids its use in most all diseases, and insists on a more natural diet ; in this important undertaking, she is supported by the rigorous system of hydropathy, which demands the abolition of all stimulating drink, and allows nothing but water.

But, if coffee can not be abstained from, or if a person is only using it moderately, without experiencing bad consequences, it would be well to recommend a good preparation of the bean, by roasting it but slightly, grinding it never long before it is used, and adding to the decoction one-half of boiled milk. In this way, coffee may be drank with comparative impunity.

*Tea*, as a daily drink, meets with the same objections as coffee, and is only tolerated in its use for the sick, because these have not been sufficiently estranged from such stimulants to do without it.

To healthy persons we would recommend to abstain from its use entirely, as tea is neither nutritious, nor capable of fostering the digestion.

If any tea shall be used, let it be black tea, as this con-

tains less of the tannin, or the astringent principle, which forms the principal ingredient of all kinds of teas.

*Chocolate*, or *cocoa*, is decidedly preferable to coffee or tea as a daily drink, although it is, to some extent, an indigestible article, which needs good digestive powers and exercise to render it wholesome. It is frequently allowed to patients who have no abdominal diseases, or do not suffer from weakness of the stomach.

*Milk* is very nutritious, but not always digestible. Although nature designed infants to subsist for some time on milk, it is no stronger proof, on that account, of its greater digestibility; as it is well known that the infant's stomach is not yet weakened by stimulating food or drink, and is better prepared for the digestion of milk than that of adults: moreover, the infant draws the milk in its warm state, where the oily and cheesy parts are still organically held in perfect solution; besides, it ejects the cheesy part most always afterward, keeping only the saccharine portion of the milk, or the whey, which is the most digestible.

The practice of drinking milk at dinners is very injurious; it is more wholesome toward evening, particularly in the summer, and in the morning, when bread-and-milk forms a good diet for healthy persons.

The reason why country people can eat and drink milk with impunity, while persons living in cities dare not do the same, lies simply in the constant exercise of the former, combined with less care and hasty pursuits of life.

After drinking milk, no sour articles must be drank or eaten, as a fermentation will ensue in the stomach with all its morbid consequences.

*Buttermilk* is a healthy beverage, but it must be used with care, not allowing any acids or fruits to follow after it.

*Spiritous liquors*, wine, brandy, etc., must be considered as medicines, never as drinks, which might be adopted for

daily consumption with impunity. And, in a medicinal capacity, they are truly beneficial, while, in the other, they have proved to be the greatest curses of mankind.

But enough has been said and argued on this point, and it would be useless here to repeat the proofs, that the habitual use of spiritous liquors ruins both body and soul.

As a medicine, they are prescribed by the physician ; for instance, brandy, after having eaten indigestible articles, or as a stimulus in Asiatic cholera ; wine, after debilitating diseases, or after great loss of blood ; and ale or porter for nursing women, etc. We advise strictly to adhere to the principle, "Touch not, taste not, handle not."

#### SLEEP.

This interruption of the voluntary activity of the bodily powers is as necessary for their renovation and strength as food and drink ; in fact, in sleep the solidification of lost particles takes place — a process, without which the system soon would lose its existence in evaporation. It is in the sleep that the blood gives off to the different organs particles of solid mass, which fill up the places of those wasted during the wakeful state, and, after every organ is thus renewed and supplied, treasures up the balance in the cellular tissue in the form of fat ; in the sleep, all the nourishing qualities of the blood are carried to the places where most needed ; this is the reason why, after a healthy sleep, the system feels invigorated again to renewed action. If we would eat and fill our blood with nourishment, yet have not sufficient bodily exercise to waste strength, we would soon produce a disturbance of the equilibrium, which must exist in our system between loss and supply. In the following order we should regulate the hours of life ; first, *work* (mental and bodily exercise for every healthy individual, without exception) ; second, *eating* (the principal meal



ought to be had toward evening, after the labor is done); third, *sleep* (soon after eating, in order to permit the nourishment in the blood to crystalize into the different solids of the system.)

The time which should be passed in sleep varies according to age, occupation, and constitution. Children need more sleep than those of a maturer age, where the growing, or solidification of the system is not so much required. Vigorous, mental activity needs longer sleep to recruit than mere bodily labor; because the nervous energy, which facilitates solidification, becomes more exhausted by mental than bodily fatigue. The lymphatic constitution needs more sleep than the nervous and bilious, because the former having less nervous energy, requires longer time to convert, during the sleep, the blood into muscle, nerve, sinew, etc.

There is no normal quantity of hours to repose in sleep; as a general rule, it may range from six to seven hours in twenty-four; circumstances alter these, however, frequently; every one must try to find out which number of hours will suit for him.

The best time for sleep is, without doubt, from nine or ten o'clock, before midnight, until three or four o'clock in the morning. Early rising has so many advantages in point of health and happiness, that it hardly needs a further recommendation here.

The place of sleep ought to be chosen from among the best in a house. The lower floors are never fit to contain bed-rooms, as the confined air, during the night, in the lower part of a house, is unfit for respiration. The sleeping apartment should be the best located and the largest room in a house; an eastern and southern exposure is the best; during the day, its windows must be kept open, and the bed-clothing aired; the best surface to repose on is the hair matrass, which combines all the necessary requisites

for this purpose. During the sleep, the cover ought to be sufficient; if possible, let it consist of quilts, whose number is regulated as to comfort.

Some persons have the erroneous idea that, to lay on the back during sleep, is the most healthy position one can assume; the most natural and healthy position is to lay on the side, the very best is the right side. But this varies as to habit, and, if so, does not signify disease. By laying on the back, the spine is pressed and heated, beyond what may be good for it. Lay with the head supported, by one pillow, slightly elevated above the rest of the body.

#### AIR AND EXERCISE.

The quality of the air which fills our lungs has the greatest influence on the healthy state of our system; as on it depends the thorough change which the blood must undergo, from the venous to the arterial. The atmospheric air consists of one-fifth part oxygen and nearly four-fifths nitrogen, and a small quantity of carbonic acid gas. The first, or oxygen, is the only life-sustaining fluid, while the nitrogen and carbonic acid gas, each by itself, are life-destroying; but, in the above mixture of the three, the air becomes just adapted to the wants of our physical existence. Alter this proportion in the least, and sickness and death must be the inevitable consequence. It is of the highest importance, therefore, to have the air which we breathe as nearly like that which nature has prepared for us as possible. This, however, we can only accomplish by a thorough system of ventilation, which carries off the vitiated air, and permits fresh air to enter our apartments freely. The rooms for the sick are no exception to this most necessary rule. The more air a patient can have, the sooner he will recover. The only precaution necessary in such cases, is to prevent a draught from passing over the patient.

We direct the attention of our readers particularly to this subject, and invite them to make themselves acquainted with works treating extensively on the necessity and the art of ventilation ; as in many a situation in life, it will be of the utmost importance for them to carry the theories of ventilation into practical execution ; for instance, in building dwelling-houses, churches, halls for meetings, etc. We would here make one remark in regard to churches, which receive their ventilation at the sides, by opening the windows at the time of meeting. This seems to me a practice fulfilling well enough its immediate object—the ventilation—but doing harm in another direction, namely, creating a draught, which passes over part of the assemblage, thereby rendering their seats very uncomfortable, and dangerous for health and life. An edifice with high ceilings ought to be always ventilated from the top of the ceiling, where the heated air will sooner be found than at the side or below.

In short, we would advise every body to try always to breathe as pure and fresh air as possible ; he may be in health or in sickness.

Every thing which has a tendency to deteriorate the air must be avoided and removed. Exhalations from marshy grounds, damp cellars, or any place where animal or vegetable decomposition is going on, are frequent sources of disease and death. Every thing must be avoided which renders the air moist and damp, or prevents a free circulation all around the places where we dwell. The more air, the better ; particularly in childhood, where the healthy development of the system depends so much on the lively stimulus which the plentiful consumption of the oxygen gives to the energies of the system. An infant, instinctively, wants to be carried in the open air, where it delights to breathe the invigorating fluid. Parents should send their

children in the free, open air as often as possible; as nothing more contributes to health and a good constitution than exercise in the open air.

*Activity, motion, exercise*, is a principal object and fundamental condition of organic life. Without it, there is no energetic expression of the functions of the organs, which become torpid, and act sluggishly. Without exercise, the muscle never will swell and become strong and well-formed; the joint never will become supple. Without exercise, the appetite never will be stimulated, the nerves invigorated, and the blood system put in vigorous motion. Exercise, however, brings life and energy to every part of the system. Look at the blacksmith's arm—how muscular, strong, and healthy. Bodily exercise and labor is required of every one—he may be rich or poor, learned or ignorant—to keep healthy, and become strong.

It should be the particular care of parents, besides mental culture, not to neglect the bodily education of their children. The training in all the arts of life must have reference to the wants and necessities of the present and future well-being of those dear ones, placed under our especial guardianship. Gymnastic exercises now form a branch of the education of our youth, and should be practiced regularly and constantly, commencing with the lighter exertions, until the severer feats can be performed with ease. Walking, and running at a moderate gait, is perhaps the best and most suitable exercise for adults, besides that exercise which the daily occupations offer. It ought to be borne in mind, however, that no severe exercise must be undertaken directly after meals; as a full stomach, for digestion, requires suspension of muscular activity.

#### CLOTHING AND OCCUPATION.

In no department of civilized life has the fancy of man

committed so many errors, leading to destruction of health and life, as in that which relates to the external covering of the body. In choosing a garment, one must always first remember its purpose and use, before he follows the dictates of fashion. If fashion was directed more by the utility than by the oddity and fancifulness of clothing, its reign might be tolerated ; but this is, unfortunately, not the case. It is, for instance, fashionable among ladies to wear thin-soled shoes ; yet there is no greater destroyer of human life than a thin-soled shoe—not even pestilence and vice.

The object of clothing the body, beside for decency, is to shield it from the heat and cold of weather. To this purpose all other considerations, as for beauty, convenience, etc., ought to yield the preference ; and even healthy persons should not easily depart from a rule so imperative and necessary.

Cotton, linen, and wool, manufactured into different kinds of stuff, are the articles mostly used for clothing. Linen is the best article for summer, and should not be laid aside in winter, except by persons subject to rheumatism and frequent colds.

Cotton has won its favor almost into universal use, and its applicability for summer and winter wear is generally admitted. Yet it is not a sufficient substitute for the woollen goods, particularly flannel, when persons liable to rheumatic attacks need, besides protection from cold, a covering which stimulates the skin to a higher degree of action, and absorbs, at the same time, its perspiration. This the woollen fabrics, especially flannel, etc., do, better than any other, and are, therefore, indispensable to persons of a tender skin, and to those liable to great exposure to the extremes of cold and heat. Wool is a bad conductor of caloric, and, therefore, the best calculated to retain the heat

generated within the body upon the surface, as well as to protect it from external heat.

The quantity of clothing must be regulated according to the seasons ; but in such a manner that its change dare never be in advance of the coming season, but only after it has fully set in. A change of under-clothes before June is seldom without serious consequences ; while the summer dress may be retained until late in the fall and winter, without great inconvenience and detriment.

Clothing must be made so that it does not incommode the body either in motion or rest.

The human system is so admirably constructed in all its parts, that none presses or weighs heavily upon the other ; and in this way, only, the perfect harmony and healthy condition of all its functions is rendered possible. For the same reason, however, does it not permit any external pressure, by clothing, etc., without serious injury. If the boots or shoes are too tight, painful corns appear, and embitter the hours of the sufferer ; if the waist is laced too tight, lung complaints are its fearful and destructive consequences. It would be unnecessary to enlarge upon this subject ; it is known too well by every one, at the present time.

In general, keep the head and breast cool ; the feet and stomach, however, warm. If the dress has become wet, hasten to change it. Be not careless in this respect ; as frequently negligence in undressing, after having become wet, has been the cause of lingering diseases, or even death ; particularly the feet require our whole attention in this respect. Young girls, at the age of change, must be especially watched well as regards the protection for their feet and abdomen.

After violent exercise, we must cool off by degrees, covering ourselves more, as we become cooler.



The *occupation* of a person has a great and increasing effect on his health, mentally as well as bodily. That this is undoubtedly the case, has been clearly proved by the statistical tables, carefully made for the purpose of showing the relative longevity of persons pursuing different avocations of life.

The following statistical table, made at Berlin, in 1834, and taken from Dr. Tarbell's "Sources of Health," is very instructive in this respect :

Of 100 Clergymen, attained the age of 70 years					
	and upward,	.	.	-	42
"	"	Farmers	.	.	40
"	"	Commercial men	.	.	35
"	"	Military men	.	.	33
"	"	Lawyers	.	.	29
"	"	Artists	.	.	28
"	"	Teachers	.	.	27
"	"	Physicians	.	.	24

It is evident, from the above, that the quiet pursuits of life, which allow of a constant and certain amount of mental and corporeal activity, but of not too exciting a character, such as the minister of the Gospel, and the farmer, follow the most healthy pursuits of life ; while the physician, the very opposite of these two, as regards regularity and uniformity of action, is the lowest in the scale of his years of life. There is no doubt that the above table is correct ; and if so, it teaches volumes to every one who has either already chosen a pursuit of life, or has yet to do it. The former must adhere the more to the strictest rules of health, if he has chosen a less favorable profession ; the latter may benefit himself infinitely by selecting

the more healthy pursuit of life, if he had to choose between two.

A greater sense of duty, and a lesser degree of unchecked ambition, might have saved many a valuable life from an early grave. It is our duty to live as long and as usefully as possible.

### HYDROPATHY.

Hydropathy is the name for a method of curing diseases by the application of cold water, in various ways, internally and externally, as circumstances may require. Although the advantage of cold water in disease was known in the earlier times, yet its systematic use for that purpose was never looked upon as an essential part of the healing art, and, therefore, no decided results were elicited. To Prissnitz, a simple farmer in Germany, belongs the honor of having first boldly proclaimed, and sustained by facts, the high qualities of cold water, as a remedy for diseases. And so rapid was the progress of this new and startling doctrine, that while its propounder is still enjoying his fame and fortune, his discovery has been already adopted in almost every part of the world.

The appearance of Hydropathy, as an auxiliary branch of medicine and hygiene, seems to have been providential in our time ; as it was never before more needed, and never would have been appreciated as much. Our generation, so weakened, physically and morally, by a thousand influences which we could hardly enumerate here, needed, with the change of medicine which was consummated in the discovery and cultivation of Homœopathy, a system of dietetic rules and strengthening appliances, which should be as much *according* to nature as those formerly prescribed had been *against* nature. In this hygienic or preventive feature of the cold water lies one of its greatest blessings ;

another one, however, presents itself in its healing powers, for a number of diseases. Although its healing effects are wonderful, and entitled to all consideration and praise, yet its universal application, as the only curative agent, superseding direct medication, is impossible ; because its principle is more negative in its effects upon disease in general. The cold water, in strengthening the powers of nature, prepares them to throw off disease, or, in other words, to produce a crisis, which enables nature, in a struggle with the disease, to throw it off. This principle of Hydropathy is true in chronic as well as acute diseases ; but in the latter, its application is more limited, and this in proportion to the acuteness and severity of the diseases. To treat Asiatic cholera, for instance, or cholera infantum, with cold water alone, might be, to say the least, a questionable and hazardous undertaking. Neither would it be necessary or right to trust, in the thousands of medical cases, to one healing agent alone, which at best only assists in keeping up sinking nature, when we have the knowledge of other agents which have a direct annihilating effect upon the disease itself. And this is Homœopathy, whose application in diseases do not prevent the use of cold water ; but, in most cases, prescribes its use, to strengthen nature, that the remedies may destroy the disease in a direct or specific manner. These two systems, properly combined and practised together, will form the most complete code of medical rule and action which has existed, as yet. Allopathy and Hydropathy can never act together ; because the principle of the former, to break down the natural strength, is too antagonistic to the objects of the latter, which tries to elevate instead of diminishing the strength of nature. There is no other alliance left, neither could any other be conceived, as natural, harmonious, and effective as that of Homœopathy and Hydropathy, each one supporting and

strengthening the doings of the other. Providence was evidently at work in permitting the former to be discovered by the most learned and philosophic mind of the age—the illustrious Hahnemann ; and the latter to be practised first by an unlearned, but unsophisticated peasant : the first being the result of deep reasoning and research in matters of nature, the latter of simple and clear observation. The power of this double lesson, given to the medical world almost at the same time, was too overwhelming not to arouse it from perversion and affectation.

Hahnemann at first broke down, with a giant's strength, the learned fetters which for ages had encompassed the best minds among the medical profession. He opened and showed the true way ; a new era had begun, and men, liberated from old prejudices, were willing to try any thing offered with some degree of plausibility, reserving for themselves the final judgment.

After such changes were wrought in the mode of thinking, it was possible that Mesmer could succeed and find followers, and that Oertel and Prissnitz could draw the attention of sufferers to the remedial powers of water.

The pretensions of Hydropathy to constitute a perfect medical system, in which light some of its most modern practitioners have tried to establish it, must and will be abandoned, as they have neither a theoretical basis nor practical demonstration. In fact, efforts of this kind, if persisted in obstinately, must finally result in detriment to the good cause itself, diminishing the reputation of the cold water, in those spheres where its application is deservedly of the greatest benefit. And this range of action is by no means so small or so inferior as to make the exclusive friends of Hydropathy seek for a larger dominion, even beyond the reach of its principle or easy application. It needs but a slight acquaintance with the workings of

Homœopathy, in comparison with Hydropathy, to come to the conclusion that the former could sooner dispense with the assistance of the latter, than *vice versa*, but that a judicious union of both insures the most blessed results.

In a great many cases, the cold water may be applied alone, without giving any medicines. This is particularly the case in those chronic forms of diseases, which have been treated or produced by allopathic medicines; such as chronic rheumatism and gout, syphilis, mercurial diseases, dyspepsias, chronic diarrheas, constipation, etc. But even in such cases a recourse to a full use of Hydropathy is not necessary, if the patient seems to possess yet a sufficient amount of natural strength to produce a reaction. Homœopathic medicine, rightly administered, will be found efficacious in a majority of such cases, as the practice of thousands of homœopathic physicians proof daily to an evidence.

But where the patient's constitution has been weakened by the disease, strength and material power are necessary to overcome the disease. In such cases, the cold water is the only agent which can safely and quickly accomplish that.

In many acute diseases, the cold water can be used to mitigate the intensity of the symptoms, in subduing local irritation and fever, in softening hardened surfaces, and repelling congestions. In other acute diseases, it may be applied as an accessory remedy, without interfering in the least with the specific action of the homœopathic medicines, as in sore throat, inflammation of the brain and other organs. In acute diseases of the skin, such as measles, scarlet, etc., the application of cold water on the diseased surfaces must not be attempted, except prescribed and attended to by a physician, as repressions of these eruptions on internal organs may take place, followed by dangerous consequences.

We will mention here those diseases, in the treatment of which cold water may be applied, as an accessory, without hesitation. The mode of application for each disease is explained further, below.

*Inflammation of Internal Organs.*

If there is inflammation of the brain, eye, throat, bowels, liver, spleen, or kidneys, apply, besides giving the necessary homœopathic remedies for these diseases internally, cold bandages externally, as near the organs affected as possible. Their application is described below. In the commencement of those diseases, the cold bandage applied in the evening is very beneficial in its results. If applied after the disease has taken too deep a hold on the system, it is of less benefit.

*Congestions of the Head, Breast, and Abdomen. (Piles.)*

Are frequently mitigated by the application of the cold water bandages, often changed, and foot-baths; the sitz-baths, if the abdomen is the seat of the disease.

*Rheumatism in its various forms, Gout, Tic Douloureux, and Sciatica.*

Are diseases, where hydropathic applications may be used beneficially with the homœopathic treatment. The mode of applying the water varies in these diseases according to the constitutionality of the patient and the severity of the case. The cold water bandage, wet sheet, and local baths are mostly necessary, although sometimes warm or lukewarm applications of water may be found more serviceable.

*Constipation and Chronic Diarrhea.*

Allows of mitigation by hydropathy; the former by the



wet sheet, drinking of cold water, and a good deal of exercise; the latter by drinking cold water and using sitz-baths, with exercise and a suitable diet.

*Different Modes of Applying the Cold Water.*

In regard to this subject we prefer to lay before our readers what one of the most experienced hydropathists—Dr. Weiss, of Freywaldau—has communicated, in his Handbook of Hydropathy:

1. *Ablutions.*

These may be local or general; they are performed in the following manner: The naked hands, or, better, a large sponge or woolen cloth is dipped into a vessel containing cold water, placed upon a chair. The sponge is to be gently pressed, and then conveyed, for some few minutes, rapidly over the whole surface of the body; water may also, at the same time, be poured over the head; but not every one is able to bear the latter application, especially in the winter. Another method of performing ablution with cold water, consists in wrapping a linen sheet, dipped into cold water, round the body, and thus washing all the parts; this process is more powerful, abstracting more heat from the body. In pursuing this, or any other mode of ablution, it is advisable to stand in a spacious vessel, so that the water which runs off may not wet or soil the room in which the operation is conducted.

The best time, undoubtedly, for these ablutions, is the morning. They are to be performed immediately after rising from bed, when the temperature of the body is raised by the heat of the bed. The sudden change favors, in a great measure, the reaction which ensues, and excites the skin, rendered more sensitive by the perspiration during the night, to renewed activity. In some cases, and under cer-

tain conditions, more than one of these ablutions becomes necessary; the same operation may then be repeated at different intervals. In most cases, a second ablution before going to bed will suffice. Local ablutions will have to be repeated most frequently, where we wish to produce increased reaction; even in these cases, the temperature of the body, or its natural warmth, should be restored before proceeding to a second ablution. To increase the beneficial effects of this washing, it should be accompanied by friction during the process; this is also essential immediately after it. Quite as necessary is exercise in the open air, if circumstances will in any way permit it. Very great invalids only may be allowed, after washing, to retire to bed.

Ablutions are, for the most part, preparations for a more powerful system of treatment, in order to accustom the body, by degrees, to water which is absolutely cold; tepid ablutions are, therefore, to be recommended at first, especially to irritable and weakly individuals, or such as have never brought cold water in contact with their bodies.

Ablutions continued for a quarter of an hour, or longer, act as a stimulant and refrigerant; those of a shorter duration have a strengthening and exhilarating effect, and also the property of equalizing the circulation of blood, as may readily be perceived after general ablution of the whole body.

Cold ablutions are fitted for all constitutions; they are best adapted for purifying and strengthening the body; for women, weak subjects, children, and old age. Even in pulmonary complaints they produce alleviation, and even perfect amendment, where these diseases have not made too great a progress.

The room, in which the ablution is performed, may be slightly heated, for debilitated patients, in winter, to prevent

colds in consequence of too low a temperature of the apartment; this exception is, however, only admissible for very weakly persons. Generally speaking, ablution may be performed in a cold room, especially where persons get through the operation quickly, and can, immediately afterward, take exercise in the open air. After ablutions, as regards mildness of operation, follow

## 2. *Shower-Baths.*

These baths are taken in a machine, or box (Schneider's bath), constructed for the purpose. The internal arrangement of this machine, or bath, is such, that, on opening the cock closing the pipes which communicate with a reservoir, the water is brought in contact with different parts of the body at the same time, in the form of many fine streams. Very weakly or irritable people may begin with tepid water, and they will soon accustom themselves to cold water, as these baths produce a very grateful impression. Those who can not obtain a proper machine, may stand in an empty bathing vat, or other vessel, sufficiently large, while an assistant, standing on a chair, pours water over them from a common watering-pot, which answers the purpose perfectly.

The action of these baths consists in a general shock to the nervous system, and to the skin; in consequence of which, the secretion and excretion is promoted, and the whole economy benefited. As the action of shower-baths is closely allied to that of ablutions, they are justly preferred to these, by many people, because their effect is milder, and more grateful, and the water, in the form of rain, is brought in contact with all parts of the body at the same time. They are to be recommended, chiefly, in the diseases requiring repeated sweatings for their cure; for patients who, in consequence of congestions, and diseases of the

chest, can not bear the full baths after the process of sweating. These baths deserve recommendation to families. Children may be best accustomed to cold water in these machines, where the temperature can at first be raised, and then gradually decreased.

### 3. *Partial, or Half-Baths.*

All baths, where the common bathing vats are half, or three parts filled with water, are thus denominated. The half-baths serve, frequently, as a preparation for the full-baths (Vollbäder), or for a more active system of treatment; they have, therefore, a higher temperature, between fifty-nine and seventy-seven degrees of Fahrenheit. They should be continued from five minutes to an hour, or more, according to the purposes we have in view. If these baths be intended as a preparation for more active treatment, they must be of short duration, as also for persons who can not bear cold baths, or full-baths. If our object be to produce a derivative effect, to remove congestions from other organs, the duration of these baths must be regulated by their effects. The patient must remain in them until revulsion is produced. Neither the temperature, nor the length of time, can be determined beforehand; this must be regulated always by the constitution, the nature of the disease, or the obstinacy of the case: generally speaking, a quarter, half, or a whole hour will suffice.

If our intention be to call forth a higher degree of reaction, or even fever, by these baths, the temperature must be lower, their duration extend from one to three hours. Baths continued for so great a length of time must only be used by the advice of a practical hydropathist, as they not only considerably derange the organs of digestion, but produce sufferings with which the patient was unacquainted before.

The whole time the patient is in these baths, he should continually rub himself with the water contained in the vat, extending the friction to the parts above and under the water, that he may not take cold. If this mode of application be intended as preparatory to the use of cold baths, or to produce increased reaction, it is advisable to pour a few buckets of cold water over the patient before he rises from the bath.

Neither half-baths, nor any other kind of bath, are to be taken on a full stomach. Exercise, further, is especially to be taken after long continued half-baths; it must not, however, be neglected after the use of cold water in any form.

#### 4. *Full-Baths.*

For these baths, spacious and deep receptacles are necessary; they should admit of freedom of motion, and fresh water (if possible) should uninterruptedly flow into them.

That the body must be, in a certain measure, prepared for their use, I have already mentioned; and even after a proper preparation, as caution, it is desirable to wash the head, or pour cold water over it before entering the bath. This rule is to be strictly observed where the patient has perspired for a length of time, or where he is suffering from congestions of the head and chest. The latter circumstance requires our especial consideration, where these attacks are increased after the first baths, notwithstanding the necessary preparation. Such patients must be spared the further use of them, to prevent dilatation, or rupture of bloodvessels. Only young, robust persons may, without injury, venture, after protracted sweating, to plunge suddenly into the bath without washing the head. It is, moreover, necessary that every patient should enter the bath as soon as possible after gently and quickly cooling his head and

chest; for all unnecessary delay is attended with pernicious consequences.

The length of time the patient should remain in the bath varies in different cases; half a minute, or one minute, is generally sufficient; an experienced hydropathist only may prescribe full-baths for a lengthened period. During the bath, the patient must exercise the members of the body. Immediately after leaving the bath, he should quickly dry the whole body, using friction to promote reaction. If possible, he should perform the rubbing himself; and, where this is impracticable, he should be aided by an assistant. Exercise, after dressing, is next required, where circumstances will allow it, in the open air, to further reaction also.

Cold full-baths are indicated in all those diseases where augmented reaction, invigoration, or a shock to the nervous system is to be produced, where the warmth of the whole body is to be equalized; where all secreting and secerning vessels are to be invigorated; where the circulation of the blood is to be determined to the skin for the elimination of morbid matter.

These baths are to be avoided, or used, at least, with caution, in all congestions, in inflammations of internal vital organs, in diseases of the chest without exception, in certain head affections, in cases where a very active crisis ensues, and in all those where violent excitements or shocks would prove injurious.

#### 5. *General Plunging-baths.*

The plunging-bath should be taken, like the former bath, in a large reservoir, or trough, filled with water. The patients, generally such who can not of themselves enter the bath, are placed upon a chair above it, arranged by means of ropes and pulleys to move upward and downward,



so that the patients in the sitting posture may be plunged into the water as deeply and as often as is necessary. As a substitute for an apparatus of this kind, the patient may be laid upon a sheet, held by several persons, which is quickly plunged into the water, and again withdrawn. The intention of these plunging-baths is pretty much the same as that of the former; but their action is more stimulating, and the shock severer than in those cases where the patient can enter the bath unassisted, and use exercise while bathing. Hence one, for the most, five plunges, suffice to cool the body; in obstinate nervous fevers, however, they are occasionally to be repeated several times in the course of the day.

#### 6. *Local-baths.*

By this term, we understand baths of tepid or cold water, into which a portion of the body is immersed for a certain period. Their action is more powerful than that of local ablutions. We will commence with

#### 7. *Head-baths.*

They are applied, according to the object we may have in view, to different parts of the head, in the form of eye-baths, ear-baths, or derivative baths. The use of these baths is attended with some inconvenience. A large dish or basin is to be filled with water, and placed on the floor at the upper end of a mattress. The patient lays himself at full length upon the mattress, and immerses the part of the head to be bathed into the water; *e. g.* if it be an eye-bath, the whole face is to be introduced into the vessel filled with water. In this case, it is necessary, moreover, to open and shut the eyelids frequently, in order to bring the water in contact with the eyeball. The duration of these cooling and strengthening baths is from two to six minutes, but

both their duration and temperature are subject to variations. In some diseases of the eye, as we shall hereafter show, cold baths are not only inefficient, but they even augment the affection and sufferings considerably. Under these circumstances, we must endeavor to discover the temperature best adapted to the case, and make use of other deterrentive baths as auxiliaries. These and all other varieties of topical baths are to be repeated, according to circumstances, three to five times a day.

We use for these purposes, also, boat-shaped eye-baths, resembling liqueur-glasses, and adapted to the shape of the eye; but baths in these vessels are less worthy of recommendation than the former bath, inasmuch as they contain a less quantity of water, and press, moreover, on the appendages of the eye.

The ear-bath is taken in a similar manner: the side affected, or, where both ears are diseased, both sides are to be alternately immersed into the water, so that it may enter the meatus. These baths should, with few exceptions, be cold; their duration should extend from six to ten minutes. A more protracted use of them is not advisable, because it would produce more injury than benefit. The same applies to all head-baths.

Where head-baths are used as derivatives for violent and local pains of the head, the side of the head opposite to that affected is immersed in the water for three or four minutes. This may be repeated several times during the day, if necessary.

If the head-baths are to have an anodyne or soothing effect in gouty or rheumatic affections, they should be used at a temperature of fifty-nine degrees to seventy-two degrees of Fahrenheit. One side of the head, and then the occiput, afterward the other side is to be plunged into the water, and this proceeding is to be repeated several times

until the desired alleviation ensues. In cases of necessity, this process may be repeated several times in the course of the day.

### 8. *Sitting-baths.* (Sitz-baths.)

For these baths we use water of various degrees of temperature. The vessel for this purpose is rather inconvenient, but so constructed that the patient can remain for the necessary time in it in the sitting posture. The dimensions of the vessel should be the following: height of the pedestal, four to five inches; the inner depth of the vessel nine to ten inches; height of the back, six to eight inches; whole breadth of the vessel, twenty-two to twenty-four inches. These baths are made of wood or tin; the latter, however, deserve the preference. The vessel in which the bath is taken should be filled with water, until it reaches the navel of the patient when in the sitting posture. In especial cases, a greater or less height of water may be requisite. During the bath, the upper part of the body is to remain covered, the shirt should be turned up, and the legs and feet are to be enveloped in a woolen coverlet. While the person is in the bath, he should rub the abdomen with a woolen cloth, to increase the action of the skin, and to facilitate the passage of flatulent collections. The action of sitting-baths varies, partly according to the length of their duration, partly according to the temperature of the water. Where they are desired to have a tonic action, the temperature should be from fifty degrees to fifty-nine degrees of Fahrenheit, and they should be continued from ten to fifteen minutes. Of course, they are to be repeated frequently during the day. To act as a stimulant, and to produce more powerful reaction, they must be continued for the same length of time; but their temperature must not exceed forty-one degrees to forty-four degrees of Fah-

renheit. In summer, this temperature may be obtained by the aid of ice.

Where the sitting-baths are to act as derivatives, determining the blood from parts which suffer from congestion, the patient must remain twenty minutes to half an hour in the bath. It is sometimes necessary, during the bath, to adapt cold applications (*umschläge*) to the parts affected; this is the more necessary, if the congestions increase during the bath.

If the sitting-baths be intended to produce a solvent effect, a moderate temperature of fifty-nine degrees to sixty-eight degrees of Fahrenheit, and rather a lengthened continuation of them, say from half an hour to an hour or more, will be required. It is, moreover, advisable, that patients suffering from obstructions or hemorrhoids, should sit in deeper water; it may, in this case, extend beyond the umbilicus.

For sitting-baths in rivers, a peculiar apparatus is necessary. A board with a round hole, about eight inches in diameter, is fixed about five or six inches below the level of the water; on this aperture the patient is to sit. At a distance of more than one fathom from this board the water is to be confined by means of any suitable contrivance, so that it may be made to flow with force against the abdomen during eight, at the most, fifteen minutes. In all abdominal diseases, and in debility of the organs of generation, these baths are of extraordinary benefit.

Whatever object we may aim at in the use of these varieties of sitting-baths, we must not lose sight of the rule—that they are never to be taken immediately after eating, (unless especial cases call for an exception), as they will otherwise derange the digestion, and produce irregularities in the evacuations. The best time for the use of these baths, is an hour before dinner, or before going to bed. In

the latter case, they offer the advantage of securing a night's rest to the patient. Generally speaking, two sitting-baths a day will suffice ; in particular cases, especially, if not persevered in for a long time, five to six may be taken during the day. Exercise in the open air is to be strictly recommended both before and after these baths.

### 9. *Foot-Baths*

Are taken in a small tub, or in the same vessel as the sitting-baths. The temperature and volume of water must depend on the action we wish to produce. If these baths are to act as stimulants, they must be very cold, not exceeding forty-one degrees of Fahrenheit. The water should have a depth of one, at the most two inches, and the bath must not be continued longer than five minutes. They may be repeated several times during the day, according to circumstances. Rubbing, immediately after the baths, to promote speedy reaction, and also exercise after them, are indispensable. If the object of the foot-bath be to determine from the head or chest, the vessel must not be too large, in order that the water may become somewhat warm during the bath, which may be continued, in that case, from twenty minutes to one hour. In this case, the water must also not be as cold as for the former purpose, and should cover the ankles. If congestion increase during the bath, cold applications should be at the same time applied to the parts affected. Friction and motion of the feet are absolutely necessary during and after the bath.

A variety of other partial baths, adapted to the different members of the body, are brought into operation in a general course of treatment by cold water. The effect desired is obtained partly by the duration of the bath, partly by the temperature of the water, as I have before mentioned in speaking of local baths in general. Where the baths

are to act as stimulants, the water must be very cold, not exceeding forty-one degrees of Fahrenheit; they may, however, be repeated several times in the course of the day, if requisite. They are to be continued from five to eight minutes. By friction, immediately after the bath, we arrive more speedily at the desired end.

If the intention of these baths be to produce a soothing effect, their temperature must not be below fifty-nine degrees of Fahrenheit; they may, however, be continued for a longer period of time; fifteen to thirty minutes will, in most cases, suffice; they may also be repeated on the return of the attacks of pain. Where their action is to be derivative, the temperature should be from forty-six to fifty-nine degrees of Fahrenheit, and they should be continued for half an hour to one hour or longer. If they are to produce this effect gradually, and not rapidly, we frequently resort, at the same time, to applications (*umschläge*) to the part affected. Cooling applications, frequently repeated, will often lead to the desired result, and, at other times, applications producing warmth will be most serviceable; we should, therefore, consider whether the object be to produce cold or heat.

#### 10. *The Drop-bath.*

This term is applied to single drops of water falling from a height of several fathoms. For this form of bath, a vessel is used filled with very cold water, and furnished with a very small aperture, through which the water passes in the form of drops. The small aperture should be partially closed by a plug, to prevent the drops from following each other in rapid succession. By these means their operation is considerably increased, and it becomes yet more potent if we allow the drops to fall upon a particular part at certain periods, and rub the part during the intervals. The



reaction about to commence will, indeed, be thus interrupted, but will afterward make its appearance in a more powerful and energetic form.

The violent excitement and irritation of the nervous system produced by these baths, render it necessary to restrict the use of them to half an hour; nor are they, indeed, adapted for vital parts, or such as are abundantly supplied with nerves.

They are often used with more effect in obstinate and chronic cases of paralysis, than the douche or affusion, with which they may alternate. Powerful and continued friction with a horse-hair glove is never, in this case, to be neglected after the baths.

### 11. *The Douche.*

This description of bath is prepared with the aid of mechanical contrivances, by means of which a stream of water is made to fall upon the body with more or less force. In many respects it is most advantageous to make use of a natural fall of water for this purpose; we can then conduct the water simply into a channel, giving it a fall of twelve to twenty feet, and to the stream a calibre of half an inch to five inches. These simple douches are far less disagreeable to the sensations of the patient than those of artificial construction. In the former, the patient can turn himself freely and alter his position so as to expose any part of the body to the stream. Douche-rooms, admitting, by their construction, of the access of the air from above, produce an agreeable sensation, especially in summer, and are very beneficial in their action. After the first time of using these baths, the dreadful ideas which many patients preconceive of them quickly disappear.

Our chief consideration in the use of the douche should be to guard against applying it to the body when quite

cold, or when in a state of perspiration after active exercise. The patient, after undressing in a moderate temperature, steps below the falling stream, attempting to receive it in the palms of his hands, that the whole force and volume of the water may not fall upon his body immediately, which is not, to say the least, at all times agreeable. After having thus prepared himself for the more potent shock, he must expose himself to the full stream and in such a manner that the whole column of water fall chiefly on the neck and spine. From time to time he must equably expose the other members of the body to the stream ; but the affected parts chiefly, and for a greater length of time. He should be careful not to allow the stream to fall perpendicularly on the head, chest, or on the region of the liver, especially if these parts be weak, or affected with disease.

The duration of the douche must be regulated by the constitution of the patient, and the effect we wish to produce ; it should never be continued for less than one, nor more than twelve minutes.

Where a natural douche is not to be obtained, we make use of the well-known machines resembling a fire engine in shape and construction. By means of these contrivances, we may bring one or more streams of any given force into operation. Our care must be directed to the selection of cold and fresh water for this purpose ; finally, the same rules are to be observed which we laid down for the use of the former douche.

The period of the day at which these different douches (excepting those for the eye and ear) are used, varies. They are only to be taken, as an exception, fasting, or immediately after sweating, and never on a full stomach, nor oftener than once or twice daily.

Rather active exercise should be taken after the douche,

until the peculiar sensation of reaction has totally disappeared, or an uncomfortable sensation of cold accompanied by head-aches, fainting, etc., will be experienced, instead of an agreeable and beneficial glow. It is, moreover, not advisable to drink cold water immediately after the douche, because a rapid generation of heat is thus impeded, and inflammations of the stomach and bowels might be caused.

The douche, the most powerful stimulant known in hydropathy, is always applicable where excitement is necessary but attention must be paid, in every case, to age, constitution, and to the vital powers. Cases are not unfrequent, where the nature of the diseases calls for this stimulant to complete the cure, but where it must be avoided in consequence of pregnancy, pulmonic affections, etc. No mode of applying cold water is more abused than the douche; in most hydropathic establishments, even at the present day, this is, unfortunately, too often the case. But very few experienced hydropathists regard it as a stimulant requiring caution. For the most part, it is looked upon as a remedy indispensable to every cure, is brought into use injudiciously, and without mature deliberation, and continued to the detriment of all organs; the beneficial results, therefore, which a judicious use of it would produce, are, necessarily, oftentimes frustrated.

## 12. *Local Applications (Umschläge.) Wet Bandages.*

By these applications, two different effects may be produced; viz., that of cooling the part to which they are applied, or that of raising its temperature. Where they are intended as refrigerants or derivatives, the cloths must be of a size suited to the part inflamed; they should be folded six or eight times, dipped into very cold water, gently pressed before application, and are to be renewed every four or ten minutes, according to the degree of inflamma-

tion. If we can not obtain water sufficiently cold in summer-time, ice must be added to it until its temperature sinks to forty-one or forty-four degrees of Fahrenheit. This low temperature is especially necessary in dangerous inflammations of important organs; *e. g.*, of the brain. The bandages must then be continued without intermission, day and night, until danger is averted. Neglect in changing the wet cloths (*umschläge*) at the proper time, will cause fatal results. One omission of the change is sufficient to frustrate their beneficial operation; for violent reaction is only to be subdued by continued cold.

The warming applications of cold water consist of pieces of linen folded two or three times, and dipped into cold water; they must be well pressed out before application, and not changed until they begin to dry—this is the indication for their repetition. This variety of application must be not only well adapted to the part, but it is to be further secured against the access of the air by a dry bandage, in order that the reaction it produces may generate a degree of heat in the part covered exceeding the temperature of the body. The combined action of the moisture and heat thus produced is that of a solvent of morbid matter, dispersing swellings and indurations, which are thus rendered more fit for absorption and elimination. These applications are not only well adapted for the removal of tumors, but may be applied also, with great benefit, in various cases of affections of the abdomen. Their efficacy in all derangements of the digestive organs, and diseases of the liver, etc., is proved.

The applications to the abdomen vary somewhat in form from the former bandages. A piece of linen, two yards in length, and rather more than one foot in breadth, is to be doubled, sewn together, and furnished with strings, to enable us to adapt it closely to the body, and secure it in its

situation. In using the bandage thus arranged, about one foot of it in length should be wetted with fresh water, well wrung out, and applied round the body ; thus, the wet portion will extend once, the dry part twice round the body. These abdominal belts are to be worn according to the urgency of the case, sometimes for several hours in the day, or through the whole night ; in some instances, again, for weeks, both by day and night.

13. *Applications round one-half of the body, or round the whole body. Wet linen sheets.*

For this purpose, the bed must be prepared in the same manner as for the process of sweating. The wet sheet is laid upon the extended blanket, the patient lays himself at full length upon the former, whereupon it is folded round him, so as to come in close contact with every portion of the body. The patient is now to be enveloped in the blanket and the bed covering (feather beds.)

The wet sheets are of remarkable utility in all febrile diseases. In acute fevers they must be changed according to the degree of heat, every quarter or half hour, until the dry, hot skin of the patient becomes softer, and more prone to perspiration. When this symptom is observed, the renewal of the wet cloths may be delayed for a longer period, until perspiration actually ensues. The patient must then remain for several hours in this state, until uneasy sensations, and other inconveniences, render it necessary to extricate him ; but it is more advisable to keep him in the loosened envelopment until perspiration ceases spontaneously, when a tepid ablution, or half-bath, should follow. In acute eruptions of the skin, measles, scarlatina, small-pox, etc., the wet sheets are not less serviceable, when the eruption can not make its way to the surface in consequence of the dry state and heat of the skin, and of the violence

of the fever; or where the rash has receded suddenly, owing to other disturbances. In both cases, the wet sheets are of essential service; one application of them suffices, sometimes, to reëstablish the eruption. If the rash fail to make its appearance after the first or second envelopment, cold effusion is to be preferred. There are cases in which the use of this remedy may be deemed objectionable, and a continuance of the wet sheets may appear more proper; we must, then, examine the skin carefully before every change, to see whether the eruption be nearer the surface, the skin softer, and the heat abated. In the latter case, the applications are to be discontinued, that the reaction of the skin may not be disturbed. The wet sheets, followed by tepid ablutions, can not be sufficiently recommended in all diseases of children. Many severe complications are averted, or relieved, at least, by them; or, where this is not the case, the disease itself is brought more speedily to a favorable termination.

The envelopment in wet sheets is not only of great advantage in acute diseases, but is, also, an admirable remedy in a variety of chronic cases, attended with an irritable, rough, and inactive skin, and in a multitude of skin-diseases; but in all those cases, a frequent change of the sheets is seldom necessary. In using the envelopment, we generally raise the temperature of the patient, and occasionally allow him to perspire, according to the circumstances of the case. Determination to the head during the process must be removed by cold applications to that part. If the feet remain cold for a long time in the wet cloths, and show no disposition to become warm, they are to be extricated and wrapped in the dry blanket only.

#### 14. *Sweating.*

In the treatment by water, perspiration is brought about



in the following manner: The patient is undressed and laid upon a woolen coverlet or blanket, extended on the bed. A servant wraps first the one side of the blanket round the body of the patient, drawing it close in all directions; grasping now firmly with the one hand the portion in which the patient is enveloped, he draws, with the other hand, the blanket round the patient, and tucks this portion also closely under him. Care must be taken that the coverings be in close contact with the body, especially at the neck, that the heat given off by the body may be well retained, and can not escape. It is the excess of caloric thus confined which induces exhalation from the skin. Individuals who are to perspire for several hours, and can not retain their urine long, may be furnished with an urinal, placed between the legs at the time of wrapping them up. The head may be included in the covering, or enveloped even in the pillow, so that the face alone is free, provided the patients do not suffer from congestions. The head must remain uncovered on the other hand, if a disposition to congest be observed. Slight excitement of the vessels before the outbreak of perspiration generally passes off spontaneously; but where this favorable termination does not ensue, a cooling bandage (*umschläge*) is to be laid on the head of the patient, who must, at the same time, drink a little cold water. Warming applications (*umschläge*) are to be wrapped round all parts affected with nodes, gouty swellings, etc., before enveloping the patient. The use of these auxiliaries is to allay pain, which is generally more violent before the appearance of perspiration, and to excite a more copious exhalation from the parts to which they are applied.

Patients who are very restless in the blanket, and thus loosen it, should be confined more closely in the encasement by additional cloths and girths, as they would other-

wise have to remain wrapped up during half a day or longer. Persons thus enveloped being totally helpless, an assistant should be always in attendance to open the windows as soon as perspiration ensues, and to administer, every ten or fifteen minutes, as much cold water as is necessary to promote perspiration. The action of the skin must be kept up uninterruptedly, and the perspiration should pass off in form of vapor, or in drops, as long as the case requires it.

The result of this mode of treatment is pretty certain, but the time necessary for the production of perspiration is not the same in all cases; for some individuals perspire sooner than others. The season, and many other inevitable circumstances, exert, moreover, a considerable influence on the skin; thus, some patients, especially in summer, will perspire in a quarter of an hour, while others require three to five hours for that purpose. In febrile and inflammatory diseases, we frequently can not produce perspiration in less than twelve or twenty-four hours, although we change the wet sheets often, and resort to other auxiliary means.

The best time for exciting perspiration in chronic cases is in the early hours of the morning, from four to five o'clock at the latest. In acute diseases, the time depends on the fever itself, or on renewed exacerbations; it must be produced, therefore, when required, without reference to time. A repetition of the process twice on the same day, is only admissible and advisable as an exception. The indication for such practice would occur more frequently in chronic than in acute diseases; because in the latter perspiration should be kept until it ceases spontaneously, or is interrupted by other circumstances. Two sweats daily can only be recommended to the robust, and even these persons will not be able to endure the protracted sweating required

in some few troublesome cases, without injuring the skin and other important organs.

We can not determine *a priori* how long each patient should sweat. The ordinary duration in chronic cases is from half an hour to three hours, daily; but moderate perspiration is to be encouraged for a longer time in acute diseases. Critical perspiration requires the longest duration, and should not be disturbed; we should rather, in this case, do every thing in our power to aid and promote it, until it cease spontaneously.

When the patient has remained in a state of perspiration long enough, the woollen covering should be loosened about his feet and legs, to enable him to walk. The attendant then raises him in bed, and leads him to the place where he is to bathe or wash. Sedan chairs, or other contrivances, will be necessary to convey patients quickly to the bath, in serious cases. No danger is to be dreaded from the transition from heat to cold (as experience in many thousand cases has proved), if the necessary precautionary rules laid down for the use of full-baths be duly observed; for, in this mode of producing perspiration, important internal organs are not excited.

After the bath, patients who can walk, or take other exercise, must not return to bed; but should dress quickly, go into the open air, drink the quantity of water prescribed, and afterward take their breakfast. Those patients, on the other hand, who can take no corporeal exercise, must be rubbed, after the bath, for some time, the friction extending over the whole surface of the body. This peculiar mode of inducing perspiration is to be brought into use in all diseases where morbid matter is to be eliminated from the system, because the skin, as daily experience teaches us, is the organ best adapted for this purpose. At the commencement of the treatment, the perspiration is clear

and aqueous for some time ; at a later period it becomes more viscid, then varies in color, and assumes a powerful odor. The urine, at the same time, or later, in the course of treatment, assumes the same properties in color and odor ; open sores and the breath participate in this change. These symptoms usually make their appearance at different intervals, in a protracted case. Critical excretions seldom continue without intermission from their first commencement, until all extraneous substances are removed from the body ; it is true that this phenomenon is observed sometimes, but it is always a rare occurrence, and the critical discharges will not then unfrequently continue two to four months.

There can be no doubt that an incalculable quantity of morbid matter is removed from the body in the act of perspiring ; for it is often to be recognized by the organs of sense, by its color and odor during the process. Most striking is, at times, the odor of valerian, turpentine, iodine, musk, assafœtida, sulphur, and mercury. The colors observed are, generally, confined to red and yellow ; brown and blue are less frequently met with : the latter occurs often in the deposits in the urine, which may assume, also, all other colors.

### III. MATERIA MEDICA.

#### 1. ACONITE.

Acute local inflammations ; active, sanguineous congestions ; evil consequences of a chill in a dry, cold air ; affections in consequence of fright or of anger ; measles ; erysipelatous inflammations ; inflammatory fevers, even with bilious or nervous symptoms ; mental alienations, with fixed ideas of approaching death ; cerebral congestion, with dizziness ; croup, first period ; hooping-cough, first period.

GENERAL SYMPTOMS.—Shooting pains, or rheumatic, which are renewed by wine, or other heating articles ; sufferings which, particularly at night, seem insupportable ; attacks of pain, with thirst and redness of the cheeks ; uneasiness, as if in consequence of suppressed perspiration, or in consequence of a chill.

*Skin.*—Skin dry and burning ; scarlet rash ; measles ; nettle-rash.

*Sleep.*—Sleeplessness from anxiety, with constant agitation and tossing ; starting in sleep.

*Fever.*—Dry, burning heat, with extreme thirst, sometimes, especially at the beginning of the disease, preceded by shiverings with trembling ; heat, chiefly of the head and face, with redness of the cheeks, shuddering over the

entire body; shivering, for the short time that they may be uncovered during the heat; pulse hard, frequent, and accelerated.

*Moral Symptoms.*—Great agitation and boasting with anguish, discouragement that can not be consoled, cries, tears, groans, complaints, and reproaches; apprehensions and fear of approaching death; a great disposition to be angry, to be frightened; alternate paroxysms of laughter and tears; inquietude under disease, and despair respecting a cure; delirium, chiefly at night.

*Head.*—Weight and fullness in the forehead and temples, with pressing outward, as if every thing was going to issue through them; congestion of blood to the head; aggravation of the pains in the head by movement, by speaking, by rising from a recumbent position, and by drinking; amendment in the open air.

*Eyes.*—Eyes red and inflamed, with deep redness of the vessels, and intolerable pains.

*Ears.*—Buzzing in the ears.

*Nose.*—Bleeding from the nose.

*Face.*—Red spots on both cheeks.

*Throat.*—Pain in the throat, with deep redness of the parts affected, and difficult deglutition; burning and pricking in the throat, chiefly when swallowing.

*Stomach.*—Sensation of swelling, tension and pressure, as from a weight in the pericordial region and in the stomach.

*Abdomen.*—Tension and pressure in the hypochondriac region; pressure in the hepatic region; painful sensitiveness of the abdomen to the touch, and to the least movement.

*Fæces.*—Loose, watery stools; white stools, with red urine.



*Urine.*—Urine scanty, burning, dark red, and with a brick-colored sediment.

*Larynx.*—A constant desire to cough, produced by an irritation or a tickling in the larynx; short and dry cough, principally at night; spitting of blood, with the cough; shootings and pains in the chest when coughing.

*Chest.*—Short breathing, chiefly during sleep, and on getting up; breathing difficult, anxious, and attended with groans; painful stitches in the chest, chiefly when breathing, coughing, and moving; stitches in the side; palpitation of the heart, with great anxiety.

*Trunk.*—Painful stiffness in the nape of the neck.

## 2. ANTIMONIUM CRUD.

Gastric and bilious sufferings; gastralgia; corns and callous indurations in the feet.

GENERAL SYMPTOMS.—Drawing, darting, and tension, principally in the limbs; the symptoms are aggravated in the heat of the sun, after having drank wine, after a meal, night and morning; amelioration during repose in the fresh air; great sensibility to cold.

*Head.*—Dizziness, with nausea.

*Eyes.*—Red, inflamed eyelids; inflammation of the eyes, with itching and nocturnal agglutination of the eyelids; redness in the corners of the eyes.

*Ears.*—Deafness, as if one had a band over his ears.

*Mouth.*—Tongue loaded with a white coating.

*Appetite.*—Bitter taste; thirst chiefly in the night; loss of appetite.

*Stomach.*—Eructations, with taste of food.

*Abdominal Region.*—Inflation of the abdomen, with a sensation of fullness, chiefly after a meal; violent, cutting pains.

*Fæces.* — Diarrhea, with cutting pains, generally watery ; constant secretion of yellowish-white slime from the anus.

*Urine.* — When coughing, involuntary emission of urine.

*Larynx.* — Sensation as if a foreign substance were in the larynx, with inability to expectorate.

*Trunk.* — Rheumatic pains in the nâpe of the neck.

*Arms.* — Rheumatic pains in the arms.

*Legs.* — Corns on the soles of the feet, and callous excrescences on the tips of the toes ; callous excrescence under the nail of the great toe.

### 3. ARNICA.

Affections in consequence of mechanical injuries (falls, commotion, blows, etc.) ; wounds, principally those inflicted by blunt instruments ; bites ; excoriation of bed-ridden patients ; bruises, dislocations, sprains, and fractures ; accidents resulting from a strain ; stings of insects ; corns, by an external application of it, after having pared them.

GENERAL SYMPTOMS. — Pains, as from dislocation ; fainting fit, with loss of consciousness, in consequence of mechanical injuries ; paralytic state, on the left side, in consequence of apoplexy.

*Skin.* — Red, bluish, and yellowish spots, as if from contusions.

*Head.* — Whirling giddiness, with obscuration of the eyes, chiefly when getting up, moving the head, or walking.

*Eyes.* — Pain like excoriation in the eyes and eyelids, with difficulty of moving them.

*Face.* — Face pale and hollow.

*Mouth.* — Tongue dry, or loaded with a white coating ; putrid smell from the mouth in the morning.

*Stomach.* — Vomiting of dark, coagulated blood.

*Abdomen.* — Pain, as from contusion in the sides.

*Anus.* — Involuntary stools, chiefly in the night.

*Urinary Passages.*—Involuntary emission of urine; urine of a brownish red, with brick-colored sediment.

*Respiratory Organs.*—Cough, with expectoration of blood; even without cough, expectoration of black, coagulated blood after every corporeal effect.

*Chest.*—Respiration short, panting, difficult, and anxious; shootings in the chest and sides, with difficulty of respiration, aggravated by coughing, breathing deeply, and by movement; pain as of a bruise, and of compression of the chest.

*Trunk.*—Pains, as from a bruise, and dislocation in the back, in the chest, and in the loins.

*Arms.*—Pain, as of dislocation in the joints of the arms and hands; want of strength in the hands, on seizing any thing.

*Legs.*—Pains, as if from fatigue or from dislocation; tension in the knee, as if from contraction of the tendons; inflammatory, erysipelatous swelling of the feet with pain, and aggravation of the pain by movement; hot, painful, hard, and shining swelling of the great toes.

#### 4. ARSENIC.

GENERAL SYMPTOMS.—Burning, chiefly in the interior of the parts affected, or sharp and drawing pains; nocturnal pains, so unbearable that they excite to despair and fury; aggravation of suffering in the evening in bed, on lying on the part affected, or during repose; mitigation by external heat and movement of the body; want of strength, excessive weakness, and complete asthenia, even to prostration.

*Skin.*—Skin dry as parchment, or cold and bluish; ulcers with raised and hard edges; fetid smell, ichorous suppuration, ready bleeding, putridity, and bluish or greenish color of the ulcers.

*Sleep.*—Nocturnal sleeplessness, with agitation and constant tossing.

*Fever.*—Coldness over the whole body, sometimes with cold and clammy sweat; pulse irregular, or quick, weak, small and frequent, or suppressed and trembling.

*Moral Symptoms.*—Anxiety, inquietude, and excessive anguish, principally in the evening in bed.

*Face.*—Face pale, hollow, and cadaverous; lips bluish or black, dry and chapped.

*Mouth.*—Offensive smell from the mouth; tongue brownish or blackish, dry, cracked, and trembling; ulceration of the tongue on the anterior edge; thrush of the mouth.

*Throat.*—Inflammation and gangrene of the throat.

*Appetite.*—Violent burning, choking, and unquenchable thirst, with inclination to drink constantly, but little at a time; want of appetite.

*Stomach.*—Vomitings after drinking and eating; vomiting of mucous, bilious, or serous matter, of a yellowish, greenish, brownish, or blackish color; burning internal heat, diarrhea, and fear of death; excessive pain in the epigastrium, chiefly when touched; insupportable heat and burning in the pericordial region, and in the stomach.

*Abdominal Region.*—Swelling of the abdomen, as in ascites; violent cutting pains, cramp-like pains in the abdomen; colic, chiefly after eating and drinking, or in the night, and often accompanied with vomiting or diarrhea, with coldness, internal heat, or cold sweat.

*Fæces.*—Violent diarrhea, of greenish, yellowish, whitish color, or brownish and blackish; burning in the rectum and in the anus.

*Larynx.*—Dry cough, in the evening after lying down, with a wish to rise; also, after drinking, with difficulty of respiration.

*Chest.* — Breathing short ; difficult, stifling dyspnœa, and attacks of suffocation, sometimes with cold sweat, spasmodic constriction of the chest or of the larynx, anguish, great weakness, coldness of the body, pain in the pit of the stomach, and paroxysms of cough ; oppression of the chest when coughing, when walking, and when going up stairs ; violent and insupportable throbbings of the heart, chiefly when lying on the back, and especially at night.

### 5. AURUM.

Mercurial (and syphilitic) sufferings ; inflammation and caries of the bones, chiefly from the abuse of mercury.

*GENERAL SYMPTOMS.* — Hysterical spasms, sometimes with alternate tears and laughter.

*Moral Symptoms.* — Great anguish, which advances even as far as a disposition to suicide.

*Head.* — Fatigue of the head from intellectual labor.

*Nose.* — Running from the nose, of a fetid, greenish-yellow matter.

*Face.* — Swelling of the bones of the forehead, of the upper jaw and of the nose.

### 6. BELLADONNA.

Scarlatina ; sleeplessness ; inflammatory fevers, with nervous, gastric, or rheumatic affections ; erysipelas.

*GENERAL SYMPTOMS.* — The least contact, and sometimes, also, the slightest movement, aggravates the sufferings ; dread of every movement, and of all exertion ; over-excitement, and too great sensibility of all the organs.

*Skin.* — Swelling, with heat and scarlet redness of the whole body, or of several parts, chiefly the face, the neck, the abdomen, and the hands ; red hot, and shining swelling of the diseased parts.

*Sleep.* — Constant desire to sleep ; nocturnal sleepless-

ness, in consequence of excessive anguish or great agitation ; when sleeping, frequent starts, with fright, groans, cries ; on waking, headache, and aggravation of sufferings.

*Fever.*— Dry, burning heat ; pulse strong and quick, or full and slow, or small and slow, or small and quick, or hard and tense ; sweat of the parts that are covered only.

*Head.*— Fullness, heaviness, and violent pressure on the head, as if going to burst, with desire to lie down ; dartings into the head, as if from knives ; strong pulsation of the arteries of the head ; bending the head backward ; boring with the head into the pillow while sleeping.

*Eyes.*— Aching pains in the eyes and the sockets, extending into the head ; inflammation of the eyes, with injection of the vessels ; dread of light.

*Ears.*— Piercing pressure, sharp pain, pinching, squeezing, and shooting in the ears ; swelling of the parotids.

*Face.*— Face pale, sometimes suddenly alternating with red ; burning heat of the face ; dark, or scarlet, or bluish redness of the face ; swelling of the sub-maxillary glands, and those of the neck.

*Teeth.*— Sharp and drawing pains, at night or in the evening ; the touch and the open air aggravate the toothache.

*Mouth.*— Sensation of great dryness, or actual and excessive dryness and choking in the mouth ; tongue red, hot, shining, dry, and cracked, or loaded with whitish mucus ; redness of the edges of the tongue ; paralytic weakness of the tongue, with difficult and stammering speech.

*Throat.*— Excoriating, scraping, and shooting pains in the throat and in the tonsils, principally when swallowing ; inflammation and swelling of the throat, of the velum palati, of the uvula, and of the tonsils ; suppuration of the tonsils ; complete inability to swallow, even the least liquid,



which frequently is forced out through the nostrils ; sensation of choking, and spasmodic constriction in the throat.

*Appetite.*— Burning, excessive, and intolerable thirst, often with dread of all drink ; drinking with trembling precipitation.

*Stomach.*— Spasmodic hiccough.

*Abdominal Region.*— Cramp-like, contractive and constrictive pains in the abdomen, and especially around the navel, with a sensation as if the parts were squeezed or seized with the nails ; the pains force one to bend himself, and are accompanied by vomiting or by inflation and protrusion of the transverse colon in the form of a pad ; soreness of the whole abdomen, as if every thing in it were excoriated.

*Urine.*— Frequent discharge of urine, copious, pale, and watery ; difficulty of retention and involuntary emission of urine.

*Genital Organs.*— Violent pressure toward the genital parts, as if all were going to fall downward, principally when walking or sitting upright ; menses too copious and too early.

*Larynx.*— Loss of voice ; cough, chiefly at night, or in the afternoon, in the evening in bed, mostly dry, short, and sometimes convulsive ; when coughing, cutting in the abdomen ; the least movement, when in bed at night, renews the cough.

*Chest.*— Oppression of the chest, difficult respiration, dyspnœa, and short breath, sometimes with anxiety ; respiration short, anxious, and rapid ; pressure on the chest.

*Trunk.*— Painful swelling, and stiffness of the neck and the nape.

## 7. BRYONIA.

Rheumatic and arthritic affections, even with inflamma-

tory fever and swelling ; local inflammations ; inflammatory fevers, with nervous, gastric, or bilious affection, and strong excitement of the sanguineous and nervous system ; typhoid fevers, in the inflammatory period.

**GENERAL SYMPTOMS.**—Tension, drawing pains, acute drawings, and stinging, especially in the limbs, and chiefly during movement, with insupportable pains on being touched ; red, shining swelling of some parts of the body ; aggravation of the pains and sufferings, at night and from movement ; amelioration during repose ; desire to remain in a recumbent posture.

*Sleep.*—Restlessness, especially before midnight, caused by heat, agitation of blood, and anxiety ; sleep, disturbed by thirst, with bitter taste in the mouth when waking ; inability to remain lying on the right side.

*Fever.*—Cold and shivering of the body, even in bed, accompanied by pains in all the limbs ; shiverings ; with trembling, often with heat in the head, redness of the face, and thirst ; before the shiverings, vertigo and headache ; universal dry heat, external and internal, almost always with a great desire for cold drinks.

*Moral Symptoms.*—Irascibility and passion ; delirium, with ravings about the transactions of the day.

*Head.*—Great fullness and heaviness of the head, with raking pressure toward the forehead, and when stooping, a sensation as if every thing were going to fall out through the forehead ; headache, aggravated by movement.

*Nose.*—Swelling of the nose ; frequent bleeding of the nose, sometimes in the morning, or when the menses are suppressed, or even when sleeping.

*Face.*—Face red and burning ; swelling of the face, sometimes on one side only, or under the eyes, and at the root of the nose ; lips swollen and cracked, with bleeding.

*Teeth.*—Toothache, with desire to lie down ; mitigated

by lying on the parts affected ; jerking, drawing toothache, with a sensation as if the teeth were too long or loose.

*Mouth.* — Dryness of the mouth, with burning thirst ; tongue dry, loaded with a white, dirty, or yellow coating.

*Throat.* — Sensation of dryness, and great dryness in the throat.

*Appetite.* — Bitter taste in the morning ; repugnance and disgust for food.

*Stomach.* — Nausea and desire to vomit, especially after eating ; vomiting as soon as one has drank, and especially on drinking after a meal ; shootings in the stomach during movement ; burning in the pit of the stomach.

*Abdominal Region.* — Pains in the liver, when touched, on breathing or coughing ; cramp-like pains, or cuttings and shootings in the abdomen.

*Fæces.* — Constipation ; diarrhea in the summer.

*Urine.* — Urine scanty, reddish, brownish, and hot.

*Genital Organs.* — Menses suppressed ; metrorrhagia of a deep red blood, with pain in the loins and in the head ; swelling of the labium, with a black and hard pustule.

*Larynx.* — Cough, mostly dry, excited by a tickling in the throat ; cramp-like, suffocating cough, after having eaten or drank, and often with vomiting of food ; cough, with stinging in the sides of the chest, or with aching pains in the head, as if it were going to split ; cough, with expectoration of mucus of a dirty-reddish color.

*Chest.* — Respiration impeded by stinging in the chest ; stings in the chest and in the sides, especially when coughing or breathing deeply, allowing one to lie only on the back, and aggravated by any movement whatever ; heat and burning pain in the chest, with anxiety and tightness ; beatings of the heart, frequently very violent, with oppression.

*Trunk.* — Pain in the loins, like a painful weight ; shoot-

ings in the loins and in the back, aggravated by cough and respiration ; rheumatic heaviness and tension in the nape of the neck, and in the neck.

*Arms.*—Tractive pains in the joints of the shoulders and the arms, with tension, stinging, and shining, red swelling ; pain of dislocation in the joints of the hands on moving them.

*Legs.*—Drawing pains in the thighs ; swelling of the legs, extending to the feet ; pain as of dislocation, in the foot when walking.

### 3. CALCAREA CARBONICA.

Muscular weakness, difficulty of learning to walk, atrophy, and other sufferings of scrofulous children ; rickety affections ; polypus ; fistulous ulcers ; delirium tremens ; difficult dentition in children, with convulsions.

GENERAL SYMPTOMS.—Agitation of blood, mostly in plethoric individuals, often in the head and in the chest ; epileptic convulsions, with cries ; the symptoms are aggravated after washing and laboring in the water, in the evening, after a meal, and every second day.

*Skin.*—Nettle-rash, chiefly disappearing in the fresh air ; swelling and induration of the glands, with or without pain.

*Sleep.*—Restlessness from flow of ideas, or in consequence of frightful images, which appear as soon as the eyes are shut ; dreams, frequent, vivid, anxious, fantastic, frightful.

*Fever.*—Excessive internal coldness ; frequent attacks of transient heat, with anguish and beating of the heart.

*Moral Symptoms.*—Melancholy and disposition to weep and to be frightened.

*Head.*—Piercing in the forehead, as if the head were going to burst, hammering pains in the head after a walk in

the open air ; icy coldness in and on the head ; falling off of the hair : tumors in the hairy scalp.

*Eyes.* — Ulcers, spots, and opacity of cornea ; red and thick swelling of the eyelids.

*Ears.* — Purulent discharge from the ears ; polypus in the ears ; cracking in the ears, when swallowing and chewing ; hardness of hearing ; inflammatory swelling of the parotids.

*Face.* — Swelling of the upper lip ; painful swelling of the sub-maxillary glands.

*Stomach.* — Regurgitation of sour substances ; sour vomitings.

*Abdominal Region.* — Swelling and induration of the mesenteric glands ; enlargement and hardness of the abdomen.

*Fæces.* — Evacuations like clay ; diarrhea during dentition, of a sour smell, fetid, or yellowish, in infants.

*Genital Organs.* — Menses premature and too copious.

*Larynx.* — Cough at night, violent and dry, sometimes even spasmodic.

*Chest.* — Pain as from excoriation in the chest, especially on breathing and being touched ; palpitation of the heart.

*Trunk.* — Hard and painful swelling of the glands of the neck.

*Arms.* — Drawing pains in the arms, also at night ; boils on the hands and the fingers.

*Legs.* — Drawing lancinations, or cutting, acute pains in them ; swelling of the knees ; corns on the feet, with burning pain as of excoriation.

## 9. CAMPHOR.

GENERAL SYMPTOMS. — Convulsions and cramps ; tetanus ; attacks of epilepsy, with rattling in the throat, red and puffed face, convulsive movements of the limbs and even

of the tongue, eyes, and muscles of the face, with hot and clammy perspiration on the hairy scalp and forehead ; after the fit, drowsy sleepiness ; sinking of all strength ; fainting fits ; the majority of the symptoms appear during motion, or else at night, or are aggravated by cold, the open air, and touch ; often disappear as soon as attention is called to them.

*Skin.* — Skin bluish and cold.

*Fever.* — Coldness over the whole body, with deadly paleness of face.

*Head.* — Congestion of the head ; inflammation of the brain ; spasms which draw the head to one side.

*Stomach.* — Vomiting and diarrhea, with failing strength ; sensation of burning and heat in the stomach.

*Abdominal Region.* — Cramps in the abdomen.

*Urine.* — Retention of urine.

*Genital Organs.* — Absence of sexual desire ; impotence ;

*Trunk.* — Cramps in the calves of the legs.

#### 10. CANTHARIS.

GENERAL SYMPTOMS. — Burning pains, as if from exco-riation, in all the cavities of the body ; the sufferings show themselves chiefly on the right side, and are mitigated in a recumbent posture ; erysipelatous inflammations.

*Moral Symptoms.* — Anxious inquietude, with agitation which forces one to keep constantly in motion.

*Teeth.* — Ulceration of the gums.

*Mouth.* — Aphthæ in the mouth.

*Throat.* — Sore throat, with burning in the throat in swallowing ; inflammation and ulceration of the tonsils and the throat, aggravated by drinking water.

*Urine.* — Urgent and ineffectual effort to make water, with painful emission, drop by drop ; burning smarting on making water ; sharp, tearing, and cutting pains, succes-



sive pullings and pulsations in the urinary organs ; inflammation and ulceration of the kidneys, of the vesica and of the urethra.

*Genital Organs.*—Draggings in the spermatic chord when making water ; inflammation and gangrene of the genital parts ; sexual desire greatly increased, with erections painful, frequent, and of long continuance.

*Larynx.*—Inflammation of the larynx.

*Chest.*—Shootings in the chest and in the sides ; burning pains in the chest.

### 11. CAPSICUM.

GENERAL SYMPTOMS.—Repugnance to movement ; the symptoms show themselves chiefly in the evening and at night, and are aggravated by the open air, contact and cold, as well as by beginning to move ; great sensibility to fresh air, and to a current of air.

*Fever.*—Cold and shivering every time after drinking.

*Head.*—Headache, as if the cranium were going to burst, when walking or moving the head ; attack of one-sided headache, pressive and shooting.

*Face.*—Redness of the face ; dull pressure on the cheek-bone.

*Throat.*—Burning blisters in the mouth and on the tongue ; sore throat, with painful deglutition and drawing in the pharynx.

*Larynx.*—Hoarseness ; cough more violent in the evening and at night, with pressure in the throat and ears.

*Trunk and Extremities.*—Acute drawing pains in the bones.

### 12. CARBO VEGETABILIS.

Evil effects from the abuse of mercury, or of cinchona ; sufferings caused by warm and damp weather ; sensibility

to changes of weather; intermittent fevers, even those which the abuse of cinchona has rendered obstinate; Asiatic cholera, with total absence of pulse.

**GENERAL SYMPTOMS.** — Burning pains in the limbs and in the back; the majority of symptoms appear while walking in the open air; soreness of all the limbs, especially in the morning, when one has just risen; sudden prostration of the physical powers; liability to take cold.

*Fever.* — Shivering and coldness in the body; cold sweat on the limbs and face.

*Moral Symptoms.* — Inquietude and anxiety, especially in the evening.

*Head.* — Heaviness of the head.

*Nose.* — Violent coryza, with hoarseness and raucity of the chest.

*Teeth.* — Opening, retraction, excoriation, and ulceration of the gums; bleeding of the teeth and gums.

*Mouth.* — Dryness, or accumulation of water in the mouth.

*Throat.* — Scraping and burning pain in the throat, the palate, and the gullet.

*Appetite.* — Salt taste in the mouth and of food; after a meal, great inflation of the abdomen.

*Stomach.* — Rising of fat food; sour risings, after a meal.

*Fæces.* — Evacuations liquid, pale, or mucous; involuntary evacuation.

*Urine.* — Diminution of the secretion of urine.

*Larynx.* — In the morning and in the evening, hoarseness, aggravated by prolonged conversation, and chiefly in cold and damp weather.

*Chest.* — Dyspnoea on walking; pain as from excoriation of the chest.

*Trunk.* — Rheumatic, drawing pains, acute pulling and

shootings in the back, the nape of the neck, and the muscles of the neck.

*Arms.*—Pullings and acute drawing pains in the forearms, the wrist, and the fingers: paralytic weakness of the wrists and of the fingers.

*Legs.*—Drawing and paralytic pain in the legs; cramps in the legs and in the soles of the feet.

### 13. CAUSTICUM.

Paralysis, especially that which manifests itself only on one side, or that which is the result of a suppression of some morbid secretion, or of some eruption, such as scabies, etc.

*Eyes.*—Sight confused, as if a gauze or mist were before it; darkness, which seems to dance before the eyes; lights and sparks before the eyes.

*Nose.*—Eruption on the point of the nose; fluent coryza with nocturnal cough, rough hoarseness, and headache.

*Throat.*—Accumulation of mucus in the throat, and behind the palate, with expectoration by hawking and gagging.

*Stomach.*—Aching, squeezing as if from claws, with constriction and cramp-like pains in the stomach and the epigastrium.

*Fæces.*—Chronic constipation; itching in the anus.

*Urine.*—Involuntary emission of urine, day and night, even when coughing, when sneezing, and when walking.

*Larynx.*—Rough hoarseness, morning and evening; cough dry, hollow, shaking, with sensation of burning and pain as from excoriation of the chest.

*Chest.*—Palpitation of the heart; oppression of the heart, with melancholy; stitches in the heart.

*Trunk.* Painful stiffness in the back, especially on rising from the chair.

*Arms.* — Contraction and induration of the tendons of the fingers.

*Legs.* — Tensive stiffness in the joints of the legs and of the feet.

#### 14. CIIAMOMILE.

Different affections of women and of children, chiefly lying-in women and new-born infants; bad effects from the abuse of coffee and of narcotic palliatives; suffering in consequence of a chill; affections arising from sudden grief or a fit of passion; convulsive and spasmodic attacks; excoriation of the skin; disposition for every wound to ulcerate; bilious and gastric affections; excoriation of the nipples; erysipelas on the breasts; catarrhal cough, with hoarseness, chiefly in children.

*GENERAL SYMPTOMS.* — Rheumatic, drawing pains, chiefly at night it bed; pains with thirst, heat and redness of one of the cheeks; over-excitement, and excessive sensibility of the nervous system, with great sensibility to pain.

*Sleep.* — Nocturnal sleeplessness; when sleeping, starts with fright, cries, tossing, tears, talking.

*Fever.* — Burning heat and redness (often only in one) of the cheeks, chiefly at night, with groaning, tossing.

*Moral Symptoms.* — Disposition to weep and to be angry, with great sensibility to offense; quarrelsome and choleric humor.

*Head.* — Headache on waking in the morning, or while asleep.

*Eyes.* — Eyes inflamed and red, with pressive pains, chiefly on moving the eyes and on shaking the head; blearedness in the eyes and nocturnal agglutination.

*Ears.* — Shootings extending into the ears, with disposition to be angry at trifles; buzzing in the ears; inflamma-

tory swelling of the parotids, as well as of the sub-maxillary glands and of those of the neck.

*Face*.—Face hot, red, burning, or redness and heat of one cheek, with coldness and paleness of the other.

*Teeth*.—Tooth-ache, mostly of one side, and chiefly at night, in the heat of the bed, with insupportable pains which almost drive one to despair, frequently after eating any thing hot (or cold), and chiefly after having taken coffee.

*Throat*.—Sore throat ; deep redness of the parts affected.

*Appetite*.—Bitter taste in the mouth and of food ; excessive thirst for cold drinks.

*Stomach*.—Bitter, bilious vomiting.

*Abdominal Region*.—Flatulent colic, with inflation of the abdomen ; excessively painful colic, pullings and cuttings in the abdomen.

*Anus and Fæces*.—Diarrhea, at night, with slimy or greenish fæces, or mucus.

*Urine*.—Menstrual colic, before the catemenia ; pressure toward the uterus, as if from the pains of child-birth ; metrorrhagia, with discharge of deep-red blood and of clots, and accompanied with labor-pains.

*Larynx*.—Dry cough, produced by a constant titillation in the larynx ; anger excites cough (in children.)

*Chest*.—Attacks of flatulent asthma, with anxiety and fullness in the pericordial region.

*Legs*.—Cramps in the calves of the legs, chiefly at night.

#### 15. CHINA. — CINCHONA.

GENERAL SYMPTOMS.—Pains and sufferings, excited or aggravated by touch, at night, or after a meal ; great general weakness, with trembling ; great tendency to perspiration when moving and sleeping.

*Skin*.—Yellow color of the skin.

*Sleep.*—Painful, frightful dreams, which continue to produce agitation after waking.

*Fever.*—Shiverings, with shuddering, or feverish trembling, commonly without thirst; the thirst generally takes place only before or after the shiverings; easy perspiration during sleep; nocturnal debilitating sweats.

*Moral Symptoms.*—Hypochondriacal dejection; excessive irascibility, with pusillanimity.

*Head.*—Headache as if from suppressed coryza; pain as from a bruise in the brain; pressive headache; acute jerking or pressive pains in the head; headache as if the head were going to burst; congestion to the head; sensibility of the exterior of the head, and even of the roots of the hair, when touched.

*Eyes.*—Sparkling, black, dancing spots, and obscuration before the eyes.

*Nose.*—Bleeding of the nose and of the mouth.

*Face.*—Complexion pale, earth-like, sometimes of a blackish yellow.

*Teeth.*—Dull and distressing pains in the carious teeth.

*Mouth.*—Tongue with a yellow or white coating.

*Appetite.*—Bitter taste of food and drink; desire for a variety of food, and for dainties, without knowing exactly which; great weakness of digestion.

*Stomach.*—Eructations, with taste of food; pressure in the stomach, with cramp-like pains, especially after having eaten.

*Abdominal Region.*—Hardness and swelling of the liver; swelling and hardness of the spleen; piercing in the spleen; dropsical swelling of the abdomen, with asthmatic sufferings and fatiguing cough.

*Fæces.*—Slimy, watery, yellowish diarrhea; loose evacuations, with excretion of undigested food; loose evacua-



tions, chiefly after a meal or at night ; crawling in the anus, as if from worms.

*Urine.*—Urine deep-colored, with sediment like brick-dust.

*Genital Organs.*—Congestion to the uterus ; leucorrhea, with cramp-like contraction.

*Larynx.*—Violent, convulsive cough, sometimes even with inclination to vomit ; cough, with expectoration streaked with blood.

*Chest.*—Shooting in the chest, when coughing and breathing ; stitches in the side ; violent congestion to the chest.

*Arms.*—Paralytic, jerking, tearings in the muscles and bones of the upper and lower extremities, excited by touch.

## 16. CINA.

Scrofulous affections ; acute hydrocephalus of children ; wetting the bed ; hooping-cough ; chiefly in scrofulous children or in those suffering from worms.

GENERAL SYMPTOMS.—Convulsions and distortion of the limbs ; epileptic convulsions, with cries, turning on the back, and violent movements of the hands and feet ; external pressure aggravates or renews the sufferings.

*Sleep.*—Nocturnal sleeplessness, with agitation, tears, cries, heat and anguish.

*Moral Symptoms.*—Child cries when it is touched.

*Eyes.*—Pupils dilated.

*Nose.*—Desire to put the fingers into the nose ; stoppage of the nose.

*Teeth.*—Grinding of the teeth.

*Appetite.*—Hunger a short time after a meal.

*Fæces.*—Loose evacuations, of the consistence of pap ; discharge of ascarides and of worms by the anus ; loose, involuntary, whitish evacuations.

*Urine.*—Wetting the bed ; urine soon becomes turbid.

*Larynx.*—Cough, with sudden starts and loss of consciousness.

*Extremities.*—Contraction and jerking of the hand and of the fingers ; cramp-like extension of the legs.

#### 17. COCCULUS.

GENERAL SYMPTOMS.—Semi-lateral sufferings ; aggravation of the sufferings from sleep, conversation, drinking and eating, especially from coffee or smoking tobacco.

*Head.*—Vertigo, as if from intoxication, or when rising up in the bed, with desire to vomit, which forces one to lie down again ; pain in the head, which is, as it were, empty and hollow.

*Appetite.*—Excessive loathing of all food and drink.

*Stomach.*—Attacks of nausea unto fainting ; vomiting and nausea from the motion of a carriage, or on the sea ; violent, cramp-like, and squeezing pains, as if from a claw, and cramps in the stomach, sometimes a short time after a meal ; pinchings in the epigastrium, with difficulty of respiration.

*Abdominal Region.*—Sensation in the abdomen, as if it were hollow and empty ; contractive pinchings in the lower part of the abdomen ; flatulent, cramp-like colic, especially at night, aggravated by coughing, or by stooping forward ; disposition to protrusion of inguinal hernia.

*Fæces and Urine.*—Hard and difficult evacuations ; frequent desire to make water, also in pregnant women.

*Genital Organs.*—Premature catamenia, with cramps in the abdomen.

*Chest.*—Cramps in the chest ; rattling, and sensation of emptiness in the chest.

*Arms.*—Paralysis of the arms and lower limbs.

## 18. COFFEA.

Excessive nervous excitability; excessively painful neuralgia; sleeplessness from nervous excitement; evil consequences of unexpected or excessive joy.

GENERAL SYMPTOMS.—Painful susceptibility of parts affected; mental and physical excitability; sleeplessness from excitement of the imagination, flow of ideas, and fantastic visions; desire to lie down and to shut the eyes, without being able to sleep.

*Head and Throat.*—Pains in the head, as if the brain were bruised; sore throat, with great and painful sensibility.

*Stomach and Fæces.*—Cramps in the stomach; abdominal pains, which even drive one to despair, especially in women; diarrhea, also, during teething.

## 19. COLCHICUM.

Rheumatic and arthritic affections; dropsical affections; dysentery.

GENERAL SYMPTOMS.—Rheumatic and arthritic tearing in the limbs, especially in warm weather; the sufferings are aggravated by intellectual fatigue, by touch, too brilliant a light, and from the commencement of the night till morning.

*Stomach.*—Nausea, even to syncope, from the smell of fresh eggs or fat meat; sensation of coldness or of burning in the stomach.

*Abdominal Region.*—Dropsical swelling of the abdomen.

*Fæces.*—Dysenteric diarrhea of white, transparent, gelatinous mucus; discharge of much slime from the rectum; prolapsus recti.

*Urine.*—Scanty discharge of urine of a deep color, with tenesmus and burning.

*Arms.*—Tearings in the upper and lower extremities.

*Legs.*—Hot swelling of the legs; tingling of the toes, as if they had been frozen.

## 20. COLOCYNTHIS.

Evil effects from mental emotions, with indignation and mortification.

**GENERAL SYMPTOMS.**—Painful cramps and cramp-like contractions, in the internal or external parts.

*Skin.*—Troublesome itching, with great restlessness in the whole body, especially in the evening in bed, followed by perspiration.

*Head.*—Attacks of semi-lateral headache.

*Stomach.*—Colic and diarrhea, however little is eaten.

*Abdominal Region.*—Inflammation of the abdomen, as if from tympanitis; cramp-like pain and constriction in the intestines, especially after a fit of passion; excessively violent colic, with cutting, cramp-like, or contractive pains, which compel one to bend double, with restlessness in the whole body, and with a sensation of shuddering in the face, which seems to proceed from the abdomen; coffee and tobacco-smoke diminish the colic.

*Feces.*—Loose evacuations of a greenish yellow; dysenterical evacuations, with colic.

*Urine.*—Diminished secretion of urine.

## 21. CONIUM.

Affections of old men; inveterate affections resulting from contusion, principally in the glands; tetters; scirrhous indurations and cancerous ulcers, especially when caused by contusion; cataract, caused by a blow; cancer in the lips; asthma, especially in old men.

**GENERAL SYMPTOMS.**—Pains, as from fatigue, in the limbs and joints, when in a state of repose; the symptoms

appear during repose, and by any movement ; sensation of fatigue, especially early in the morning in bed ; want of energy and nervous debility ; great fatigue and other sufferings, from walking in the open air.

*Skin.* — Nettle-rash, in consequence of violent bodily exercise ; brown or red itching spots over the whole body, which disappear and return.

*Fever.* — Slow fever, with total want of appetite ; pulse irregular ; nocturnal sweat.

*Moral Symptoms.* — Ill-humor and moroseness.

*Head.* — Vertigo on rising, on looking behind.

*Eyes.* — Obscuration of the sight ; black spots and colored bands before the eyes in a room.

*Face.* — Cancerous ulcer on the lip.

*Genital Organs.* — Swelling of the testes.

*Larynx.* — Dry, convulsive cough ; cough like whooping cough, with sanguineous expectoration, or in violent fits, during the night.

*Chest.* — Short respiration when walking ; difficulty of respiration, also in the morning on waking.

*Arms.* — Shoulders painful, as if they had been bruised and excoriated.

*Legs.* — Weariness of the knees ; coldness and strong disposition to chilliness of the feet.

## 22. CROCUS.

GENERAL SYMPTOMS. — Discharge of black, slimy blood, from different organs ; striking alternation of the most opposite physical and mental symptoms.

*Moral Symptoms.* — Strong tendency to sadness, sometimes alternately with great gaiety and joyous humour ; severity and mildness of character alternating.

*Throat.* — Sore throat, as if caused by elongation of the

palate, or as if there were a plug in the throat, during deglutition, and at other times.

*Abdominal Region.*—Movements in the abdomen, as if from something alive.

*Fæces.*—Metrorrhagia of black and slimy blood.

*Chest.*—Movements, as if something alive were in the chest.

*Arms.*—Chilblains in the hands, and on the toes.

### 23. CUPRUM.

Spasmodic affections and convulsions; encephalitis; Asiatic cholera; whooping-cough; spasmodic asthma.

GENERAL SYMPTOMS.—Tonic spasms, with loss of consciousness, throwing of the head backward; the convulsions generally begin in the fingers and in the toes; convulsive startings at night when sleeping; violent convulsions, with great display of strength; symptoms which appear periodically, and in groups.

*Head.*—Whirling vertigo, as if the head were going to fall forward; stupifying depression in the head, with crawlings in the vertex; pains in the occiput and nape of the neck, on moving the head.

*Eyes.*—Convulsions and restless movements of the eyes.

*Face.*—Spasmodic distortion of the face; lips bluish.

*Teeth and Mouth.*—Foam in the mouth.

*Stomach.*—Violent vomitings, with cramps in the abdomen, aggravated by touch and by movement.

*Abdominal Region.*—Spasmodic colic.

*Fæces.*—Violent diarrhea.

*Chest.*—Respiration accelerated; short, difficult respiration, with spasmodic cough and rattling in the chest; suffocating fits; cramps in the chest, which interrupt the respiration and the voice.

*Trunk and Arms.*—Cramps of the fingers and of the toes.



## 24. DROSERA.

Catarrh and hoarseness; hooping-cough; affections of the respiratory organs, in consequence of croup.

*Larynx.* — Crawling in the larynx, which excites a short cough and shootings as far as the throat; sensation of dryness or roughness, and of scraping in the bottom of the gullet, with inclination to cough; hoarseness and very low voice; dry, spasmodic cough, with inclination to vomit; fatiguing cough, like hooping-cough, with bluish face, wheezing respiration, attacks of suffocation, bleeding from the nose and mouth, and anxiety; vomiting of food during the cough and afterward.

*Chest.* — Difficulty of respiration when coughing or speaking.

## 25. DULCAMARA.

Affections, in consequence of taking cold in general; tetters of different kinds; nettle-rash.

GENERAL SYMPTOMS. — Aggravation of sufferings, chiefly in the evening or at night, and during repose mitigated by movement; swelling and induration of the glands; dropsical swelling of the whole body, limbs, and face.

*Skin.* — Dry furfuraceous, humid, scaling, or suppurating tetters; redish tetters, bleeding after being scratched, with painful sensibility to the touch, and to cold water.

*Fæces.* — Diarrhea, as if after a chill, of greenish or brownish mucus; nocturnal, watery diarrhea with colic.

*Larynx.* — Catarrh and hoarseness, as if from having taken cold; moist cough.

*Arms.* — Tettery eruption on the hands.

## 26. EUPHRASIA.

Catarrhal ophthalmia.

*Eyes.*—Inflammation of the eyelids, with headache ; abundant flow of corrosive tears.

*Nose.*—Violent flowing of coryza, with abundant secretion of mucus, excessive confusion of the head, and corrosive tears in the eyes.

*Chest.*—Cough in the morning, with flowing coryza, and abundant expectoration of mucus.

## 27. FERRUM.

Congestion of blood and hemorrhagia ; diarrhea, especially in children ; ascarides.

GENERAL SYMPTOMS.—Dropsical swellings, with shooting pains ; the symptoms manifest themselves at night, are aggravated by being seated, and mitigated by gentle movement ; great lassitude and general weakness.

*Sleep.*—Excessive and drowsy fatigue, with agitated sleep at night.

*Moral Symptoms.*—Anxiety, with beatings in the epigastrium ; gayety, alternating with sadness.

*Head.*—Periodical hammering, and pulsative headache ; congestion of the head.

*Face and Throat.*—Fiery redness of the face ; lips pale ; spitting of blood.

*Stomach.*—Vomiting of food, especially at night, or immediately after a meal ; pressure in the stomach, especially after eating meat.

*Fæces.*—Aqueous and acrid diarrhea ; undigested fæces.

*Genital Parts.*—Metrorrhagia ; abortion ; milk-like and acrid leucorrhea.

*Larynx.*—Spasmodic cough, ceasing immediately after a meal ; cough after a meal, with vomiting of food.

*Chest.*—Difficulty of respiration ; palpitation of the heart.

## 28. GRAPHITES.

Tetters of several kinds especially on the face ; wens. ;

GENERAL SYMPTOMS.—Great disposition to take cold, and fear of the open air and of currents of air.

*Skin.*—Tetters, and other humid or scabby eruptions, sometimes with secretion of corrosive serum, or with itching in the evening and at night; unhealthy skin, every injury tends to ulceration; deformity and thickness of the nails.

*Eyes.*—Inflammation of the eyes, injection of the veins, swelling and abundant mucous secretion from the eyelids.

*Ears.*—Dryness of the internal ear; scabs, tetters, running and excoriation behind the ears; hardness of hearing mitigated by the motion of a carriage.

*Nose.*—Dry scabs on the nose; nostrils excoriated, cracked, and ulcerated; fetid smell from the nose.

*Face.*—Flushes of heat in the face; erysipelatous inflammation and swelling of the face, with eruption of vesicles; one-sided paralysis and distortion of the muscles of the face, with difficult articulation; ulcers on the internal surface of the lips; scabby eruption on the chin and around the mouth.

*Throat.*—Sore throat, even at night, as if there were a plug within it.

*Appetite.*—Weakness of digestion; inflation of the abdomen after a meal.

*Abdominal Region.*—Immoderate expulsion of fetid wind, preceded by pinchings.

*Fæces.*—Obstinate constipation, with hard fæces; large hemorrhoidal excrescences in the anus.

*Urine.*—Wetting the bed.

*Menses.*—Suppression of menses; in place of menses, flow of blood from the anus.

*Arms.*—Arthritic nodosities on the fingers; tettery excoriation between the fingers; thickness of the nails of the fingers.

*Legs.*—Tetters on the thighs, hams, and tibia ; cold feet, even in the evening in bed ; tettery excoriation between the toes ; thickness and deformity of the toe-nails.

## 29. HELLEBORUS.

Dropsical affections, especially some kinds of anasarca, and chiefly those which proceed from the repercussion of eruptions, such as measles ; scald-head, with obstruction of the glands of the neck ; acute hydrocephalus.

GENERAL SYMPTOMS.—Convulsions.

*Sleep.*—Sleepiness, with eyes half open and pupils turned upward.

*Moral Symptoms.*—Dullness of the internal senses.

*Head.*—Stupefying pain and sensation of a bruise in the head ; painful heaviness of the head ; disposition to bury the head in the pillow, when sleeping.

*Fæces.*—Watery and frequent evacuations.

*Chest.*—Difficult respiration, as if from hydrothorax ; constriction of the chest.

## 30. HEPAR SULPHURIS.

Evil consequences of the abuse of mercury ; scrofulous affections ; icterus ; scald-head ; eruptions and tetters on the face ; scrofulous catarrhal ophthalmia ; ulcers on the cornea.

GENERAL SYMPTOMS.—Pains as from excoriation or a bruise on different parts when touched ; swelling, inflammation, and ulceration of the glands ; aggravation of the pains at night.

*Skin.*—Eruption of pimples and tubercles, painful when touched ; unhealthy skin, every injury tends to ulceration ; suppurations.

*Sleep.*—Jerking at night, as if from want of air.

*Fever.* — Dry heat at night; great disposition to perspire in the day-time.

*Head.* — Pain in the head, as if a nail were driven into it; pain, as from ulceration in the head, directly above the eyes; tuberosities on the head, with pain as from excoriation when touched.

*Eyes.* — Stinging in the eyes; inflammation of the eyes and eyelids, with pain as from a bruise and excoriation when touched; specks and ulcers on the cornea; spasmodic closing of the eyelids.

*Ears.* — Heat, redness, and itching in the ears; scabs behind and on the ears.

*Nose.* — Pain as from a bruise and excoriation in the nose when touched.

*Face.* — Face burning and of a deep red; erysipelatous inflammation and swelling of the face; pains in the bones of the face when touched.

*Teeth.* — Toothache, with jerking and drawing pains, aggravated by closing the teeth, by eating, and in a hot room; swelling and inflammation of the gums, which are painful to the touch.

*Mouth.* — Salivation; sore throat, as if there were a plug in it.

*Appetite.* — Desire only for acid or pungent things.

*Stomach.* — Eructations, with burning sensation in the throat.

*Faces.* — Whitish diarrhea, of an acid odour, especially in children; dysenteric evacuations, greenish, or of a clay-color, with evacuation of sanguinous mucus.

*Urine.* — Wetting the bed; discharge of mucus from the urethra.

*Genital Organs.* — Flow of prostatic fluid, especially after making water, and during a difficult evacuation; excoriation between the thighs; leucorrhea, with smarting.

*Larynx.* — Hoarseness ; dry cough in the evening, from any part of the body becoming cold, or when lying on the bed ; attacks of dry, rough, and hollow cough, with anguish and suffocation, often causing one to weep.

*Chest.* — Anxious, hoarse, wheezing respiration, with danger of suffocation when lying down.

*Arms.* — Pain, as from a bruise, in the bones of the arm ; arthritic swelling of the hand, fingers, and joints of the fingers, with heat, redness and pain, as from dislocation, during movement ; skin of the hands cracked, rough, and dry ; panaris.

*Legs.* — Swelling of the knees ; cracks in the feet.

### 31. HYOSCYAMUS.

Convulsions and other spasmodic affections, chiefly in pregnant or parturient women, as well as in children, and in consequence of worms ; excessive nervous excitement, with sleeplessness : encephalitis ; acute hydrocephalus.

*GENERAL SYMPTOMS.* — Convulsions, with cries, great anguish.

*Fever.* — Burning heat of the body, especially of the head.

*Moral Symptoms.* — Desire to run away ; loss of consciousness, with eyes closed, and raving about business ; delirium ; perversion of all actions.

*Head.* — Headache, as if from concussion of the brain ; pressive and benumbing pain in the forehead.

*Eyes.* — Eyes red, fixed, convulsed, spasmodic closing of the eyelids ; nocturnal blindness ; weakness of sight, as if from incipient amaurosis.

*Mouth and Throat.* — Redness of the tongue ; constriction in the throat and inability to swallow liquids.

*Appetite and Stomach.* — Vomiting of food and drinks immediately after a meal.



*Fæces*.—Involuntary evacuations.

*Genital Organs*.—Metrorrhagia, of a bright, colored blood.

*Larynx*.—Cramp-like cough at night, especially when lying down, sometimes with redness of the face and vomiting of mucus.

*Chest*.—Pressure in the right side of the chest, with great anxiety and shortness of breath when ascending stairs ; spasms in the chest, with short breathing.

*Arms*.—Hands clenched, with retraction of the thumbs (in the convulsive fits.)

*Legs*.—Painful cramps in the thighs and calves of the legs.

### 32. IGNATIA.

Spasmodic affections, especially in consequence of fright or contradiction, and chiefly in hysterical women, or in children ; melancholy and other mental affections caused by affliction ; difficult teething of children, with convulsions ; prolapsus recti, also in children.

GENERAL SYMPTOMS.—Violent pain, merely on being touched, in different parts ; attacks of cramp and of convulsions ; hysterical debility and fainting fits ; the pains are removed always by change of position.

*Skin*.—Excoriation of the skin ; itching when becoming warm in the open air.

*Sleep*.—Violent spasmodic yawnings, especially in the morning.

*Fever*.—Absence of thirst during the heat and perspiration.

*Moral Symptoms*.—Tenderness of character, and delicacy of conscience ; love of solitude.

*Head*.—Pressive headache, especially above the root of the nose, aggravated or relieved by stooping ; the head-

aches are aggravated by coffee, brandy, tobacco-smoke, noise, and strong smell; headache as if a nail were driven into the brain.

*Eyes.* — Convulsive movements of the eyes and eyelids.

*Face.* — Convulsive jerkings, and distortion of the muscles of the face; convulsive twitching of the corners of the mouth.

*Throat.* — Sore throat, as if there were a plug in it; shootings in the throat, extending sometimes to the ear, chiefly when not swallowing; inflammation, swelling, and induration of the tonsils, with small ulcers; impeded deglutition (of drinks.)

*Appetite.* — Dislike to milk and tobacco-smoke; painful inflation of the abdomen after a meal.

*Stomach.* — Hiccough every time after eating and drinking; periodical attacks of cramp in the stomach, aggravated by pressing on the part affected.

*Abdomen.* — Shootings and pinchings in the abdomen, especially in the sides; flatulent colic, especially at night.

*Fæces.* — Hard evacuations, with frequent ineffectual efforts; prolapsus of the rectum during evacuation; itching and crawling in the anus.

*Catamenia.* — Cramp-like and compressive pains in the region of the uterus, with attacks of choking.

*Larynx.* — Short cough, as if from a feather in the throat, becoming more violent the more one coughs.

*Chest.* — Difficult respiration, as if hindered by a weight upon the chest; choking when running; palpitation of the heart at night.

*Arms.* — Insupportable pains in the bones and joints of the arms, as if the flesh were being loosened; convulsive jerkings in the arms and in the fingers.

*Legs.* — Convulsive jerkings of the legs.

## 33. IODIUM.

Scrofulous and lymphatic affections ; abdominal phthisis ; inflammatory swelling of the knee.

GENERAL SYMPTOMS. — Swelling and induration of the glands ; atrophy and emaciation until reduced to a skeleton.

*Fever.* — Pulse quick, small, and hard.

*Head.* — Congestion to the head, with beating in the brain.

*Face.* — Complexion pale, yellowish ; swelling of the submaxillary glands.

*Teeth.* — Softening of the gums.

*Stomach.* — Eructations, generally acid, with burning sensation.

*Abdominal Region.* — Abdominal pains, which return after every meal ; enlargement of the abdomen, which renders it impossible to lie down without the danger of suffocation.

*Fæces.* — Loose, soft evacuations, alternating with constipation ; violent, frothy diarrhea.

*Catamenia.* — Catamenia, at one time too late, at another too soon.

*Larynx.* — Inflammation of the throat and trachea, with contractive sore pain ; copious secretion of mucus in the trachea, with frequent hawking ; dry cough ; cough in the morning.

*Chest.* — Difficulty of respiration when going up stairs.

*Trunk.* — Swelling of the exterior of the neck ; swelling of the glands of the neck, nape, and armpits ; hard and large goitre.

## 34. IPECACUANHA.

Gastric and bilious fevers ; intermittent fevers ; gastric uneasiness, especially when caused by indigestion ; Asiatic

and sporadic cholera ; gastric affections, with vomiting and diarrhea ; asthmatic affections.

**GENERAL SYMPTOMS.**— Attacks of uneasiness, with dislike to all food, and excessive and sudden debility.

*Fever.*— Coldness, especially of the hands and feet ; thirst only during the shivering or chill.

*Head.*— Attacks of headache, with nausea and vomiting.

*Appetite.*— Great repugnance and dislike to all food.

*Stomach.*— Nausea ; vomiting of drink and undigested food, of bilious, greenish, or acid matter, and sometimes immediately after a meal ; vomiting, with diarrhea.

*Feces.*— Loose evacuations like matter in a state of fermentation ; diarrhea, with nausea, colic and vomiting ; dysenteric evacuations, with white flocks, and followed by tenesmus.

*Genital Organs.*— Metrorrhagia, with discharge of bright red and coagulated blood.

*Larynx.*— Cough, especially at night, with painful shocks in the head and stomach, with disgust and inclination to vomit and vomiting, or with fits of suffocation, stiffness of the body, and bluish face.

*Chest.*— Spasmodic asthma, with contraction of the larynx.

### 35. KALI CARBONICUM.

Ascites ; tuberculous phthisis ; dry cough at night.

*Appetite.*— After a meal, desire to sleep, paleness of the face, shivering, headache, ill-humor, nausea, sour eructations and pyrosis, colic, inflation of the abdomen and flatulency ; hoarseness and roughness in the throat.

### 36. KALI HYDRIODICUM.

*Face — Appetite.*— Toothache, aggravated by contact with any cold, mitigated by hot things ; gums swollen with lancinating pain, as from ulceration.

*Larynx—Extremities.*—Dry cough, excited by constant irritation in the trachea, with hoarseness.

### 37. LACHESIS.

Sufferings of drunkards, from the abuse of mercury; fainting fits; erysipelas.

*GENERAL SYMPTOMS.*—Great weakness of body and mind.

*Skin.*—Skin yellow, green, lead-colored, or bluish, or blackish, chiefly round the wounds and ulcers.

*Head.*—Vertigo, chiefly on waking in the morning; apoplectic fits, with blue face, convulsive movements of the limbs; headache, with congestion of blood, sparkling before the eyes; headache, every morning on waking, or after dinner, or else from every change of weather.

*Mouth.*—Tongue shining, red, and cracked; painful excoriation and inflammatory swelling of the throat; constant desire to swallow, and a sensation on swallowing as if there were a tumor, some foreign body, or a plug in the throat; ulcers on the palate, back of the mouth, and throat.

*Appetite.*—Desire for wine.

*Stomach.*—Excessive sensibility of the precordial region to the slightest touch.

*Abdominal Region.*—Inflammation and softening of the liver; pain and stitches in the region of the spleen; abdomen hard and distended, with flatulent colic.

*Fæces.*—Obstinate constipation, with hard and difficult evacuation; loose evacuations, principally at night, or after a meal, or in warm (and damp) weather, or from having taken fruits and acids.

### 38. LYCOPodium.

Obstruction of the glands; typhus fever; inertia of the intestines and obstinate constipation; chronic pneumonia; tuberculous phthisis.

GENERAL SYMPTOMS. — Excessive sensibility to fresh air ; great tendency to take cold.

*Skin.* — Excoriation of the skin of children.

*Moral Symptoms.* — Melancholy and disposition to weep ; aversion to speaking.

*Face.* — Frequent flushes of transient heat in the face ; swelling of the submaxillary glands.

*Mouth.* — Dryness of the mouth without thirst.

*Throat.* — Dryness of the throat ; inflammation of the throat and palate, with shooting pain.

*Stomach.* — Pressure in the stomach after every meal ; swelling of the epigastrium, with painful sensibility to the touch.

*Abdominal Region.* — Tension round the hypochondria, as if caused by a hoop ; induration of the liver ; fullness and distension of the stomach and abdomen ; obstructed flatulency.

*Feces.* — Obstruction of the abdomen.

*Genital Organs.* — Leucorrhea, milky, yellowish, redish and corrosive ; excoriation and running sores of the nipples.

*Larynx.* — Nocturnal cough, which affects the head, diaphragm, and stomach.

*Chest.* — Short respiration during almost every exertion ; palpitation of the heart, especially during digestion ; painful eruption and liver spots on the chest.

*Trunk.* — Swelling of the glands of the neck and shoulder, with shooting pain.

*Arms.* — Nocturnal aching pains in the arms ; dryness of the skin of the hands.

*Legs.* — Swelling of the knees ; ulcers in the legs ; pain in the soles of the feet when walking.

### 39. MERCURIUS VIVUS.

Swelling and inflammation of the glands ; inflammatory



fevers, with disposition to perspire profusely ; rheumatic or catarrhal headache ; scrofulous, rheumatic, catarrhal (and arthritic) ophthalmia ; syphilitic ophthalmia ; rheumatic and catarrhal otalgia ; rheumatic prosopalgia and toothache ; dysentery ; mucous or bilious diarrhea ; influenza.

GENERAL SYMPTOMS. — Rending and drawing, or stinging pains in the limbs, principally at night, in the heat of the bed, which renders the pain insupportable ; nocturnal aching pains ; rheumatic pains, with profuse sweat, which affords no relief ; the whole body feels as if it had been bruised, with soreness in all the bones ; emaciation and atrophy of the whole body ; excitability and sensibility of all the organs.

*Skin.*—Enlargement, inflammation, and ulceration of the glands, with pulsative and shooting pains, hard, red, and shining swelling ; violent and voluptuous itching over the whole body, principally in the evening, or at night, augmented by the heat of the bed, and sometimes with burning after being scratched.

*Fever.*—Copious, excessive, and colliquative sweats, both day and night, in the morning ; sweat, with nausea and desire to vomit ; great fatigue.

*Head.*—Vertigo, principally on getting up, or on raising up the head ; fullness and pressure in the head, as if the forehead were squeezed by a band, or that the cranium would split ; heat and burning, or tearing and drawing pains, or stinging in the head, often only one-sided and extending to the ears, teeth, and neck.

*Eyes.*—Itching, tickling, and burning in the eyes ; eyes red and inflamed ; eyelid red, inflamed, swollen, ulcerated on the margins, and covered with scabs.

*Ears.*—Tearing, stinging, and drawing pains in the ears, increased by the heat of the bed ; purulent otorrhea, with tearing in the affected side of the head and in the face.

*Face.*—Bloatedness and swelling of the face ; tearing in the bones and muscles (of one side) of the face ; obstruction and inflammatory swelling of the submaxillary glands, with stinging or pulsative pains.

*Teeth.*—Tearing, stinging, or pulsative pains in the carious teeth, or in the roots of the teeth, often spreading as far as the ears, and in the entire cheek of the side affected, sometimes, also, with painful swelling of the cheek or of the submaxillary glands, with salivation and shivering ; appearance or aggravation of toothache, principally in the evening or at night, in the heat of the bed, where it is insupportable ; renewed by the fresh air, as well as by eating, and taking any thing hot or cold into the mouth ; retraction and swelling of the gums, principally at night, with burning pain and sensation of excoriation on touching them and on eating ; gums livid, discolored, and very sensitive ; ulceration of the gums.

*Mouth.*—Putrid smell from the mouth ; inflammatory swelling of the inside of the mouth ; burning pain, vesicles, blisters, aphthæ, and ulcers in the mouth ; accumulation of tenacious mucus ; profuse discharge of excessively fetid saliva ; tongue moist, coated with white and thick ; inflammatory swelling and ulceration of the tongue, with stinging pains ; entire loss of speech.

*Throat.*—Stinging pains in the throat and tonsils, principally when swallowing ; inflammatory swelling and redness of all the back parts of the mouth and throat ; constant desire to swallow ; inability to swallow the least liquid, which escapes through the nostrils ; the pains in the throat commonly extend as far as the ears, the parotides, the submaxillary glands and those of the neck : they are aggravated, for the most part, by empty deglutition, as well as at night, in the fresh air and when speaking, and they are accompanied with salivation.

*Appetite.*— Acid and mucous taste ; dislike to all food, principally solid nutriment, meat.

*Stomach.*— Violent, empty eructations ; excessive tenderness of the stomach and precordial region ; pressure as if from a stone in the pit of the stomach.

*Abdominal Region.*— Painful sensibility of the hepatic region, with stinging, burning pains ; complete icterus ; obstruction and inflammatory swelling of the inguinal glands.

*Fæces.*— Loose and dysenteric evacuations, principally at night, with colic and cuttings ; tenesmus and burning in the anus ; nausea and eructations ; shivering and shuddering, exhaustion and tremor of all the limbs ; scanty evacuations of sanguinolent mucus ; evacuations which are mucous, or bilious, or putrid, or acid, or of a greenish or brownish color ; evacuation of acrid and burning fæcal matter ; discharge of blood or of mucus from the rectum ; discharge of ascarides and lumbrici.

*Urine.*— Frequent, copious emission of urine, like diabetes ; corrosive and burning urine.

*Larynx.*— Catarrh, with febrile shivering ; continued hoarseness and loss of voice ; pains in the head and chest when coughing, as if these parts were about to burst.

*Chest.*— Difficulty of respiration, with attacks of suffocation at night, or in bed, in the evening when lying (on the left side) ; shootings in the chest and side, or extending as far as the back, principally when breathing, sneezing, and coughing ; pain as from excoriation and of ulceration in the chest.

*Trunk.*— Obstructions and inflammatory swelling of the glands of the neck.

*Arms.*— Sharp pains in the shoulders and arms, principally at night and when moving these parts.

*Legs.*— Sharp and piercing pains in the hip-joints, as

well as the thighs, principally at night, and when moving ; cedematous, transparent swelling of the thighs and legs.

#### 40. NATRUM MURIATICUM.

Intermittent fevers, also those which have changed their character from strong doses of cinchona ; chronic constipation.

*Fever.* — Continual shivering and want of vital heat ; fever, with aching pains ; yellowish complexion, headache, weakness, bitter taste in the mouth, ulceration at the angles of the mouth, absence of appetite ; the fever generally commences in the morning, followed by heat and thirsts ; typhus fever, with debility, dryness of tongue, and violent thirst ; pulse irregular and often intermittent.

*Moral Symptoms.* — Irascibility ; easily provoked.

*Head.* — Headache in the morning, on turning and while moving the body or the head, when running or in the cold air, as if the head were to split, with nausea and vomiting ; abundant falling out of the hair.

*Eyes.* — Inflammation of the eyes ; the letters are confused when reading ; the sight is weak, as if from incipient amaurosis ; black specks, luminous marks, and sparks before the eyes.

*Ears.* — Hardness of hearing ; humming in the ears.

*Nose.* — Loss of smell ; dryness of the nose.

*Face.* — Lips dry, chapped, cracked, or excoriated and ulcerated, with scurf, and burning and smarting eruption ; tettery eruption round the mouth ; frequent swelling of the submaxillary glands.

*Teeth.* — Piercings, boring, and pulsation in carious teeth ; fistula of the gums.

*Mouth.* — Ulcers and vesicles on the tongue and in the mouth, with burning, smarting, and pain, on coming in contact with food and drink.

*Throat.*—A sensation, during deglutition, as if there were a plug in the throat; inflammation of the throat, with stinging pain and ulceration.

*Fæces.*—Constipation; burning in the anus and rectum, during and after the evacuations; painful and burning hæmorrhoidal tumors about the anus.

*Genital Organs.*—Menses too early and too profuse, or retarded and scanty.

*Larynx.*—Cough excited by a tickling in the throat, or in the epigastrium, day and night; pains in the head when coughing, as if the forehead were about to burst.

*Chest.*—Short breathing, especially when walking quickly; anxious and violent palpitation of the heart from every movement of the body, but principally when lying on the left side; irregular and intermittent palpitation of the heart.

#### 41. NITRI ACIDUM.

Aching pains, inflammation, caries, and other affections of the bones, especially from the abuse of mercury; syphilitic affections; mercurial ulcers; nervous excitability, especially from the abuse of mercury; syphilitic and mercurial angina, also with ulceration; chronic relaxation of the bowels.

GENERAL SYMPTOMS.—Stinging pains as if caused by splinters, especially when touched; pains from change of weather; riding in a carriage mitigates the majority of the symptoms.

*Face.*—Painful swelling of the submaxillary glands.

*Mouth.*—Offensive and putrid smell of the mouth; salivation, sometimes from attacks of fever.

*Throat.*—Ulcer with stinging pain in the mouth and throat; inflammation of the throat, with stinging pains, especially on swallowing liquids, as from excoriation.

*Fæces.*— Loose evacuations, sometimes mucous or putrid ; sanguinolent, dysenteric evacuations, with tenesmus ; swelling of hemorrhoidal tumors about the anus, and which bleed at every evacuation.

*Genital Organs.*— Ulcers, like chancres, on the prepuce and glands.

*Larynx.*— Dry cough, especially in the evening on lying down ; purulent yellowish expectoration with the cough.

*Chest.*— Short breath.

*Trunk.*— Stinging in and between the shoulder-blades ; swelling of the glands of the neck and axillæ ; suppuration of the axillary glands.

*Legs.*— Chilblains on the toes.

#### 42. NUX VOMICA.

Sufferings from the abuse of coffee, wine, or other spiritous or narcotic drugs ; bad effects from passion or excessive study, prolonged watching, or a sedentary life ; periodical and intermittent affections ; gastralgia ; gastritis ; gastrico-mucous or bilious affections ; dyspepsia, also with vomiting of food ; vomiting of drunkards, of pregnant women ; incarcerated hernia ; obstinate constipation ; blind and bleeding hemorrhoids.

GENERAL SYMPTOMS.— Repugnance to the open air ; great desire to remain lying down or sitting.

*Sleep.*— Too short sleep, with difficulty in going to sleep again after midnight, and inability to remain in bed after three o'clock in the morning ; on waking in the morning, pain in the limbs as if they were bruised, great lassitude, with desire to remain lying down, and fits of stretching and of convulsive yawning.

*Fever.*— During the shivering, skin, hands and feet, face or nails are cold and bluish ; during the heat, vertigo,



headache, shivering when moving in the least, or when uncovered in the slightest degree, thirst.

*Moral Symptoms.*—Hypochondriacal, sorrowful, and sad humor; ill-humor, vexation, and anger; dislike to, and unfitness for, bodily and mental labor.

*Head.*—Vertigo, with sensation of turning and of wavering of the brain; heaviness, pressure, and sensation of expansion in the head, as if the forehead were about to burst, especially above the eyes.

*Eyes.*—Eyes inflamed, with redness and swelling, also the eyelids.

*Nose.*—Obstruction of the nose, sometimes on one side only, and often with itching in the nostrils and discharge of mucus; obstruction in the head, principally in the morning, or at night, and dry coryza with heat and heaviness in the forehead, and obstruction of the nostrils.

*Teeth.*—Putrid and painful swelling of the gums.

*Mouth.*—Fetid, putrid, and cadaverous smell from the mouth; tongue loaded with a white coating, or dry, cracked, brownish or blackish.

*Throat.*—Swelling of the uvula; bitter taste of the mouth; desire for brandy.

*Stomach.*—Frequent, bitter, and acid eructations; violent hiccough; vomituration and violent vomiting of mucous and sour matter, after drinking or eating, or in the morning; pressure in the stomach as if by a stone; tension and fullness in the epigastrium; tight clothes are insupportable.

*Abdominal Region.*—Flatulent colic, sometimes in the morning, after eating or drinking, with pressing pains, as if by stones.

*Fæces.*—Frequent but ineffectual and anxious desire to evacuate; obstinate constipation, as if from inactivity of the intestines; incomplete evacuations; small, loose, aque-

ous or mucous and sanguinolent evacuations ; hemorrhoides, with excoriating, stinging, burning pain, and pressure in the anus and rectum.

*Urine.* — Ineffectual desire to urinate ; frequent emission of watery and pale urine ; burning pain in the neck of the bladder when making water.

*Larynx.* — Catarrhal hoarseness and painful roughness of the larynx and chest ; accumulation of tenacious mucus, which it is impossible to detach ; pains, as from excoriation in the larynx when coughing.

*Chest.* — Asthmatic constriction of the chest at night, in bed, when going up stairs, choking, anxiety.

*Trunk.* — Pains, like those caused by a bruise in the back and loins ; rheumatic, drawing, and burning pains in the back.

#### 43. OPIUM.

Recent affections rather than those of long standing ; nervous torpor and want of vital reaction against the medicines that have been administered ; sufferings of drunkards ; affections of old men ; bad effects of fright, with continued fear, or of sudden joy ; typhus ; delirium tremens ; ileus ; constipation, principally that caused by torpor of the intestinal canal, after frequent diarrhea, or from want of exercise, and especially in the case of vigorous persons, or those who are plethoric or well fed, as well as in the case of children and pregnant women ; tympanitis ; suppressed or false and spasmodic labor-pains.

GENERAL SYMPTOMS. — General insensibility of the whole nervous system.

*Sleep.* — Lethargy, with snoring and mouth open ; excessive desire to sleep, with absolute inability to go to sleep.

*Fever.* — Pulse generally full, slow ; fever, with lethargic sleep, snoring.

*Face.*—Face dark red, sometimes brownish, hot, and bloated.

*Stomach.*—Vomiting of fæcal matter and of urine.

*Abdominal Region.*—Abdomen hard and distended, as in tympanitis.

*Fæces.*—Constipation, long continued; involuntary evacuations.

*Chest.*—Noisy, stertorous, and rattling respiration; attacks of suffocation on making an effort to cough.

#### 44. PETROLEUM.

Gastric affections; sea sickness; sufferings caused by the motion of a carriage; chilblains on the hands and feet.

*Appetite.*—Repugnance to meat and fat; heaviness and pressure in the stomach after a meal.

*Stomach.*—Nausea from the motion of a carriage.

#### 45. PHOSPHORUS.

Physical and nervous weakness caused by protracted influences injurious to the vital economy; hemorrhage and congestion of blood; cholera; chronic and colliquative diarrhea; chronic laryngitis; disposition to croup.

GENERAL SYMPTOMS.—The majority of the symptoms manifest themselves morning and evening, in bed, as well as after dinner, while several others appear at the beginning of a meal and disappear after it.

*Skin.*—Lymphatic abscesses, with fistulous ulcers, which have callous margins, and secrete a fetid and colorless pus, with hectic fever; copious bleeding from small wounds.

*Sleep.*—Unrefreshing sleep; in the morning it appears as if one had not slept enough; nocturnal heat; hectic fever, with dry heat toward evening, especially in the palms

of the hands, sweat and colliquative diarrhea, circumscribed redness of the cheeks, etc.

*Moral Symptoms.* — Anguish and uneasiness, especially when alone.

*Head.* — Vertigo, with nausea and pressing pains in the head; congestion to the head, with beating; falling off of the hair.

*Eyes.* — Inflammation of the eyes; nocturnal agglutination of the eyes; black spots before the sight.

*Nose.* — Unpleasant dryness of the nose; continued discharge of yellow mucus from the nose.

*Face.* — Face pale, wan, dirty, earthy, with hollow eyes, surrounded by a blue circle.

*Stomach.* — Sour regurgitations of food; nausea of various kinds, especially in the morning; spasmodic pain and contraction in the stomach.

*Abdominal Region.* — Spasmodic colic; sensation of coldness, with heat and burning in the abdomen; flatulent colic, with grumbling.

*Faces.* — Prolonged looseness of the bowels.

*Larynx.* — Aphonia; cough excited by a tickling and itching in the chest, or with hoarseness and sensation as if the chest were raw; cough, with purulent and saltish expectoration, or of blood.

*Chest.* — Obstructed respiration of various kinds; congestion to the chest; palpitation of the heart.

*Arms.* — Trembling in the arms and hands.

*Legs.* — Drawing and tearing in the knees; swelling of the feet.

#### 46. PHOSPHORIACIDUM.

Phthysical and nervous weakness, caused by violent acute diseases, by loss of the fluids and other debilitating

causes, especially when they have rapidly undermined a constitution that was previously strong; bad effects from vexation, with care and inquietude, or from disappointed love; epidemic diarrhea.

*Fever.*—Malignant fever with great weakness, apathy, stupidity, aversion to conversation, diarrhea.

*Fæces.*—Purging, but not debilitating evacuations; loose, serous, or undigested evacuation.

*Urine.*—Frequent and profuse emission of aqueous urine, which immediately deposits a thick and white sediment; urine as in diabetes mellitus.

*Larynx.*—Cough, with vomiting of food, and headache.

#### 47. PLATINA.

Different affections of females, and principally of irritable women; hysteria; facial neuralgia; metrorrhagia, after accouchment, or caused by miscarriage; constipation of lying-in women.

GENERAL SYMPTOMS.—Affections caused by fright, by vexation, or by a fit of passion; the majority of the symptoms are aggravated by repose.

*Moral Symptoms.*—Sadness, with great inclination to weep; involuntary weeping, with great fear of death, which is believed to be very near; pride and self-conceit, with contempt for all others.

*Face.*—Burning heat and glowing redness in the face; torpor in the whole side of the face; cramp and tensive pressure in the zygomatic process.

*Teeth.*—Toothache, with pulsative and digging pain.

*Fæces.*—Constipation, sometimes very obstinate.

*Urine and Genital Organs.*—Sensation of bearing down toward the genital organs, with pressure in the abdomen; metrorrhagia of thick, deep colored blood, with drawings in the groins.

*Chest.* Short breath, with constrictive oppression of the chest; anxious palpitation of the heart.

#### 48. PULSATILLA.

Affections of persons of mild character, inclined to pleasantry, and to laughter or weeping, with a mild countenance, and of phlegmatic temperament, inclining to melancholy, lymphatic constitution, with pale complexion, blue eyes, and light hair, freckles, disposition to take a cold in the head, or to other mucous discharges, etc.; bad consequences from the abuse of sulphur-waters, of mercury, cinchona, chamomilla, or from the fat of pork; rheumatic and arthritic affections, with swellings; inflammatory otalgia; dyspepsia, with vomiting of food; mucous or bilious diarrhea; inflammatory swelling of the testes; organic affections of the heart; inflammatory swelling of the legs and feet.

**GENERAL SYMPTOMS.**—Sharp, drawing, and jerking pains in the muscles, aggravated at night, or in bed in the evening, as well as by the heat of the room, mitigated in the open air, and often accompanied by torpor, with paralytic weakness or hard swelling of the parts affected; shifting pains which pass rapidly from one part to the other, often with swelling and redness of the joints.

*Skin.*—Red spots like measles or nettle-rash; chilblains, with bluish-red swellings, heat and burning, or pulsative pains.

*Head.*—Vertigo, as during intoxication; headaches in the evening after lying down, or at night, compression sometimes mitigates them.

*Eyes.*—Pressive, or sharp, shooting pain in the eyes, with inflammation; styes on the eyelids.

*Ears.*—Shootings, with itching, or sharp, jerking pains and contraction in and round the ears; the pains sometimes



come on by fits, attack the whole head, appear almost insupportable, and almost cause the loss of reason; inflammatory swelling, heat, and erysipelatous redness of the ear and auditory duct, as well as the surrounding external parts; tingling, roaring, and humming in the ears; hardness of hearing.

*Nose.*—Obstruction of the nose and dry coryza, principally in the evening and in the heat of a room.

*Face.*—Face pale, and sometimes with an expression of suffering.

*Teeth.*—Sharp, shooting pains in the teeth, or drawing, jerking pains, as if the nerve were tightened, then suddenly relaxed, or pulsative, digging, and gnawing pains, often with pricking in the gums, and shivering worse in the evening, or afternoon, in the heat of the bed, or of a room, mitigated by cold water or fresh air.

*Mouth.*—Tongue loaded with a thick coating, of a whitish or yellowish color.

*Throat.*—Pain, as if from excoriation in the throat, as if it were all raw, with sensation as if these parts were swollen, principally when swallowing, or accumulation of tenacious mucus, which covers the parts affected.

*Appetite.*—Bitter or sour taste in the mouth, immediately after having eaten; sensation of derangement in the stomach, similar to that caused by fat pork or rich pastry; after eating, nausea, and pressure in the pit of the stomach, and many other sufferings.

*Stomach.*—Nausea and vomitings take place in the evening or at night, and after eating or drinking, with shivering, paleness of face, colic; pressive, spasmodic, contractive, and compressive pains in the stomach and precordial region, after a meal or in the evening; pulsations in the pit of the stomach.

*Feces.*—Loose evacuations, with colic and cuttings, of

greenish, bilious, or watery matter; blind and bleeding hemorrhoides, with itching, smarting, and pain, as if from excoriation.

*Genital Organs.*—Inflammatory swelling of the testes, and of the spermatic cord (sometimes only on one side), with pressive and drawing pains extending into the abdomen; metrorrhagia; black menstrual blood, with clots of mucus, or discharge of pale and serous blood; menses irregular, too tardy, or entirely suppressed, with colic, hysterical spasms in the abdomen; nausea, and vomitings, shiverings, and paleness of face; leucorrhœa, thick like cream.

*Larynx.*—Shaking cough in the evening, at night, aggravated when lying down, accompanied with a desire to vomit, or by a choking, as if caused by the vapor of sulphur; moist cough, with expectoration of white, tenacious mucus, or of thick, yellowish matter.

*Chest.*—Spasmodic constriction of the chest, or larynx, in the evening, or at night when lying horizontally; frequent and violent palpitation of the heart.

*Trunk.*—Rheumatic, tensive, and drawing pain in the nape of the neck.

*Arms.*—Sharp, jerking, and drawing pains in the shoulder-joint, as well as in the arms, hands, and fingers.

*Legs.*—Pain, as from sub-cutaneous ulceration in the legs and soles of the feet; swelling of the knees, with sharp, drawing, and shooting pains; great fatigue in the legs and in the knees, with trembling.

#### 49. RHEUM.

Diarrhea in children; sourness, colic, sleeplessness and cries of children.

*Fæces.*—Loose evacuations of a sour smell, preceded and followed by constrictive pinching in the abdomen.

## 50. RHUS TOXICODENDRON.

Rheumatic affections, with swelling; vesicular erysipelas; bad effects from a strain, dislocation, concussion, and other mechanical injuries, especially when attended with sufferings in the joints and synovial membranes; gastric affections; diarrhea and dysentery; coxalgia and spontaneous dislocation.

GENERAL SYMPTOMS. — Rheumatic and arthritic drawings, tension and tearing in the limbs, increased to the highest degree during repose, in bad weather, at night, and in the heat of the bed; red and shining swellings.

*Sleep.* — Disturbed sleep, with anxious and frightful dreams.

*Fever.* — Malignant fever, with loquacious delirium, violent pains in all the limbs, excessive weakness, dry or black tongue, dry, brownish, or blackish lips, heat and redness of the cheeks.

*Head.* — Heaviness and pressive fullness in the head; beating and pulsations in the head, especially in the occiput; balancing and sensation of fluctuation in the head at every step, as if the brain were loose.

*Eyes.* — Inflammation of the eyes and lids; swelling of the whole eye and surrounding parts.

*Ears.* — Swelling and inflammation of the parotids, with fever.

*Face.* — Erysipelatous inflammation and swelling of the face, with pressive and tensive shootings, and burning crawling; vesicular erysipelas, with yellow serum in the vesicles; humid eruption and thick scabs on the face.

*Mouth.* — A yellow and sometimes also a sanguinous saliva flows from the mouth at night.

*Appetite.* — Thirst, from a sensation of dryness in the mouth.

*Fæces*.—Loose, sanguinous, serous, or slimy evacuations; obstinate diarrhea.

*Larynx*.—Cough, excited by a tickling in the ramifications of the bronchi, short and dry.

*Chest*.—Shootings and lancinations in the chest and its sides.

*Trunk*.—Pains in the loins, as if beaten; stiffness of the nape and neck.

*Arms*.—Warts on the hands and fingers.

#### 51. RUTA.

Sufferings caused by mechanical injuries (contusion, fall, dislocation, injury of the bones, or of the periosteum.)

GENERAL SYMPTOMS.—Pains in the limbs, joints, and bones, as if beaten, or as if caused by a contusion, fall, or bruise, especially in the periosteum of the limbs.

#### 52. SAMBUCUS.

Spasmodic asthma and cramps in the chest; asthma Milari; croup.

GENERAL SYMPTOMS.—Desire to sleep, without being able to sleep; fever, with excessive perspiration, especially at night.

*Larynx*—*Chest*.—Attack of suffocating cough, with cries in children; wheezing and quick respiration; spasmodic paroxysm of suffocation at night, with tears, great agitation and tossing of hands; great difficulty in breathing.

#### 53. SANGUINARIA.

Arthritic and rheumatic pains.

GENERAL SYMPTOMS.—Pain in the tips of the fingers, as from ulceration; sensation of heaviness in the head; pains in the head which pass rapidly from one place to another, like electric shocks.

## 54. SECALE.

Gastric and bilious affections; Asiatic and sporadic cholera; diarrhea, especially in old men; metrorrhagia of weak women.

*Sleep.*—Great desire to sleep, and deep, lethargic sleep.

*Eyes.*—Eyeballs sunk deep in the sockets.

*Face and Teeth.*—Face pale, discolored, yellow, wan, with the eyes hollow and surrounded by a blue circle.

*Stomach.*—Excessively painful sensibility and anguish in the pit of the stomach; burning sensation in the stomach and abdomen.

*Fæces.*—Loose, frequent evacuations, with serous or slimy fæces, and sudden prostration of strength; involuntary evacuations.

*Urine.*—Metrorrhagia, of a black, liquid blood, flowing, especially during a slight movement; miscarriage.

*Legs.*—Contraction of the legs and toes; troublesome cramps in the calves of the legs and soles of the feet, especially at night; mortification of the toes.

## 55. SEPIA.

GENERAL SYMPTOMS.—The symptoms disappear during every violent exercise, except on horseback, and are aggravated during repose, as also in the evening; uneasiness and throbbing in all the limbs; violent ebullition of blood during the night, with pulsation in the whole body; great tendency to take cold, and sensitiveness to cold air.

*Skin.*—Itching in different parts, which changes to a burning sensation; brown, reddish, and liver spots on the skin; deformity of the nails.

*Fever.*—Profuse perspiration from the slightest movement; sadness and dejection with tears; susceptibility and peevishness.

*Head.*—Attacks of headache, with nausea and vomiting when shaking or moving the head, and also at every step; semi-lateral headache; violent congestion of blood to the head.

*Eyes.*—Pressure on the eyeballs; inflammation, redness, and swelling of the eyelids, with styes.

*Ears.*—Shootings in the ears.

*Nose.*—Scabby and ulcerated nostrils.

*Face.*—Yellow color of the face; yellow streak on the nose and cheeks in the form of a saddle; yellow color and herpetic eruption round the mouth.

*Teeth.*—Toothache when compressing the teeth, when touching them, and when speaking, as also from the slightest current of cold air.

*Throat.*—Hawking up of mucus, especially in the morning.

*Appetite.*—Putrid, or sour taste; repugnance and dislike to food.

*Stomach.*—Throbbing in the pit of the stomach.

*Abdominal Region.*—Enlargement of the abdomen (in women who have had children.)

*Fæces.*—Ineffectual desire to evacuate; greenish diarrhea, often with putrid or sour smell, or especially in children; prolapsus recti; protrusion of hamorrhoides.

*Urine.*—Wetting the bed during the first sleep.

*Genital Organs.*—Bearing down in the uterus; prolapsus uteri; leucorrhœa, of a yellow or greenish red water.

*Larynx.*—Dry cough, which seems to arise from the stomach, especially when in bed in the evening, and often with nausea and bitter vomiting; cough excited by a tickling, and accompanied with constipation, detached with difficulty.

*Chest.*—Dyspnœa, oppression of the chest, and short breath when walking and ascending, as well as when lying



in bed, in the evening, and at night; ebullition of blood in the chest, and violent palpitation of the heart.

#### 56. SILICEA.

Obstruction, inflammation, induration, and ulceration of the glands; inflammation, softening, ulceration, and other diseases of the bones; ulcers, almost of all kinds; ulceration of the mammæ; chronic coryza and obstinate disposition to take cold in the head; panaritium.

*Skin.*—Mild and malignant suppurations, especially in membranous parts; carbuncles.

*Head.*—Vertigo, with nausea, which mounts from the back to the nape of the neck and head; tearing pains in the head, often semi-lateral; moist scald-head.

*Eyes.*—Redness of the eyes; swelling of the lachrymal gland; black spots before the sight.

*Ears.*—Hardness of hearing; swelling and induration of the parotids.

*Nose.*—Scabs, pimples, and ulcers in the nose.

*Face.*—Ulcers on the red part of the lip; herpes on the chin; swelling of the submaxillary glands.

*Appetite.*—After a meal, sour taste in the mouth; pressure in the stomach, water-brash, vomiting.

*Stomach.*—Nausea every morning.

*Fæces.*—Constipation, and slow, hard fæces.

*Trunk.*—Inflammatory abscess in the lumbar region; swelling and deviation of the spine; swelling of the glands of the nape of the neck, on and under the axillæ, sometimes with induration; suppuration of the axillary glands.

*Arms.*—Burning sensation in the ends of the fingers; paniritium.

*Leg.*—Inflammatory swelling of the knee; caries in the tibia; offensive smell from the feet.

## 57. SPIGELIA.

Prosopalgia; vermiculous affections; diseases of the heart.

*Head.*—Pains in the head, aggravated by the slightest movement, or least noise.

*Eyes.*—Pains in the eyes, as if the eyeballs were too large; inflammation of the eyes, and ulceration of the eyelids; hanging down of the eyelids, as from paralysis.

*Face.*—Jerking tearings, burning sensation and pressure in the region of the zygomatic process, or violent pains which can not bear the slightest touch, or the least movement.

*Teeth.*—Jerking toothache, or pulsative tearings, especially in the carious teeth, aggravated by cold water, or by contact with the open air; offensive exhalation from the mouth.

*Fæces.*—Discharge of mucus from the anus, without fæces; lumbrici and ascarides from the rectum; itching and sensation as if something were creeping in the rectum and anus.

*Chest.*—Danger of suffocation from the least movement, and especially when raising the arms; violent palpitation of the heart, perceptible to the sight and hearing, often attended with anxious oppression of the chest, aggravated by stooping the chest forward, and by sitting down.

## 58. SPONGIA.

Croup (before hepar, or alternate with it.)

GENERAL SYMPTOMS.—Swellings and indurations of the glands.

*Face.*—Face bloated, red, or bluish, with anxious expression of countenance.

*Larynx.* — Hoarseness ; pain in the larynx, on touching it and on turning the head ; sensation of obstruction in the larynx, with impeded respiration ; hollow, dry, barking or whistling cough, day and night, increased toward the evening.

*Chest.* — Wheezing respiration ; respiration quick, anxious, and difficult, sometimes with a fit of suffocation and mucous rattling in the chest ; ebullition of blood in the chest after the slightest effort, and the least movement, with obstructed respiration.

#### 59. STANNUM.

*Larynx.* — Hoarseness and roughness in the throat ; dry, violent, shaking cough ; cough excited by lying on the right side, with frequent expectoration of mucus, greenish, sweetish, or saltish.

*Chest.* — Pain in the chest, as if from excoriation ; sensation of weakness in the chest, as if it were empty, especially after speaking or expectorating.

#### 60. STAPHYSAGRIA.

Bad effects from crosses, with anger and indignation, or from vexation, with uneasiness.

*Moral Symptoms.* — Hypochondriacal humor, with indifference to every thing ; ill-humor, irascibility, and excessive anger, so as to fling about violently whatever is at hand.

*Eyes.* — Itching and biting smarting in the internal canthi ; inflammation in the eyes, which are surrounded with pimples.

*Ears and Nose.* — Coryza, with ulcerated nostrils.

*Face.* — Painful swelling and induration of the submaxillary glands.

*Teeth.* — Toothache, immediately after a meal, after cold drinks, or inhaling cold air, worse in the morning.

## 61. STRAMONIUM.

Convulsions, St. Vitus' dance, and other spasmodic affections; hydrocephalus.

GENERAL SYMPTOMS.—Convulsive movements and jerks, especially when touched, or from fixing the eyes on brilliant objects (such as a candle, a mirror, or water), or else appearing periodically.

*Sleep.*—Deep sleep, with snoring.

*Fever.*—Small, quick pulse.

*Moral Symptoms.*—Ungovernable fury, desire to bite, to strike.

*Head.*—Congestion of blood to the head, with heat in it.

*Eyes.*—Eyes fixed and sparkling; cloudiness of the sight; illusions of vision.

*Face and Teeth.*—Face bloated and puffed up with blood; grinding of the teeth.

*Trunk and Extremities.*—Opisthotonus; convulsive movements of the arms, above the head; clenched fists.

## 62. SULPHUR.

GENERAL SYMPTOMS.—Great sensitiveness to the open air and to the wind; the majority of the sufferings are aggravated, or appear at night, or in the evening, and also during repose, when standing for a long time, and by exposure to cold air; they disappear when walking or moving the parts affected, and also in the warmth of a room; but the heat of the bed renders the nocturnal pains insupportable.

*Skin.*—Itching in the skin, at night in bed; scabious eruptions, with burning itching; desquamation and excoriation of the skin in several places; ulcers, with elevated margins, surrounded by itchy pimples; inflammation, swelling, and induration, or suppuration of the glands.

*Fever.* — Frequent and profuse perspiration day and night; aptness to perspire when working, partial perspiration, principally on the head.

*Head.* — Fullness, pressure, and heaviness in the head, principally in the forehead; congestion of blood to the head, with pulsative, clucking, hammering sensations, and feeling of heat in the brain; quotidian, periodical, and intermittent headaches, appearing principally at night, or in the evening in bed, or in the morning; movement, walking, the open air, and meditation often excite or aggravate the headaches.

*Eyes.* — Itching, tickling, and burning sensation in the eyes and eyelids; inflammation, swelling, and redness of the sclerotica, conjunctiva and eyelids; pustules and ulcers round the orbits, as far as the cheeks; great sensitiveness of the eyes to the light.

*Ears.* — Obstruction and sensation of stoppage in the ears, on one side only; humming and roaring in the ears.

*Nose.* — Inflammation, ulceration, and scabs in the nostrils.

*Face.* — Heat and burning sensation in the face, with dark redness of the whole face, circumscribed redness of the cheeks.

*Teeth.* — Appearance or aggravation of toothache, principally in the evening, at night, or in the open air, and also from a current of air, from cold water, eating and masticating.

*Mouth.* — Aphthæ in the mouth and on the tongue.

*Throat.* — Pressure, as if from a plug, or from a tumor in the throat.

*Appetite.* — Dislike to sweet and acid things.

*Abdominal Region.* — Shootings in the abdomen, principally in the left side when walking; pains in the abdomen, principally at night, or after eating and drinking, mitigated by bending forward.

*Fæces.*—Constipation, and hard, knotty, and insufficient evacuations; hemorrhoides.

*Genital Organs.*—Excoriation between the thighs and groins.

*Larynx.*—Moist cough, with profuse expectoration of thick, whitish, or yellowish mucus; when coughing, pain as if from excoriation, or shootings in the chest.

*Chest.*—Obstructed respiration, dyspnæ and attacks of suffocation, principally when lying down at night, and also during sleep; periodical spasms in the chest; shootings in the chest or sternum, or extending as far as the back, or into the left side.

*Trunk.*—Pinching and burning sensation between the shoulder-blades.

#### 63. TABACUM.

GENERAL SYMPTOMS.—Cramps in the limbs.

*Stomach.*—Frequent nausea, especially when moving; the vomiting is renewed from the slightest movement.

*Fæces.*—Violent diarrhea, fetid, or yellowish green, also at night, accompanied and followed by violent tenesmus, and burning in the anus.

*Arms.*—Cold perspiration on the hands; cramps in the fingers.

#### 64. TARTARUS.

Varioloids; gastric and bilious affections; asphyxia of new-born infants.

*Skin.*—Eruption of pustules, like varioloids, with red areola, afterward covered with a crust and leaving a scar.

*Sleep.*—Invincible drowsiness, with deep and stupifying sleep.

*Fever.*—Pulse quick, weak, or full.

*Head.*—Chronic trembling of the head.



*Appetite.* — Bitter taste in the mouth ; excessive desire for acid things.

*Stomach.* — Continued nausea ; much vomiting, with violent efforts.

*Fæces.* — Slimy diarrhea.

*Larynx.* — Cough, with vomiting of food after a meal ; hollow cough, with rattling of mucus in the chest.

*Chest.* — Paralysis of the lungs ; rattling of mucus in the chest.

#### 65. THUJA.

Rheumatic and arthritic affections ; syphilitic and sycotic affections.

*Head.* — Vertigo, when rising from a seat and when lying down.

*Eyes.* — Burning sensation in the eyes ; inflammatory swelling of the eyelids, with hardness.

*Face.* — Boring and digging pain in the face and cheek-bones ; gnawing in the (carious) teeth, with painful sensitiveness of the whole side of the head, greatly aggravated by contact with cold things, or by mastication.

*Mouth.* — Ranula, on both sides of the tongue, transparent, bluish red, gray, and, as it were, gelatinous.

*Urine.* — Sensation, as if a drop were flowing into the urethra ; burning sensation in the urethra during the emission of urine ; itching in the urethra.

*Genital Organs.* — Condylomata on the glans and prepuce, moist, itching.

#### 66. URTICA URENS.

Burns ; dysentery ; nettle-rash.

#### 67. VERATRUM.

Sporadic or Asiatic cholera ; diarrhea, of different kinds, also those produced by cold drinks, when one is over-heated.

GENERAL SYMPTOMS. — Pains in the limbs, which are rendered insupportable by the heat of the bed, and disappear completely, when walking; sudden, general, and paralytic prostration of strength.

*Fever.* — General coldness of the whole body, and cold, clammy perspiration; fever, with external coldness; pulse slow, and almost extinct.

*Head.* — Attack of headache, with paleness of the face, nausea, and vomiting; cold perspiration on the forehead.

*Nose.* — Icy coldness of the nose.

*Face.* — Face pale, cold, hippocratic, wan, with the nose pointed, and a blue circle round the eyes; cold perspiration on the face; lips dry, blackish, and cracked.

*Teeth.* — Grinding of the teeth.

*Mouth.* — Sensation of coldness on the tongue; tongue dry, blackish, cracked.

*Appetite.* — Immediate vomiting and diarrhea, however little is eaten.

*Stomach.* — Violent nausea, with desire to vomit, with excessive thirst; violent vomiting, with continued nausea, great exhaustion and desire to lie down; vomiting of black bile and blood; vomiting, with diarrhea; the least drop of liquid and the slightest movement excites the vomitings; burning sensation in the pit of the stomach.

*Abdominal Region.* — Excessively painful sensitiveness of the abdomen, when touched; cramps in the abdomen, and cuttings, as if from knives; burning sensation through the whole extent of the abdomen, as if from hot coals.

*Fæces.* — Loose, blackish, greenish, brownish evacuations; unnoticed evacuation of liquid fæces.

*Urine.* — Retention of urine; urine diminished.

*Genital Organs.* — Menses suppressed, with delirium.

*Chest.* — Chest very much oppressed; cramp in the

chest, with painful constriction ; violent palpitation of the heart, which causes heaving of the chest.

*Arms.* — Icy coldness of the hands ; cramps in the fingers.

*Legs.* — Violent cramps in the calves of the legs and feet ; icy coldness of the feet.

#### 68. ZINCUM.

GENERAL SYMPTOMS. — Tearing in the limbs, aggravated whenever one is over-heated or takes exercise ; pain, as if from excoriation ; the symptoms are aggravated in an extraordinary degree by Chamomile, Nux vomica and wine, substances which also excite them, particularly the nocturnal uneasiness and constipation ; the majority of the symptoms manifest themselves after dinner and toward the evening ; violent trembling of the whole body, particularly after every mental emotion ; heaviness, lassitude and excessive weakness, principally when walking, or on waking in the morning.

*Skin.* — Herpes and herpetic ulcers.

*Sleep.* — Sleep by day and continued desire to sleep, especially in the morning, or after a meal, with inclination to yawn ; unrefreshing sleep ; fantastic, frightful, agitated, or disgusting and terrific dreams, with talking and cries during sleep, excessive coldness of the feet at night.

*Fever.* — Febrile shuddering, with flushes of heat ; violent trembling of the limbs, short and hot breath, and pulsation in the whole body.

*Moral Symptoms.* — Fretful, peevish humor, with dislike to conversation, especially in the evening ; aversion to labor ; fickleness, with sadness toward noon and joy in the evening, and *vice versa*.

*Head.* — Pressive cephalalgia in the forehead, with confusion, or else in the temples and occiput, principally in the

morning ; shootings and tearing in the head, especially in the sides, temples, forehead and occiput, aggravated after dinner ; itching and sensation of excoriation in the hairy scalp, or pain, as if from ulceration ; baldness.

*Eyes.* — Itching, smarting, and sensation as if from excoriation in the eyes, eyelids, and internal canthi ; falling down and paralysis of the upper eyelids.

*Ears.* — Ootalgia, with tearing lacerations and external swelling, especially in children ; flow of fetid pus from the ears ; humming in the ears.

## ERRATA.

Page 37 read "poisoned," instead of "poisonous."

" 59, line 14, read "defer the farther," instead of "wait with the."

" 59, " 30, at its end, insert the word "alternately."

" 88, " 12, read "it," instead of "is."

" 327, " 5, insert after emptied "through."

" 349, " 16, read "glans," instead of "gland."

" 351, last line, read "glans," instead of "glands."

" 355, insert after chlorosis "green disease."

" 398, line 8, insert after as "this."

" 415, last line, "serum," instead of "scrum "

" 421, line 8, after ventricle insert "and auricle."


" 421, " 9, read "the auricles," instead of "these ventricles."

" 421, " 12 and 17, read "heart," instead of "ventricle."

" 422, " 5, read "is," instead of "be."

" 447, " 20, omit "which."

" 447, " 22, read "and that," instead of "such as."





## GLOSSARY OF MEDICAL TERMS.

---

- ABORTUS.** Miscarriage; abortion.
- ABSCESS.** A collection of pus seated in any particular organ or tissue.
- ADHESION.** In surgery, the reunion of parts that have been divided, by means of a special kind of inflammation, denominated the *adhesive*. In pathology, the morbid union of parts which are naturally contiguous, though not adherent, through the instrumentality of adhesive inflammation.
- ADHESIVE INFLAMMATION.** The process by which wounds are united. It is often synonymous with *union by the first intention*.
- ADYPSIA.** The absence of natural thirst.
- ALKALI.** A substance which unites with acids in definite proportions, so as to neutralize their properties more or less perfectly, and to form salts. It changes vegetable blues to green.
- ALLOPATHY.** A term used by hmcœopathic writers to designate the old practice of medicine in contradistinction to their own, now generally employed by both parties; literally implies curing one disease by another.
- AMENORRHŒA.** Absence or stoppage of the menstrual flux.
- ANASARCA.** Dropsy of the cellular tissue, or membrane, immediately under the skin.
- ANGINA.** Sore throat. The term is also applied to diseases with difficult respiration.
- ANGINA MEMBRANACEA.** Croup.
- ANGINA PAROTIDEA.** Mumps.
- ANGINA PHARYNGIA.** Inflammation of the membrane which forms the pharynx.
- ANOREXIA.** Want of appetite.
- ANTHRAX.** Carbuncle.
- ANTIPHLOGISTIC.** Applied to remedies employed in the old system against inflammation; literally, against heat.
- ANTRUM HIGHMORI.** The maxillary sinus. A hollow or cavity above the teeth of the upper jaw, in the middle of the superior maxillary bone.
- ANUS.** The inferior opening of the rectum.
- APOPLEXIA.** Apoplexy; a loss of voluntary motion and consciousness. See *Diagnosis* under this head.
- APYREXIA.** The intervals between febrile paroxysms.
- ARTHRITIS.** Gout.
- ASCARIS, plur. ASCARIDES.** Pin-worms.
- ASPHYXIA.** Apparent death.
- ASTHENIC.** Low; applied to disease; literally, want of strength.
- ASTRINGENTS.** Medicaments used in the old practice to contract the animal fibre.
- ATONY.** A want of tone or energy in the muscular power.
- ATROPHY.** A morbid state of the digestive system, in which the food taken into the stomach fails to afford sufficient nourishment. A wasting of the whole, or of individual parts of the body.
- AUSCULTATION.** The detection of symptoms by the ear in disease.
- BILIIOUS.** Connected with the secretion of bile.
- BLEPHARITIS.** Inflammation of the eyelids.
- BORBORYGMUS.** Rumbling in the intestines, caused by flatus or wind.
- BRONCHIA; BRONCHI.** The tubes into which the trachea or windpipe divides.

- BRONCHITIS.** Inflammation of the ramifications of the windpipe.
- BULIMY ; BULIMIA.** Canine, or excessive hunger.
- CADAVEROUS.** Resembling a corpse.
- CÆCUM.** The blind gut; so called from its being perforated at one end only.
- CARCINOMA.** Cancer, adj. *Carcinomatous*.
- CARDIALGIA.** Spasm of the stomach.
- CARDITIS.** Inflammation of the heart.
- CARIES.** Ulceration of the bones.
- CAROTIDS.** The name of two large arteries of the neck.
- CARPOLOGIA.** Picking at the bed clothes.
- CARTILAGE.** Gristle.
- CATAMENIA.** The menstrual flux.
- CATARRI.** Cold; used, also, to express inflammation of the mucous membrane.
- CATARRHAL OPHTHALMIA.** Simple inflammation of the conjunctiva.
- CATHARTIC.** Purgative.
- CELLULAR TISSUE.** The fine, net-like membrane enveloping or connecting most of the structures of the human body.
- CEPHALALGIA.** Headache.
- CEREBRAL.** Appertaining to the brain.
- CERVICAL.** Belonging to the neck.
- CESSATIO MENSII.** Discontinuance of the menstrual flux.
- CHLOROSIS.** Green sickness.
- CHRONIC.** Long continued, in contradistinction to acute.
- CICATRIX.** A scar left after the healing of a wound, etc.
- CLAVI PEDIS.** Corns.
- CLONIC SPASM.** A spasm which is not of long duration. It is opposed to *tonic* spasm, which see.
- COAGULA.** Clots of blood.
- COAGULABLE LYMPH.** The term given to the fluid which is slowly effused into wounds, and afterward forms the uniting medium or cicatrice.
- COLIC.** Gripping in the intestines.
- COLLAPSE.** Failing of vitality.
- COLLIQUATIVE.** Excessive discharge of any secretion.
- COMA.** Drowsiness.
- COMA SOMNOLENTUM.** Drowsiness, with relapse thereunto on being roused.
- COMATOSE.** Drowsy.
- COMPRESS.** Soft lint, linen, etc., folded together so as to form a pad, for the purpose of being placed, and secured by means of a bandage, on parts which require pressure.
- CONGESTIO AD CAPUT.** Determination of blood to the head.
- CONGESTIO AD PECTUS.** Determination of blood to the chest.
- CONGESTION.** Overfullness of the bloodvessels of some particular organ.
- CONJUNCTIVA.** The membrane lining the eyelids, and extending over the fore part of the eyeballs.
- CONTAGION.** Propagation of a disease by contact.
- CORNEA.** The anterior transparent portion of the eye. It is of a horny consistence.
- CORYZA.** Cold in the head.
- COXAGRA.** Inflammation of the hip-joint. Literally, seizure or pain in the hip.
- COXALGIA.** Literally, pain in the hip; inflammation of the hip joint.
- CRANIUM.** The skull.
- CREPITATION.** Grating sensation, or noise, such as is caused by pressing the finger upon a part affected with emphysema; by the ends of a fractured bone when moved; or by certain salts during calcination.
- CREPITANT RHONCHUS, or RALE.** The fine crackling noise heard in consequence of the passage of air through a viscid fluid. It is heard in the first stage of inflammation of the lungs.
- CREPITUS.** Crackling or grating.
- CUTANEOUS.** Appertaining to the skin.
- CUTICLE.** The outer or scarf skin.
- CYSTITIS.** Inflammation of the bladder.
- DEGLUTITION.** The act of swallowing.
- DELIRIUM.** Derangement of the brain, raving.
- DEPLETION.** Abstraction of the fluids; generally applied to venesection.
- DESCICATION.** A drying up.
- DESQUAMATION.** Falling off of the epidermis in form of scales.
- DIAPHRAGMITIS.** Inflammation of the diaphragm (muscular partition between the thorax and abdomen).
- DIAGNOSIS.** Distinction of maladies.
- DIARRHEA.** Looseness of the bowels.
- DIATHESIS.** Constitutional tendency.
- DIETETIC.** Relating to diet.
- DIPLOPIA.** Affection of the eyes, in which objects appear double or increased in number.
- DIURETIC.** Medicines which increase the secretion of urine.
- DORSAL.** Appertaining to the back.
- DRASTIC.** Powerful purgatives.
- DUODENUM.** The first intestine after the stomach, so called from its length; the twelve-inch gut.

- DYSCRASIA.** A morbid condition of the system; adj. *Dyscrastic*.
- DYSECO A.** Deafness.
- DYSMENORRHEA.** Painful menstruation.
- DYSPEPSIA.** Indigestion; literally, difficulty of appetite.
- DYSPNŒA.** Difficulty of respiration. Shortness of breath.
- DYSURIA.** Difficulty in passing urine.
- EFFUSION.** A pouring out, or escape of lymph or other secretion.
- EMACIATION.** A falling off in the flesh.
- EMETIC.** Provoking vomiting.
- ENCEPHALITIS.** Inflammation of the brain and membranes.
- ENDEMIC.** Peculiar to a particular locality.
- ENDOCARDITIS.** Inflammation of the internal parts of the heart.
- ENTERALG A.** Colic.
- ENTERITIS.** Inflammation of the intestines.
- EPHEMERAL.** Of a day's duration.
- EPHIALTES.** Nightmare.
- EPIDEMIC.** Diseases arising from general causes.
- EPIGASTRIUM.** The region of the stomach.
- EPILEPSY; EPILEPSIA.** Falling sickness.
- EPISTAXIS.** Bleeding from the nose.
- EPITHELIUM.** The cuticle.
- ERYSIPELAS.** St. Anthony's fire. Rose. A disease of the skin.
- ERYSIPELAS PNEUMONODES.** Phlegmonous erysipelas.
- ERYSIPELAS ŒDEMATODES.** Œdematous erysipelas.
- ERYSIPELAS ERRATICUM.** Wandering erysipelas.
- ERYSIPELAS GANGRENOSUM.** Gangrenous erysipelas.
- ERYSIPELAS NEONATORUM.** Induration of the cellular tissue in infants.
- EXACERBATION.** Aggravation of fever, etc.
- EXANTHEMA.** Eruption terminating in exfoliation.
- EXPECTORATION.** Discharge of any matter; phlegm; pus from the chest.
- EXUDATION.** Discharge of fluid from the skin, etc.
- FÆCES.** Alvine excrement.
- FASCIA.** In anatomy, the tendinous expansion of muscles which bind parts together, are called *fasciæ*.
- FAUCES.** The throat.
- FEBRIS.** Fever.
- FEBRIS NERVOSA.** Nervous fever, or typhus.
- FEMUR.** The bone of the thigh.
- FIRST INTENTION.** See *Union by the*.
- FISTULA.** An obstinate, tube-like sore, with a narrow orifice; adj. *Fistulous*.
- FISTULA LACHRYMALIS.** An ulcerated opening in the lachrymal sac.
- FLATUS.** Wind in the intestines. Flatulency.
- FŒTUS.** The infant in the womb.
- FOMENTATION.** The application of flannel wet with warm water.
- FUNCTIONAL DISEASES.** Those in which there is supposed to be only derangement of action.
- FURUNCULUS.** A boil.
- FURUNCULUS MALIGNANS.** Carbuncle.
- GANGRENE.** Incipient mortification. adj. *Gangrenous*.
- GASTRALGIA.** Pain in the stomach.
- GASTRIC.** Belonging to the stomach.
- GASTRITIS.** Inflammation of the stomach.
- GASTRODYNIA.** Vide *Cardialgia*.
- GLAND.** A small body met with in many parts of the body, and consisting of various tissues, bloodvessels, nerves, etc.
- GLOSSITIS.** Inflammation of the tongue.
- GLOTTIS.** Opening of the windpipe. The superior opening of the larynx.
- GRANULATION.** See *Incarnation*.
- HEMATESIS.** Vomiting of blood.
- HEMOPTYSIS.** Discharge of blood from the lungs. Spitting of blood.
- HEMORRHAGE.** Discharge of blood.
- HEMORRHOIDES.** Piles.
- HECTIC FEVER.** Habitual or protracted fever.
- HELMINTHIASIS.** Worm disease.
- HEMIPLEGIA.** Paralysis of one side of the body, longitudinally.
- HEPATITIS.** Inflammation of the liver.
- HEPATIZATION.** Structural derangement of the lungs, the result of inflammation, changing them into a substance resembling the liver; hence its name.
- HERNIA.** Rupture.
- HERNIA CONGENITAL.** Congenital hernia. Literally, hernia from birth.
- HERPES CIRCINNATUS.** Ringworm.
- HORDEOLUM.** Sty.
- HYDROCEPHALUS.** Water in the head.
- HYDROPHOBIA SYMPTOMATICA.** Symptoms resembling those arising from hydrophobic virus, appearing during the course of other diseases.
- HYPERTROPHY.** A morbid increase of any organ, arising from excessive nutrition.
- HYPOCRATIC.** Sunken and corpse-like.

- HYPOCHONDRIUM.** Region of the abdomen, contained under the cartilage of the false ribs.
- HYPOCHONDRIASIS.** Spleen disease; great depression of spirits, with general derangement; adj. *Hypochondriacal*.
- HYPOGASTRIUM.** The lower anterior portion of the abdomen.
- HYSTERIA.** Nervous affection; almost peculiar to females.
- ICHOR.** A thin, watery discharge secreted from wounds, ulcers, etc.; adj. *Ichorous*.
- ICTERUS.** Jaundice.
- ICTERUS NEONATORUM.** Jaundice of infants.
- IDIOPATHIC.** Original or primary disease.
- IDIOSYNCRASY.** Individual peculiarity.
- ILIUM.** The haunch-bone. It, together with the pubis, sacrum, and ischium, contributes to form the pelvis.
- ILEUS MISERERE.** A form of colic, a twisting pain in the region of the navel.
- INCARCERATED.** Strangulated or constricted; a term applied to rupture.
- INCARNATION.** The process by which abscesses or ulcers are healed; this takes place by means of little grain-like fleshy bodies, denominated granulations, which form on the surface of ulcers or suppurating wounds, etc., and serve the double purpose of filling up the cavities and bringing closely together and uniting their sides.
- INCUBUS.** The nightmare.
- INFECTION.** Propagation of disease by effluvia.
- INFILTRATION.** Diffusion of fluids into the cellular tissue.
- INTEGUMENTS.** The coverings of any part of the body. The skin, with the adherent fat and cellular membrane, form the common integuments.
- INTENTION.** See *Union by the first*.
- ISCHIAS.** Pain in the hip.
- ISCHURIA.** Suppression of urine.
- LACHRYMATION.** Tear-shedding.
- LACTATION.** Suckling; also the process of the secretion of milk.
- LACTEAL.** Appertaining to the process of the secretion of milk.
- LACTIFEROUS.** Conducting or conveying the milk.
- LARYNGEAL.** Belonging to the larynx.
- LARYNGITIS.** Inflammation of the larynx.
- LARYNX.** Upper part of the windpipe.
- LESIONS.** Injuries inflicted by violence, etc.
- LESION, ORGANIC.** Structural derangement, or injury.
- LEUCO-PHLEGMATIC.** Torpid or sluggish; mostly applied to a temperament characterized by want of tension of fibre; with light hair, and general inertness of the physical and mental powers.
- LEUCORRŒA.** Female sexual weakness; vulg. *Whites*.
- LOCHIA.** Discharge from the womb after delivery.
- LUMBAGO.** Rheumatism in the loins.
- LUMBAR.** Appertaining to the loins.
- LUMBRICUS.** The round or long worm.
- LUXATION.** Dislocation.
- LYMPH.** A colorless liquid, circulating in the lymphatics.
- LYMPHATIC.** As applied to temperament; same as leucophlegmatic.
- LYMPHATICS.** Absorbent vessels with glands and valves distributed over the body.
- LYMPHATIC GLANDS.** CONGLOBATE GLANDS. These are composed of a texture of absorbents, or lymphatic vessels, connected together by a cellular membrane.
- MAMMA.** The breast in the female; adj. *Mamillary*.
- MANIA.** Insanity; madness.
- MARASMUS.** A wasting away of the body.
- MATERIA MEDICA.** The title of that splendid work of the immortal Hahnemann, in which the true properties of medicaments are given, as determined by experiment upon the healthy body.
- MAXILLARY.** Appertaining to the jaws. The superior and inferior *maxillary bones* from the upper and lower jaws.
- MEGRIM.** A pain affecting only one side of the head.
- MEIBOMEAN GLANDS.** Small glands within the inner membrane of the eyelids.
- MENORRHAGIA.** Excessive discharge or blood from the uterus.
- MENSES AND MENSTUAL FLUX.** The monthly period.
- MENINGITIS SPINALIS.** Inflammation of the spinal membranes.
- METASTASIS.** The passing of a disease from one part to another.
- METEORISMUS.** Extreme inflation of the intestines.
- METORRHAGIA.** Discharge of blood from the womb.
- MIASM, or Miasma (Marsh.)** Peculiar effluvia or emanations from swampy grounds.
- MICTRITION.** Urination.

- MILIARIA.** Eruption of minute transparent vesicles of the size of millet seeds: millary eruption.
- MILIARIA PURPURA.** Scarlet-rash.
- MORBUS COXARIUS.** Disease of the hip; hip-disease.
- MUCOUS MEMBRANE.** The membrane which lines the sides of cavities which communicate with the external air, such as that which lines the mouth, stomach, etc.
- MUCUS.** One of the primary animal fluids; secretion from the nostrils.
- MYELITIS.** Inflammation of the spinal marrow.
- MYOPIA.** Short sight; near-sightedness.
- NARCOTIC.** Having the property of inducing sleep.
- NASAL.** Belonging to the nose.
- NASAL CARTILAGES.** The cartilages of the nose.
- NEPHRITIS.** Inflammation of the kidneys.
- NEURALGIA FACIALIS.** Face-ache.
- NODOSITIES.** Swellings; nodes, a swelling of the bone or thickening of the periosteum.
- NOTALGIA.** Pains in the loins.
- OCCIPUT.** The posterior part of the head.
- ODONTALGIA.** Toothache.
- ŒDEMA.** Swelling; dropsical swelling; adj. *Œdematous*.
- OLFACTION.** The act of smelling.
- OMENTUM.** The caul. The viscus consists of folds of the peritoneum connected together by cellular tissue; it is attached to the stomach, lying on the anterior surface of the bowels.
- OPHTHALMIA.** By this term is now usually understood simple inflammation of the *conjunctiva*. Catarrhal Ophthalmia.
- OPHTHALMIC NERVE.** The first branch given off from the Gasserian ganglion of the fifth pair of nerves; it divides into the lachrymal, frontal, and nasal nerves.
- OPHTHALMITIS.** Inflammation of the entire ball of the eye.
- ORGANIC DISEASE.** In pathology, diseases in which there is derangement or alternation of structure are termed organic.
- OS UTERI.** The mouth or opening of the womb.
- OSSICULA AUDITORIA.** The small bones of the ear. They are situated in the cavity of the tympanum, and are four in number: termed, the malleus, incus, stapes, and os-orbitulare.
- OTALGIA.** Earache.
- OTITIS.** Inflammation of the ear.
- OTORRHEA.** A discharge, or running from the ear.
- OZÆNA.** An ulcer situated in the nose. See *Ozæna*.
- PALATE BONES.** These are placed at the back part of the roof of the mouth, between the superior maxillary and sphenoid bones, and extended from thence to the floor of the orbit.
- PALPITATIO CORDIS.** Palpitation of the heart.
- PANARIS.** Whitlow; panaritium; paronychia.
- PANCREAS.** A gland situated transversely behind the stomach.
- PARALYSIS.** Palsy.
- PARALYSIS PARAPLEGICA.** Paralysis affecting one-half of the body transversely.
- PARENCHYMA.** The connecting medium of the substance of the lungs.
- PAROTITIS.** Inflammation of the parotid gland; the *mumps*.
- PAROXISM.** A periodical fit of a disease.
- PARTURITION.** The act of bringing forth.
- PATHOGENETIC.** The producing or creating of abnormal phenomena.
- PATHOGNOMIC.** Characteristic of and peculiar to any disease.
- PATHOLOGY.** The investigation of the nature of disease.
- PECTORAL.** Appertaining to the chest.
- PECTUS.** The chest.
- PELVIS.** The basin-shaped cavity below the abdomen, containing the bladder, and rectum, and womb in woman.
- PERCUSSION.** The act of striking upon the chest, etc., in order to elicit sounds to ascertain the state of the subjacent parts.
- PERICARDITIS.** Inflammation of the Pericardium (sac containing the heart.)
- PERINÆUM.** The space between the anus and the external sexual organs.
- PERIOSTEUM.** The membrane which envelopes the bones.
- PERITONÆUM.** The serous membrane which lines the cavity of the abdomen, and envelopes the viscera contained therein.
- PERITONITIS.** Inflammation of the peritonæum.
- PETECHIÆ.** Spots of a red or purple hue, resembling a flea bite.
- PHAGEDENIC.** A term applied to any sores which eat away the parts, as it were.

- PHARYNX.** The throat, or upper part of the gullet.
- PHASE.** Appearance, or change exhibited by any body, or by *disease*.
- PHLEBITIS.** Inflammation of the veins.
- PHLEGMATIC.** Vide *Leuco phlegmatic*.
- PHLEGMON.** An inflammation of that nature which is otherwise termed *healthy inflammation*.
- PIRENETIS.** Inflammation of the brain.
- PHTHISIS.** (Pulmonalis) Consumption, abscess of the lungs.
- PHYSIOLOGY.** The branch of medicine which treats of the functions of the human body.
- PLETHORA.** An excessive fullness of the bloodvessels.
- PLEURA.** The serous membrane which lines the cavity of the thorax or chest.
- PLEURITIS or PLEURISY.** Inflammation of the pleura.
- PLEURODYNTIA.** Pain or stitch in the side.
- PNEUMONIA, PNEUMONITIS, PERIPNEUMONIA.** Inflammation of the parenchyma of the lung.
- POLYPSUS.** A tumor most frequently met with in the nose, uterus, or vagina.
- PORRIGO SCUTULATA.** Ringworm of the scalp.
- PRÆCORDIAL REGION.** The fore-part of the chest.
- PROGNOSIS.** The faculty of predicting what will take place in diseases.
- PROLAPUS ANI.** Protrusion of the intestines.
- PROSOPALGIA.** Faceache.
- PRURIGO.** Itching of the skin.
- PSOAS MUSCLES.** The names of two muscles situate in the loins.
- PSORITIS.** Inflammation of the psosas muscles.
- PUBIS.** The pubic or share-bone.
- PUERPERAL FEVER.** Appertaining to childbed.
- PURIFORM.** Pus-like, resembling pus.
- PURULENT.** Of the character of pus.
- PUS.** Matter. A whitish, bland, cream-like fluid, found in abscesses, or on the surface of sores.
- PUSTULE.** An elevation of the scarf-skin, containing pus or lymph, and having an inflamed base.
- PYROSIS.** Heart-burn; water-brash.
- QUINSY.** Inflammatory sore throat.
- QUOTIDIAN.** Intermittent, about twenty-four hours intervening between the attacks.
- RABIES.** Madness arising from the bite of a rabid animal, generally applied to the disease showing itself in the brute creation.
- RACHITIS.** The rickets.
- RAUCITAS.** Hoarseness.
- RECTUM.** The last of the large intestines, terminating in the anus.
- REMITTENT.** A term applied to fevers with marked remissions, and generally subsequent exacerbation.
- REPERCUSSED.** Driven in.
- RESOLUTION.** A termination of inflammatory affections without abscess, mortification, etc. The term is also applied to the dispersion of swellings, indurations, etc.
- RHEUMATIC OPHTHALMIA.** Inflammation of the tunica albuginea, and of the sclerotica.
- ROSE.** A term applied to erysipelas, from its color.
- RUBEOLA.** Measles.
- SACRUM.** The bone which forms the base of the vertebral column.
- SALIVA.** The fluid secreted by the salivary glands into the cavity of the mouth.
- SANGUINEOUS.** Consisting of blood.
- SANIES.** A thin greenish discharge of fetid matter from sores, fistulæ, etc., adj. sanious.
- SCABIES. PSORA.** Itch.
- SCAPULA.** The shoulder blade.
- SCIATICA.** A rheumatic affection of the hip-joint.
- SCIATIC NERVE.** A branch of a nerve of the lower extremity.
- SCIRRHUS.** Indolent, glandular tumor, generally preceding cancer in an ulcerated form.
- SCLEROTICA.** The hard membrane of the eye; it is situated immediately under the conjunctiva.
- SCROBICULUS.** Pit of the stomach.
- SCORBUTUS.** Scurvy.
- SCROFULOUS OPHTHALMIA.** Inflammation of the conjunctiva, with slight redness, but great intolerance of light, and the formation of pimples, or small pustules.
- SECRETORY VESSELS, or ORGANS.** Parts of the animal economy, which separate or secrete the various fluids of the body.
- SEMI-LATERAL.** Limited to one side.
- SINUS.** A cavity or depression.
- SOLIDIFICATION.** Vide *Hepatitis*.
- SOMNOLENCE.** Disposition to sleep.
- SPECIFIC.** A remedy possessing a peculiar curative action in certain diseases.
- SPLEEN.** A spongy viscous organ, of a livid color, placed on the posterior part of the left hypochondrium.
- SPLENITIS.** Inflammation of spleen.



- SPLINTS.** Long, thin pieces of wood, tin, or strong pasteboard, used for preventing the extremities of fractured bones from moving so as to interrupt the process by which they are united.
- SPUTA.** Expectoration of different kinds.
- ST. ANTHONY'S FIRE.** Erysipelas.
- STERTOROUS.** Snoring.
- STOMACACE.** Canker or scurvy of the mouth.
- STRABISMUS.** Squinting.
- STRANGURY.** Painful discharge of urine.
- STERNUM.** The breast-bone.
- STETHOSCOPE.** An instrument to assist the ear, in examining the morbid sounds of the chest.
- STRICTURE.** A constriction of a tube or duct of some part of the body.
- STRUMA. SCROFULA.** The king's evil; adj. *Strumous*.
- STYE.** An inflammatory small tumor on the eyelid.
- SUB-MAXILLARY.** Under the jaw.
- SUB-MAXILLARY GLANDS.** Glands on the inner side of the lower jaw.
- SUB-MUCOUS TISSUE.** Placed under the mucous membrane.
- SUDORIFICS.** Medicines which produce sweating.
- SUGILLATION.** A bruise, or extravagated blood.
- SUPPURATION.** The morbid action by which pus is deposited, in inflammatory tumor, etc.
- SYNCOPE.** Fainting or swooning.
- SYNOCHA.** Continued inflammatory fever.
- SYNOVIA.** A peculiar, unctuous fluid secreted within the joints, which it lubricates, and thereby serves to facilitate their motions.
- SYNOVIAL MEMBRANE.** The membrane which lines the cavities of the joints, and secretes the synovia.
- TENIA.** Tape-worm.
- TARTAR.** A concretion encrusting the teeth.
- TEMPORAL.** Appertaining to temples.
- TENDON.** The white and shining extremity of a muscle.
- TENESMUS.** Painful and constant urging to alvine evacuations, without a discharge.
- TETANUS,** adj. *Tetanic*. A spasmodic rigidity of the parts affected.
- THERAPEUTICS.** That branch of medicine describes the action of the different means employed for the curing of diseases, and of the application of those means.
- THORAX.** The chest, or that part of the body situated between the neck and the abdomen.
- THRUSH.** Numerous small white vesicles in the mouth. See Thrush.
- TIC DOULOUREUX.** Face-ache.
- TINEA ANNULARIS. TINEA CAPITIS.** Ringworm of the scalp.
- TINEA FACIEI.** Milk-crust; milk scab.
- TITILLATION.** Tickling.
- TONIC.** Medicines which are said to increase the tone of the muscular fibre when debilitated and relaxed.
- TONSILS.** The oblong, sub-oval glands placed between the arches of the palate.
- TONSILLITIS.** Inflammation of the tonsils.
- TRACHEA.** The windpipe.
- TRACHEOTOMY.** An operation by opening the windpipe.
- TRAUMATIC.** Appertaining to wounds; arising from wounds.
- TREMOR.** Trembling.
- TRISMUS.** Lock-jaw.
- TRITURATION.** The reduction of a substance to minute division, by means of long-continued rubbing.
- TUBERCLE.** A small, round, eruptive swelling, anatomically speaking. In pathology, the name is applied to a peculiar morbid product occurring in various organs or textures, in the form of small, round, isolated masses of a dull whitish yellow, or yellowish gray color, opaque, unorganized, and varying in shape and consistence according to their stage of development and the texture of part in which they are engendered.
- TYPHOID.** Applied to diseases of a low character.
- UMBILICAL CORD.** The navel string.
- UMBILICUS.** The navel.
- UNION BY THE FIRST INTENTION.** The healing of wounds by adhesion; the growing together of the opposite surfaces of a wound when brought into close approximation, without suppuration or granulation. The latter process of healing is sometimes designated the *second intention*.
- URETHRA.** The urinary canal.
- URTICARIA.** Nettle-rash.
- UTERUS.** The womb.
- VARICELLA.** Pimples, quickly forming pustules, seldom passing into suppuration, but bursting at the point and drying into scabs. Chick-en-pock.
- VARIOLA.** Smallpox.
- VARIOLA. SPURIA. (Varicella.)** Chick-en-pock.

- 
- VARIX, plur. *Varices*. Swelling or enlargement of the veins.
- VENESECTIO. The abstraction of blood by opening a vein.
- VERTIGO. Giddiness, with a sensation as if falling.
- VESICLE. A small bladder-like eruption; an elevation of the cuticle containing a transparent watery fluid.
- VICARIOUS. Acting as a substitute.
- VIRUS. Contagion or poison.
- VISCID. Glutinous and gelatinous.
- VICUS, plur. *Viscera*. Any organ of the system. A bowl.
- VOMICA. An abscess of the lungs.
- ZYGOMATIC PROCESS. A thin narrow projection of bone, defining the squamous portion of the temporal bone at its base.

# INDEX.

Abdomen, dropsy of the . . . . .	337	Anthrax . . . . .	82
“ external injuries of the . . . .	339	Anury . . . . .	343
Abdominal inflammation . . . . .	289	Anus, itching of the . . . . .	335
Ablutions . . . . .	453	Aphonia . . . . .	210
Abortion . . . . .	368	Aphthæ . . . . .	383
Abscesses . . . . .	81	Apoplexia . . . . .	122
Abscess in the gums . . . . .	193	Apoplexy . . . . .	122
Acute dropsy . . . . .	254	Apparent death from hunger . . . .	25
Adherence of the after-birth . . . .	371	“ from drowning . . . . .	26
Affections of the ears . . . . .	152	“ from freezing . . . . .	28
“ “ eyes . . . . .	140	“ from hanging, choking, or suffocating by burdens and pressure . . . . .	28
“ “ face, lips, and jaws . . . . .	174	“ from noxious vapors . . . . .	29
“ “ head . . . . .	115	“ from a fall or blow . . . . .	29
“ “ mind . . . . .	110	“ of a new-born infant . . . . .	381
from fright and fear . . . . .	110	Appetite, morbid . . . . .	258
excessive joy . . . . .	110	“ want of . . . . .	257
grief . . . . .	110	Arachnoiditis . . . . .	124
home sickness . . . . .	110	Arthritis . . . . .	7
unhappy love . . . . .	110	Ascaris lumbricoides . . . . .	300
jealousy . . . . .	110	“ vermicularis . . . . .	300
mortification, insult . . . . .	110	Asiatic cholera . . . . .	325
contradiction and chagrin . . . . .	111	Asphyxia . . . . .	381
violent anger . . . . .	111	Asthma . . . . .	248
“ of the nose . . . . .	163	“ Millari . . . . .	389
“ “ penis . . . . .	349	“ of Millar . . . . .	254
“ “ stomach and bow- els . . . . .	257	“ of children . . . . .	389
“ “ teeth, gums, and mouth . . . . .	180	Backache . . . . .	9
“ “ testicles . . . . .	350	Bad taste in the mouth . . . . .	188
“ “ throat . . . . .	199	Bedridden . . . . .	72
“ “ windpipe and chest . . . . .	210	Biliousness . . . . .	267
After-birth, Adherence of the . . . .	371	Bilious colic . . . . .	279
After-pains . . . . .	372	Bilious rheumatic fever . . . . .	95
Agrypnia . . . . .	17	Birth, premature . . . . .	370
Ague in the breast . . . . .	375	“ regular . . . . .	371
Air and exercise . . . . .	442	Bites and stings of insects . . . .	36
Allopecia . . . . .	139	Bites of serpents . . . . .	37
Amaurosis . . . . .	149	Black vomit . . . . .	97, 270
Amblyopia . . . . .	149	Bladder, inflammation of the . . . .	342
Amygdalitis . . . . .	208	“ piles of the . . . . .	347
Anatomy and Physiology . . . . .	395, 397	Bleeding of the nose . . . . .	165
Anatomy of the bones . . . . .	400	Blindness . . . . .	149
Anger . . . . .	111	“ incipient . . . . .	149
Angina faucium . . . . .	199	“ sudden . . . . .	149
“ gangrenosa . . . . .	205	“ by night . . . . .	150
“ membranacea . . . . .	213	“ by day . . . . .	150
Anorexia . . . . .	257	Bloody Flux . . . . .	319
		Boils . . . . .	82

Bowels, affections of the	257	Colic with obstinate constipation	279
“ inflammation of the	291	“ caused by piles or hemorrhoids	279
“ looseness of the	313	“ inflammatory intestines	279
Breast	373	“ from worms	279
“ ague in the	375	“ from indigestion	279
“ gathered	375	“ from indignation, rage	279
Breath, offensive	188	“ from a chill	279
Bronchitis	213	“ from cold, damp weather	279
Bulimy	258	“ from bathing	279
Burns and scalds	34	“ from external injuries	280
Buzzing of the ears	157	“ in children	280
Calculus	348	“ in pregnant or lying-in women	280
Cancer	82	“ in hysterical women	280
“ of the face	179	“ at time of menses	280
“ of the nose	172	“ from low spirits	280
Canker of the mouth	190	“ during confinement	280
Carbuncles	82	“ griping	277
Cardialgia	272	“ of infants	382
Carditis	239	Common or ephemeral fever	87
Caries of the teeth	187	“ sore throat	199
Carious teeth	187	Concussions	34
Cataract	148	Confinement, duration of	373
Catarrhal cough	218	Congestio ad caput	115
“ dry	218	Congestion and stagnation of blood in the abdomen	286
“ loose	218	Congestion of blood to the head	115
Catarrh	171	“ of the chest	242
Cessation of the menses	360	Congestion or determination of blood to the head	115
Chafing of bedridden patients	72	from great joy	116
Chagrin	111	fright or fear	116
Change of life	360	anger, violent	116
Chancre	351	anger, suppressed	116
Chapped hands from working in water	72	a fall or blow	116
Chest, affections of	210	loss of blood	116
“ congestion of the	242	a cold	116
“ contusion of the	256	constipation	116
“ dropsy in the	254	drinking liquors	116
Chicken pox	63	sedentary habits	116
Chilblains	71	critical periods of girls	116
Children, treatment of	381	stoppage of menstruation	116
Chills and fever	104	dentition in children	116
Chlorosis	355	Congestive fever	94
Choking fits	389	Constipation	308
Cholera Asiatica	325	“ of infants	387
“ infantum	392	Consumption, galloping	255
“ morbus	323	“ pulmonary	255
Cholera	332	“ scrofulous	256
Chorea	21	Contagious typhus	94
Chronic Angina	205	Contradiction	111
“ dropsy of the chest	255	Contusion of the chest, by a fall or blow	256
“ dyspepsia	259	Convulsions or spasms of infants	385
“ eruptions	69	Corns	77
“ hoarseness	212	Coryza	167
“ inflammation of the eyes	145	Coryza of infants	382
Circulation of the blood	419	Costiveness	308
Clap	352	Cough	218
Clavi pedis	77	“ croupy	213, 216
Clothing and occupation	444	“ hooping	225
Cold in the head	167	Coughing up of blood	243
“ swelling	81	Courses	353
Colica	277		
Colic bilious	279		
“ flatulent or windy	279		
“ painter's or lead	279		

Coxagra - - - - -	11, 12	Enteritis - - - - -	291
Coxalgia - - - - -	11	Enuresis - - - - -	344
Coxalgia infantilis - - - - -	11	"    nocturna - - - - -	345
Cracked skin from cold - - - - -	73	"    spastica - - - - -	345
Critical period - - - - -	360	Epilepsia - - - - -	20
Cramp in the legs - - - - -	15	Epilepsy - - - - -	20
Croup - - - - -	213	Epistaxis - - - - -	165
Croupy cough - - - - -	213, 216	Eruptions and swelling of the ex- ternal ear - - - - -	152
Crustæ lactea - - - - -	75	Eruptions of the face - - - - -	174
Crying of infants - - - - -	382	Eruptions of the genital organs - - - - -	351
Cutaneous diseases - - - - -	47	Eruptions on the nose - - - - -	164
Deafness - - - - -	158	Eruptive fevers - - - - -	47
Debility during confinement - - - - -	380	Erysipelas bullosum - - - - -	49
Deficiency of milk - - - - -	376	"    erraticum - - - - -	50
Delirium tremens - - - - -	19	"    œdematodes - - - - -	52
Delivery, treatment after - - - - -	371	"    of face and lips - - - - -	174
Dentition - - - - -	386	"    of infants - - - - -	385
Determination of blood to the head - - - - -	115	"    of the nose - - - - -	163
Diabetes - - - - -	346	"    of the throat - - - - -	205
Diaphragm, anatomy and physi- ology of the - - - - -	410	"    on the scrotum - - - - -	351
Diaphragm, inflammation of the - - - - -	288	"    or St. Anthony's fire - - - - -	49
Diaphragmitis - - - - -	288	"    wandering - - - - -	50
Diarrhea - - - - -	313	Excoriations - - - - -	72
"    chronic - - - - -	315	"    of infants - - - - -	384
"    in lying-in females - - - - -	316	Exercise - - - - -	442
"    of infants - - - - -	388	External injuries - - - - -	30
Different forms of typhus - - - - -	91	"    of the abdomen - - - - -	339
Different modes of applying cold water - - - - -	453	Eyes, affections of - - - - -	140
Difficulty of speech - - - - -	197	Faceache - - - - -	175
Digestive organs - - - - -	412	Face, affections of - - - - -	174
Discharge from the womb during confinement - - - - -	379	Fainting - - - - -	24
Diseases caused by the pressure of worms - - - - -	303	Fall-baths - - - - -	457
Diseases consequent upon scarlet fever and scarlet rash - - - - -	61	Falling off of the hair - - - - -	139
Diseases during pregnancy - - - - -	365	during confinement - - - - -	380
"    of females - - - - -	353	Falling of the body - - - - -	335
"    of the heart - - - - -	239	"    of the eyelids - - - - -	151
Dislocations of joints - - - - -	31	"    of the palate - - - - -	209
Disposition to fester - - - - -	72	"    of the womb - - - - -	361
Disorders consequent upon measles - - - - -	55	False pleurisy - - - - -	237
Dizziness - - - - -	119	Far-sightedness - - - - -	150
Douche - - - - -	465	Fatigue - - - - -	23
Drop-bath - - - - -	464	Fear - - - - -	110
Dropsy of the chest - - - - -	254	Febris intermittens - - - - -	104
"    of the abdomen - - - - -	337	"    simplex - - - - -	87
"    of the brain - - - - -	388	Felons - - - - -	78
"    of the scrotum - - - - -	350	Females, diseases of - - - - -	353
Duration of confinement - - - - -	373	Festering - - - - -	72
Dysentery - - - - -	319	Fever and ague - - - - -	104
Dyspepsia - - - - -	258	Fever-cake - - - - -	299
"    recent - - - - -	259	"    and chills - - - - -	104
Dysury - - - - -	343	"    congestive - - - - -	94
Earache - - - - -	155	"    ephemeral or common - - - - -	87
Ears, affections of the - - - - -	152	"    inflammatory - - - - -	88
Ears, running of the - - - - -	159	"    intermittent - - - - -	104
Elements of the human organism - - - - -	400	"    irritative, inflammatory, torpid - - - - -	86
Elongation of the head of infants - - - - -	382	"    milk - - - - -	373
Encephalitis - - - - -	124	"    puerperal - - - - -	379
Enlarged glands - - - - -	81	"    remittent of infants - - - - -	390
Enteralgia - - - - -	277	"    rheumatic - - - - -	95
		Fevers - - - - -	84
		Fever, typhoid or nervous - - - - -	89
		"    yellow - - - - -	95

Flatulency . . . . .	265	Headache from overheating . . . . .	134
Flatulent or windy colic . . . . .	279	“ drinking ardent spirits . . . . .	134
Flooding . . . . .	359	“ loss of sleep . . . . .	134
Fluor albus . . . . .	363	“ external causes . . . . .	133
Food and drink . . . . .	429	“ gastric derangement . . . . .	133
Foot-baths . . . . .	463	“ nervous . . . . .	136
Foreign substances in the eye . . . . .	151	“ from rheumatism . . . . .	131
“ in the ear . . . . .	161	“ sick . . . . .	134
“ in the nose . . . . .	172	Head, affections of . . . . .	115
Fractures of bones . . . . .	30	Head-baths . . . . .	459
French measles . . . . .	56	Heart-burn . . . . .	265
Fright . . . . .	110	Heart, diseases of . . . . .	239
Frozen limbs . . . . .	39	“ inflammation of the . . . . .	239
Galloping consumption . . . . .	255	“ palpitation of the . . . . .	240
Gastralgia . . . . .	272	Heat of infants . . . . .	384
Gastric affections . . . . .	268	Heat spots . . . . .	384
“ bilious affections . . . . .	268	Hemorrhage of the lungs . . . . .	243
“ colic . . . . .	279	Hemorrhage with the urine . . . . .	348
“ derangement . . . . .	267	Hemorrhoids . . . . .	334
Gastritis . . . . .	289	Hemorrhoids vesica . . . . .	347
Gastroataxia . . . . .	268	Hepatitis . . . . .	295
Gastrodynia biliosa . . . . .	268	Hernia . . . . .	336
Gathered breast . . . . .	375	Herpes circinnatus . . . . .	73
General complaints during confinement . . . . .	380	Herpes . . . . .	73
General inflammatory fever . . . . .	88	Hip disease . . . . .	11
General plunging-baths . . . . .	458	Hives . . . . .	48
Genital organs, affections of the . . . . .	340	Hoarseness acute . . . . .	210
“ eruptions of the . . . . .	351	“ chronic . . . . .	212
Giddiness . . . . .	119	Hæmatemesis . . . . .	270
Glaucoma . . . . .	148	Hæmoptysis . . . . .	243
Gleet . . . . .	352	Home-sickness . . . . .	110
Glossitis . . . . .	189	Hooping-cough . . . . .	225
Goitre . . . . .	16	Humming of the ears . . . . .	157
Gonitis . . . . .	14	Humors in the ear . . . . .	152
Gonorrhea . . . . .	351	Hydrophobia . . . . .	448
Gout . . . . .	7	Hydrocele . . . . .	350
Gravel . . . . .	348	Hydrocephalus . . . . .	388
Green disease . . . . .	355	Hydrops saccatus . . . . .	338
Grief . . . . .	110	Hydrophobia . . . . .	39
Gripping colic . . . . .	277	Hydrothorax . . . . .	254
Gumbolls . . . . .	193	Hygiene . . . . .	428
Gums, affections of . . . . .	180	Hygiene and Hydrophobia . . . . .	395, 428
Hæmaturia . . . . .	348	Hypochondria . . . . .	113
Hæmaturia vesicalis . . . . .	349	Icterus . . . . .	298
Hanging jaw . . . . .	179	Ileus miserere . . . . .	292, 293
Hardness of hearing . . . . .	158	Impotence . . . . .	350
Headache . . . . .	128	Incontinence of urine . . . . .	344
“ from abuse of tobacco . . . . .	134	Incontinentia urinæ . . . . .	344
“ excess of bodily or mental labor . . . . .	134	Incubus . . . . .	17
“ grief . . . . .	134	Indigestion . . . . .	265
“ anger . . . . .	134	Indurated glands . . . . .	81
“ changeable weather . . . . .	134	Induration of the skin . . . . .	77, 78
“ from bathing . . . . .	134	Inflammation, abdominal . . . . .	289
“ suppressed eruptions . . . . .	134	“ in the abdomen after delivery, . . . . .	379
“ mechanical injuries . . . . .	134	“ of the bladder . . . . .	342
“ from catarrh and cold in the head . . . . .	130	“ of the bowels . . . . .	291
“ from congestion of blood to the head . . . . .	128	“ of the brain . . . . .	123
“ from constipation . . . . .	132	“ of the diaphragm . . . . .	288
“ from drinking coffee . . . . .	133	“ of the ear . . . . .	154
		“ chronic of the eyes . . . . .	145
		“ of the eyeball . . . . .	143
		“ of the margins of the lids . . . . .	143



Inflammation of the eyelid and the margins - - -	141	Lockjaw - - - - -	21, 22
“ of the eyelid - - -	141	Loin-ache - - - - -	9
“ of the eyes from abuse of mercury - - -	147	Looseness of the bowels - - -	313
“ of the eyeball - - -	143	Love, unhappy - - - - -	110
“ of the eyes from scrofula - - - - -	145	Lumbago - - - - -	9
“ of the eyes from syphilis or venereal disease -	146	Lungs, hemorrhage of the - -	243
“ of the eyes of infants -	383	Malignant boils - - - - -	82
“ of the face - - - - -	174	Malignant quinsy - - - - -	205
“ of the heart - - - - -	239	Mania a potu - - - - -	19
“ of the kidneys - - - -	340	Materia Medica - - - - -	395, 475
“ of the knee-joint - - -	14	Maw-worms - - - - -	300
“ of the liver - - - - -	295	Measles - - - - -	54
“ of the lungs - - - - -	229	Medicinal diseases - - - - -	45
“ of the lungs, or pneumonia in children -	233	Megrim - - - - -	126
“ of the nose - - - - -	163	Melæna - - - - -	270
“ of the palate - - - - -	193	Melancholia - - - - -	114
“ of the penis - - - - -	349	Memory, weakness of - - -	121
“ of the spine—spinal marrow - - - - -	9	Meningitis - - - - -	123
“ of the spleen - - - - -	299	Menses, cessation of the - -	360
“ of the stomach - - - -	289	Menses, suppression of the -	356
“ of the testicles - - - -	350	Menstrual colic - - - - -	280
“ of the tongue - - - - -	189	Menstruation - - - - -	353
“ of the tonsils - - - - -	208	“ of too long duration -	260
Inflammatory colic - - -	279	“ painful - - - - -	357
Influenza - - - - -	216	“ too early - - - - -	357
Insult - - - - -	110	“ too late and scanty -	358
Intermittent fever - - -	104	“ too copious - - - - -	359
Internal ear - - - - -	154	Mental exhaustion - - - -	23
Intertrigo - - - - -	72	Metritis - - - - -	379
Irritation of the skin - -	69	Mictus cruentus - - - - -	248
Ischias - - - - -	12	Milk bad, too thin, or repugnant to the child -	376
Ischury - - - - -	343	Milk-crust - - - - -	75
Itch - - - - -	74	Milk, deficiency of - - - -	376
Itching - - - - -	69	“ excessive flow of it -	376
“ of the anus - - - - -	335	“ fever - - - - -	373
“ in the ear - - - - -	152	“ leg - - - - -	180
“ of the nose - - - - -	165	“ secretion of - - - - -	373
“ on the private parts -	367	“ suppression of the -	376
Jaundice - - - - -	298	Mind, affections of the -	110
“ of infants - - - - -	384	Miscarriage - - - - -	268
Jaw-bone, swelling of - -	193	Morbid appetite - - - - -	258
Jaws, affections of - - -	174	Morbili - - - - -	54
Jealousy - - - - -	110	Morbus Coxarius - - - - -	11
Joy, excessive - - - - -	110	Mortification - - - - -	110
Kidneys, inflammation of the -	340	Mouth, affections of - - -	180
Labor-pains - - - - -	371	Mouth, canker of - - - -	190
Laryngitis catarrhalis - -	213	“ ulceration of - - - -	190
Lead-colic - - - - -	278	Mumps - - - - -	153
Lethargy - - - - -	23	Muscles - - - - -	408
Leucorrhœa - - - - -	363	Myelitis - - - - -	10
Lienteria - - - - -	318	Myopia - - - - -	150
Lips, affections of - - -	174	Nausea - - - - -	269
Liver complaint - - - - -	296	Nausea and vomiting during pregnancy -	366
“ grown - - - - -	288	nancy - - - - -	340
“ inflammation of the -	295	Nephritis - - - - -	219
Local applications - - - -	467	Nervous and spasmodic cough -	280
Local baths - - - - -	459	“ colic of women - - -	126
Lochia - - - - -	379	“ headache - - - - -	89
		“ or typhoid fever - - -	423
		“ system, anatomy of the -	48
		Nettle-rash - - - - -	175
		Neuralgia in the face - - -	136
		“ in the head - - - - -	

Neuralgia of the stomach - - -	272	Podagra - - - - -	9
Nightmare - - - - -	17	Poisoned wounds - - - -	76
Nipples - - - - -	373	Poisons - - - - -	40
" sore - - - - -	374	" animal - - - - -	43
Nose, affections of - - - -	163	" vegetable - - - -	44
" swelling of - - - -	163	" mineral - - - - -	44
Notalgia - - - - -	9	Pollution - - - - -	350
Nursing women, sore mouth of	378	Polypus of the bladder - -	347
Obstruction of the menses - -	354	" of the ear - - - -	161
" of the nose in infants	382	" of the nose - - - -	172
Occupation - - - - -	444	Pregnancy - - - - -	364
Odontalgia - - - - -	180	Premature birth - - - -	370
Offensive breath - - - - -	188	Prolapsus ani - - - - -	335
Onania - - - - -	350	Prolapsus uteri - - - -	361
Ophthalmia - - - - -	143	Prosopalgia - - - - -	175
Otalgia - - - - -	155	Prurigo - - - - -	71
Otitis - - - - -	154	" on the scrotum - - -	351
Otorrhœa - - - - -	159	Pruritus - - - - -	69, 367
Overheating - - - - -	23	Pseudo-typhus, or bilious-rheu-	
Ozæna antri Highmori - - -	171	matic fever - - - - -	95
Ozæna narium - - - - -	171	Psoitis - - - - -	14
Pain in the abdomen after delivery	379	Ptyalism - - - - -	195
Pain in the hip - - - - -	11	Puerperal fever - - - -	379
Pain in the stomach - - - -	276	Pulmonary consumption - -	255
Pains in the small of the back		Pyrosis - - - - -	265
and loins - - - - -	9	Putrid sore throat - - - -	205
Painter's or lead colic - - -	279	Quinsy - - - - -	199
Palate, elongation, swelling of	209	" malignant - - - - -	205
Palate, falling of the - - -	209	Ranula - - - - -	197
Palpitation cordis - - - - -	240	Rash - - - - -	47
Palsy - - - - -	18	Raucitas - - - - -	210
Palpitation of the heart - - -	240	Rawness of the skin in infants	284
Panaris - - - - -	78	Real pleurisy - - - - -	256
Paralysis - - - - -	18	Red gum - - - - -	384
" of the eyelids - - - -	151	Red spots on the nose - - -	164
" of the face and jaw - -	179	Regular birth - - - - -	371
" of the tongue - - - -	196	Remarks on pneumonia - -	225
Partial or half-baths - - - -	456	Remittent fever - - - - -	104
Parotitis - - - - -	153	Remittent fevers of infants -	390
Part second - - - - -	395	Respiratory and Circulatory or-	
Parturition - - - - -	371	gans, anatomy of the - -	418
Penis, affections of the - - -	349	Restlessness of infants - - -	383
Penis, inflammation of the - -	349	Retention of urine during con-	
Phthisis laryngea - - - - -	213	finement - - - - -	380
" pulmonalis - - - - -	255	Rhagades - - - - -	72
Physiology of the bones - - -	407	Rheumatism - - - - -	3, 6
Piles - - - - -	334	" acute - - - - -	4
Piles of the bladder - - - -	347	" inflammatory - - -	4
Pimples on the face - - - -	175	" chronic - - - - -	6
Pin-worms - - - - -	300	" shifting or wander-	
Placenta, adherence of the - -	371	ing - - - - -	5
Plethora pectoris - - - - -	242	Ringworm - - - - -	73
Pleurisy - - - - -	236	Ringworm of the scalp - - -	76
" real - - - - -	236	Running of the ears - - - -	159
" false - - - - -	237	Rupture - - - - -	336
Pleuritis - - - - -	236	Salivation - - - - -	195
" muscularis - - - - -	226	Scabies - - - - -	74
" serosa - - - - -	226	Scald-head - - - - -	76
Pneumonia - - - - -	229	Scalds - - - - -	14
" in infants and children -	233	Scarlet-fever - - - - -	58
" in old people - - - -	234	" " malignant - - - -	60
" notha - - - - -	234	Scarlet-rash - - - - -	57
" remarks on - - - - -	235	Sciatica - - - - -	11, 12
Pneumorrhagia - - - - -	243	Schirrus - - - - -	82

Scorbutus - - - -	193	Structure of the human organism	398
Scrofulous consumption - - -	256	Struma - - - - -	16
Scrofulous sore eyes - - -	145	Stuttering - - - - -	197
Scrotum, dropsy of the - - -	150	St. Vitus's dance - - - -	21
" erysipelas on the - - -	351	Stye on the eyelid - - - -	147
" prurigo on the - - -	351	Stymatosis - - - - -	349
" tetter on the - - -	151	Summer-complaint - - - -	392
Scurvey - - - - -	193	Sunstroke - - - - -	127
Sea-sickness - - - - -	271	Suppression of the catarrh, or cold	
Secretion of milk - - - - -	37	in the head - - - - -	169
Seminal weakness - - - - -	350	Suppression of the menses - -	356
Ship fever - - - - -	95	Suppression of the milk - - -	376
Short-sightedness - - - - -	150	Sweating - - - - -	470
Shower baths - - - - -	455	Sweating feet - - - - -	15
Sick headache - - - - -	134	Swelling of the breasts in infants	382
Sitting baths - - - - -	461	" of the cheek - - - - -	174
Sitz baths - - - - -	461	" of the ear - - - - -	152
Skin, anatomy of the - - - -	425	" of glands - - - - -	81
Sleep - - - - -	440	" of the jaw-bone - - - -	193
Sleeplessness - - - - -	17	" of the nose - - - - -	63
Sleeplessness during confinement	380	" of the palate - - - - -	269
Sleeplessness of infants - - -	383	" of the tonsils - - - - -	208
Small pox - - - - -	64	" under the tongue - - - -	197
Snake-bitten - - - - -	37	Swooning - - - - -	24
Snuffles - - - - -	382	Syncope - - - - -	24
Somnolency - - - - -	23	Synocha - - - - -	88
Sore mouth of infants - - - -	383	Syphilis - - - - -	351
" " of nursing women - - -	378	Tania - - - - -	300
" nipples - - - - -	374	Tape worm - - - - -	300
" throat - - - - -	199	Tastes, different - - - - -	189
" throat, chronic - - - - -	205	Taste, loss of - - - - -	189
Spasms in the stomach - - - -	272	Teeth, affections of - - - -	180
Spasms of infants - - - - -	386	Teeth, anatomy of the - - - -	411
Specks on the eyes - - - - -	146	Teething - - - - -	386
Spitting of blood - - - - -	243	Testicles, affections of the -	350
Spleen, enlargement of the - -	299	Tetanus - - - - -	21
Spleen, indurated - - - - -	299	Tetter - - - - -	73
Spleen, inflammation of the -	299	" dry - - - - -	73
Splenitis - - - - -	299	" running - - - - -	73
Spontaneous limping - - - - -	1, 12	" bleeding - - - - -	73
Spots on the face - - - - -	178	" ulcerating - - - - -	73
Spots on the face during pregnancy	368	" on the scrotum - - - - -	351
Spots on the nose, black red -	164	Throat, affections of - - - -	199
Sprains and concussions - - -	34	Throat, erysipelas of - - - -	205
Squinting - - - - -	150	Thrush - - - - -	383
Stagnation of blood in the abdomen	286	Tic douloureux - - - - -	175
Stammering - - - - -	197	Tinea capitis - - - - -	76
St. Anthony's fire - - - - -	49	Tongue, inflammation of - - -	189
State of the bowels after delivery	377	" swelling of - - - - -	189
Stings of insects - - - - -	36	" swelling under the - - -	197
Stitch in the side - - - - -	237	" ulceration of - - - - -	190
Stomacace - - - - -	190	Tonsils, inflammation of - - -	208
Stomach-ache - - - - -	277	Tonsils, swelling of - - - -	208
Stomach, affections of the - -	157	Toothache - - - - -	180
" cold in the - - - - -	273	Treatment after delivery - - -	371
" inflammation of the - - -	289	Treatment of children - - - -	381
" neuralgia of the - - - -	272	Trismus - - - - -	21, 22
" sour - - - - -	265	Tubercular consumption - - -	255
" spasms and pain in the - -	272	Tumors - - - - -	81
" weak - - - - -	258	Tussis convulsiva - - - - -	225
Stone in the bladder - - - - -	348	Typhoid pneumonia - - - - -	231
Strabismus - - - - -	150	Typhus - - - - -	89
Strangury - - - - -	343	" versatile - - - - -	91
		" stupid - - - - -	93

Typhus putrid - - - -	93	Vomiting bile - - - -	270
" contagious - - - -	94	" of mucus - - - -	270
Ulcerated sore throat - - -	205	" of water - - - -	270
Ulceration in the nose - - -	170	" sour - - - -	270
Ulceration of the windpipe -	213	Want of appetite - - - -	257
Ulcers on the ear - - - -	152	Warts - - - -	78
Ulcers - - - -	79	Warts on the face - - - -	178
Ulcers on the eyeballs - - -	146	Warts on the nose - - - -	164
Umschläge - - - -	467	Water-brash - - - -	265
Urinary organs, affections of the	340	Weakness of memory - - - -	121
Urinary system, anatomy of the	417	Weakness of sight - - - -	149
Urine, bloody - - - -	348	Weakness, seminal - - - -	350
Urine, incontinence of the - -	344	Weak stomach - - - -	258
Vaccination - - - -	391	Weaning - - - -	376
Varicose veins - - - -	367	Weeping or watery eyes - - -	148
Variola - - - -	64	Wet bandages - - - -	467
Varioloid, or modified small pox	68	Wet linen sheets - - - -	469
Veins varicose - - - -	367	Wetting the bed - - - -	345
Verrucæ - - - -	78	Whites - - - -	363
Vertigo - - - -	119	White swelling - - - -	11
" from disordered stomach	119	Whitlows on the fingers - - -	78
" from irritation of brain	119	Wind-colic - - - -	278
" from suppressed ulcers		Windpipe, affections of - - -	210
and eruptions - - -	119	Windpipe, ulceration of the -	213
" from riding in a carriage	120	Womb, falling of the - - -	361
" from congestion of blood		Worm-colic - - - -	305
to the brain - - -	120	Worm-fever - - - -	305
Vomiting - - - -	269	Worms - - - -	300
" after a fall - - - -	270	Wounds and bruises - - - -	32
" of blood - - - -	270	Yellow fever - - - -	95

HOMŒOPATHIC  
BOOKS AND MEDICINES.

---

H. F. DAVIS, M. D.

No. 23, Main Street, Cincinnati.

*Keeps constantly on hand and for sale, an assortment of*  
**Homœopathic Books and Medicine Chests, Cases, and Single Remedies.**

AMONG THE BOOKS, ARE

**Buchner and Gruner.** New Homœopathic Pharmacopœia and Posology ; or the Mode of preparing Homœopathic Medicines, and the Doses. Compiled and translated from the German Works of BUCHNER and GRUNER, and the French Work of Jahr. With original contributions by Chas. J. Hempel, M. D. Cloth, \$2.

**Hempel, Chas. J.** The Homœopathic Domestic Physician. By C. J. Hempel, M. D. Second Edition, revised and considerably enlarged by the Author. Cloth, 50 cents.

**Jahr's New Manual of Homœopathic Practice**, edited, with Annotations, by A. Gerald Hull, M. D. This is the fourth American edition of a very celebrated work, by the eminent Homœopathic Professor, Jahr, and it is considered the best practical compendium of this extraordinary science that has yet been composed.

Complete Repertory, one volume, bound, \$3.

**Jahr's New Manual**, originally published under the name of Symptomen-Codex. (Digest of Symptoms.) This work is intended to facilitate a comparison of the parallel symptoms of the various homœopathic agents, thereby enabling the practitioner to discover the characteristic symptoms of each drug, and to determine with ease and correctness what remedy is most homœopathic to the existing group of symptoms. Translated, with important and extensive additions, from various sources, by Charles Julius Hempel, M. D.; assisted by James M. Quin, M. D.; with revisions and chemical notes by John F. Gray, M. D.; contributions by Drs. A. Gerald Hull and George W. Cook, M. D., of New York, and Drs. C. Hering, J. Jeans, C. Neidhard, W. Williamson, and J. Kitchen, of Philadelphia. With a preface by Constantine Hering, M. D. 2 volumes, bound, 1848. \$11 00.

HOMŒOPATHIC BOOKS.

---

**Laurie, Dr. J**, *Homœopathic Domestic Medicine*, with the Treatment and Diseases of Females, Infants, Children and Adults. Fifth American edition, much enlarged with additions by A. Gerald Hull, M. D. 1849. Bound, \$1 50.

**Laurie, Dr. J.**, *Elements of Homœopathic Practice of Physic*. An Appendix to Laurie's Domestic, containing, also, all the diseases of Urinary and Genital Organs. Bound, 1849. \$1 25.

**Jahr's Clinical Guide, or Pocket Repertory**. Translated from the German, by C. J. Hempel, M. D., bound (just published.) \$1 50.

**Hahnemann's Organon of Homœopathic Medicine**. Third American edition, with improvements and additions from the last German edition, and Dr. C. Hering's introductory remarks, 1848. Bound, \$1 00.

**Rau's Organon, of the specific healing art of Homœopathy**. Translated by Charles J. Hempel, M. D. 1848. \$1 25.

**Hahnemann, Dr.**, *Materia Medica Pura*. Translated and edited by Charles Julius Hempel, M. D. 4 vols. 1846. \$6.

**E. Stapf's Additions to the Materia Medica Pura**. Translated by C. J. Hempel, M. D. \$1 50.

**Hahnemann, Dr. S.**, *The Chronic Diseases, their specific Nature and Homœopathic Treatment*. Translated and edited by Charles J. Hempel, M. D., with a Preface, by Constantine Hering, M. D., Philadelphia. 8vo. 5 vols. Bound, 1845. \$7 00.

**Transactions of the American Institute of Homœopathy**. 1846. Bound. \$1 50.

**Becker, M. D.** On Consumption. Translated from the German. 1884.  
" " 38 cents.  
" " On Diseases of the Eye. 1848. 38 cts.  
" " On Constipation. 1848. 38 cts.  
" " On Dentition. 1848. 38 cts.

**Hempel's Bœnninghausen** for Homœopathic Physicians; to be used at the bedside of the patient, and in studying the *Materia Medica Pura*. Most complete edition, including the Concordance of Homœopathic Remedies. Translated and adapted to the use of the American Profession, by C. J. Hempel, M. D. 1847. \$1 50.

**Hartmann's Acute and Chronic Diseases**, by Charles J. Hempel, M. D. 4 vols. \$6.

**Jahr, G. H. G., M. D.** *Short Elementary Treatise upon Homœopathia* and the manner of its Practice; with some of the most important effects of ten of the principal Homœopathic Remedies, for the use of all honest men who desire to convince themselves, by experiment, of the truth of the doctrine. Second French edition, corrected and enlarged. Translated by Edward Bayard, M. D. Bound. 38 cts.



**Bœnninghausen's** Essay on the Homœopathic Treatment of Intermittent Fevers. Translated and edited by Charles Julius Hempel, M. D. 1845. 38 cts.

**A Treatise on the use of Arnica.** in cases of Contusions, Wounds, Strains, Sprains, Lacerations of the Solids, Concussions, Paralysis, Rheumatism, Soreness of the Nipples, etc. By Charles Julius Hempel, M. D. 1845. 19 cts.

**Homœopathic Cookery.** Second edition. Designed chiefly for the use of such persons as are under homœopathic treatment. 50 cts.

**Rueckert's Therapeutics;** or Successful Homœopathic Cures, collected from the best Homœopathic periodicals. By C. J. Hempel, M. D. One large 8vo. vol. \$3.

**The Homœopathic Examiner.** By Drs. Gray and Hempel. 2 volumes. New Series. 1846 and 1847. Bound in two volumes. \$6.

**Chepmell's Homœopathic Domestic,** with additions and improvements, by S. B. Barlow, M. D. 1849. 50 cts.

**E. Stapf's Additions** to the Materia Medica Pura. Translated by C. J. Hempel, M. D. \$1 50.

**C. Hering's Domestic Physician.** Fourth American edition, revised, with additions from the Author's manuscript of the sixth German edition. The part relating to the Diseases of Females and Children, by Walter Williamson, M. D. 1848. \$2 00.

**Ruoff's Repertory of Homœopathic Medicine,** nosologically arranged. Translated from the German, by A. H. Okie, M. D., translator of Hartmann's Remedies. Second American edition, with additions and improvements, by G. Humphrey, M. D., etc. 1844. Bound. \$1 50.

**Epps, Dr. J., Domestic Homœopathy;** or Rules for the Domestic Treatment of the Maladies of Infants, Children, and Adults, etc. Third American from the fourth London edition. Edited and enlarged by George W. Cook, M. D. 1848. Bound. 75 cts.

**W. Williamson, M. D.,** Diseases of Females and Children. 38 cents.

**William Henderson, M. D.** Homœopathic Practice. 1846. 50 cents.

**Forbes, M. D.** Homœopathy, Allopathy, and Young Physic. 1846. 19 cents.

**Wm. Henderson, M. D.** Letter to John Forbes. 1846. 19 cents.

☞ The above three books, bound in one volume, \$1 00.

**The Family Guide** to the Administration of Homœopathic Remedies. Third edition, after the second London edition, with additions. Retail price, 25 cents.

HOMŒOPATHIC BOOKS.

---

**On Eclecticism in Medicine;** or a Critical Review of the Leading Medical Doctrines. An inaugural thesis, presented at the New York University, on the first of March, 1845. By Charles J. Hempel, M. D. 25 cents.

**Reasons** why Homœopathy should receive an Impartial Investigation from the Medical Profession and the Public. By B. F. Bowers, M. D. 18 cents.


**Defence of Hahnemann and his Doctrines,** including an Exposure of Dr. Alexander Wood's "Homœopathy Unmasked." London, 1844. 50 cents.

**Sherrill's Manual of Homœopathy.** 37½ cents.

**F. A. Guenther's** New Manual of Homœopathic Veterinary Medicine; or, the Homœopathic Treatment of the Horse, the Ox, the Sheep, the Dog, and other Domestic Animals. \$1 25.







NATIONAL LIBRARY OF MEDICINE



NLM 02521333 5